

Association of Aberrant Promoter Methylation Changes in the Suppressor of Cytokine Signaling 3 (*SOCS3*) Gene with Susceptibility to Crohn's Disease

Golshid Sanati ¹, Davood Jafari ^{2,3}, Mehrdad Noruzinia ⁴, Naser Ebrahimi Daryani ⁵,
Mohammad Ahmadvand ⁴, Shahram Teimourian ⁶, and Nima Rezaei ^{1,7,8*}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Immunology, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran
3. Immunogenetics Research Network (IgReN), Universal Scientific Education and Research Network (USERN), Zanjan, Iran
4. Department of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran
5. Department of Internal Medicine, Division of Gastroenterology, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran
6. Department of Genetics, School of Medicine, Iran University of Medical Sciences, Tehran, Iran
7. Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran
8. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Boston, MA, USA

Abstract

Background: Growing evidence supports that changes in the methylation state of Inflammatory Bowel Disease (IBD)-associated genes could significantly alter levels of gene expression, potentially contributing to disease onset and progression. We supposed that alterations in DNA methylation status at promoter region within the suppressor of cytokine signaling 3 (*SOCS3*) gene in intestinal tissues may be involved in the susceptibility to Crohn's Disease (CD).

Methods: DNA methylation status in the promoter region of the human *SOCS3* gene of intestinal tissues from 15 patients with CD and 15 age- and sex-matched healthy controls were profiled using the real-time Quantitative Multiplex Methylation Specific PCR (QM-MSP) assay.

Results: Based on methylation assay data profiling, we found that patients with CD showed a higher degree of methylation of the *SOCS3* gene promoter region than did the healthy controls (unmethylated DNA in CD *vs.* healthy controls; 0.00048 ± 0.0011 *vs.* 0.07 ± 0.142 , $p < 0.000$).

Conclusion: The data presented here demonstrate that aberrant methylation of the CpG islands within promoter regions of *SOCS3* gene in colonic mucosa of CD was associated with mucosal inflammatory status, providing insights into the involvement of methylation could contribute to the initiation of the inflammatory process and development of CD.

Avicenna J Med Biotech 2022; 14(2): 165-169

Keywords: Crohn's disease, DNA methylation, Epigenetics, Immune regulation, *SOCS3*

Introduction

Crohn's Disease (CD) is a major subtype of Inflammatory Bowel Disease (IBD) and could potentially affect whole Gastrointestinal (GI) tract ¹. CD is histopathologically characterized by the presence of persistent inflammation of the terminal ileum and colon dominates ^{2,3}. CD appears to be multifactorial and polygenic and the precise etiology of the disease is not

clear. However, it is assumed that the onset and perpetuation of the disease is caused by a complex interplay between susceptibility genes, the immune system and environmental factors ^{3,4}.

It is becoming apparent that epigenetic factors can have a significant contribution in the pathogenesis of disease ⁵. Epigenetic can be defined as inherited chang-

es in gene expression that is not involved in DNA sequencing itself and is influenced by environment⁶. Genome Wide Association Studies (GWAS) have reported more than 160 gene loci related to IBD, but only 13.6% of CD patients and 7.5% of ulcerative colitis patients are accounted for this IBD-associated genetic variants so the role of non-genetic factors such as epigenetic regulations have been bolded recently⁷. Emerging evidence suggests that epigenetic modifications, including changes in the methylation state of IBD-associated genes could significantly alter levels of gene expression, potentially contributing to disease onset and progression⁸.

Jack/stat pathway is cytokine-modulated intracellular signaling pathway which can regulate the immune system development and hematopoiesis potentially through inflammatory cytokines such as Tumor Necrosis Factor (TNF)- α , Interferon (IFN)- γ and Interleukin (IL)-6^{2,9}. STAT family of transcription factors including STAT3 have been demonstrated to have a pivotal role in transmitting inflammatory cytokine signals to the nucleus¹⁰. The suppressors of cytokine signaling (SOCS) 3 protein as a member of SOCS family is the key physiological regulators of cytokine-mediated activation of STAT3 by inhibiting the catalytic activity of JACKs¹¹. In the recent decades, SOCS3 has been one of the major interesting research subjects due to changes in the expression of SOCS3 contribute to the development of a variety of autoimmune diseases including IBDs^{2,9}.

It has been shown that in IBD patients, SOCS3 expression is considerably decreased in comparison with healthy volunteers¹²⁻¹⁴, however up to our knowledge there is no study that investigate the SOCS3 promoter region methylation as a fundamental epigenetic mechanism. To appreciate the role of DNA methylation as a mechanism underlying differences in SOCS3 expression level and consequently in the regulation of the protein, we checked promoter methylation status. The primary goal of this study was to determine the alterations in DNA methylation status at promoter region within *SOCS3* gene in intestinal tissues to clarify whether it is hyper or hypo-methylated in CD in comparison to normal population.

Materials and Methods

Patients and tissue samples

We studied extracted DNA from colon biopsy specimens of 15 CD patients (8 females, 7 males), who underwent total proctocolectomy at the gastroenterology clinics of Kasra and Laleh Hospitals in Tehran, Iran, between May 2018 and July 2019. Colonic biopsies from 15 age- and sex-matched patients (8 females, 7 males) without abnormalities at colonoscopy and histories of gastrointestinal disease served as healthy controls. Diagnosis of CD was determined according to the usual clinical criteria, endoscopic and pathohistological

criteria. The tissue samples were frozen at -80 °C immediately. The study protocol was approved by the Committees on the Ethics of Human Research of Tehran University of Medical Sciences.

DNA isolation and Bisulfite conversion system

Genomic DNA was isolated from frozen colon tissues using the High Pure PCR Template Preparation Kit (Roche) according to the manufacturer's protocol. DNA concentration and quality were measured using the NanoDrop ND-1000 spectrophotometer. Genomic DNA was modified by sodium bisulfite to convert unmethylated cytosines to uracil using the available MethylEdge™ Bisulfite Conversion System (Promega, Madison, WI) as per manufacturer's instructions. Bisulfite-modified DNA specimens were aliquoted and stored at -20 °C.

Methylation-specific PCR

The SOCS3 promoter and gene sequences, obtained from the GenBank database of National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov>), were used to design primers for methylation studies. The MethySYBR procedure requires two sequential steps of PCR reactions. MethySYBR primers for *SOCS3* gene were designed, including external forward primer (EXT-F1: 5'-GTAGGGAGGTGACGAG GTAGG-3'), external reverse primer (EXT-R: 5'-GGT TGTTCCGGGTTATTTTGT -3'), contain no CpG dinucleotides. The external primers was used in the PCR reaction amplified total DNA templates of the target gene regardless of their methylation status. This served as a reference control to normalize the amount of methylated target alleles detected by the methylation specific primer pair between samples. The initial amplification of PCR reaction was performed in 25 μ l of PCR reaction mixture composed of 1 μ l bisulphite-treated genomic DNA, SOCS3 external primers 1 μ l. The DNA was denatured at 95 °C for 5 min, followed by 30 cycles at 94 °C for 30 s, 56 °C for 30 s, and 72 °C for 30 s, with a final extension at 72 °C for 5 min.

For designing the nested methylation-specific primers, three CpG sites of SOCS3 promoter were targeted in the primer sequence. The methylation profile of the promoter CpG islands were determined based on UCSC database. Methylation-Specific PCR (MSP)-specific primers were designed by Methprimer used to amplify methylated and unmethylated DNA sequences as appropriate. Methyl forward primer (FM: 5'-GGA GATTTTAGGTTTTTCGGAATATTTTC-3'), Methyl reverse primer (RM: 5'-CCCCCGAACTACCTAAAC GCCG-3'). The methylation status of bisulfite-converted DNA was determined using a fluorescence-based, real-time polymerase chain reaction (Real time-PCR) assay. For the second-step real-time PCR reactions, 1 μ l of the diluted PCR product was used in each PCR reaction. The bisulfite treated DNA was PCR amplified in a 10 μ l reaction volume containing containing 5 μ l SYBR® Green Master Mix (Ampliqon,

Denmark), 0.25 μ l of each of the methylated primers, 3.5 μ l DDW and 1 μ l of bisulfite-treated DNA. Quantitative MSP was performed with a 7500 Real-Time PCR System (Applied Biosystems, Foster City, CA). No untreated template controls were included in each run as negative controls.

For data presentation, fully converted methylated human DNA was included in each run to serve as the 100% methylated reference for calculating the relative methylation percentages of DNA samples based on the relative $2^{-\Delta\Delta CT}$ quantitation approach.

Statistical analysis

DNA methylation data were expressed as mean \pm SD. The possible statistical differences in CpG island methylation levels between two groups were determined using the Mann-Whitney test. Chi-square statistic was used to discover if there is a relationship between two categorical variables. To calculate the measure of association, the Odds Ratio (OR) and 95% confidence interval were considered. All p-values were 2-tailed and considered significant when less than 0.05. All statistical calculations were made using SPSS 21.0 version.

Results

Because some genes are shown to be methylated with aging and sex, age- and sex-matched healthy controls were studied. The study consisted of 15 CD patients, with a mean age of 46 \pm 17.2 years (range: 17-71 years), and included 7 (46.7%) males and 8 (53.3%) females. Furthermore, fifteen were healthy controls (8 females, 7 males), with a mean age of 46 \pm 15.38 years (range: 28-78 years).

The use of SYBR green dye during quantitative PCR enables melting curve analysis of target amplicons to determine the methylation status of CpG sites on *SOCS3* gene. The resulting melting curve for *SOCS3* gene is shown in figure 1.

The promoter methylation status of *SOCS3* gene in colonic mucosa specimens of all 15 CD patients and 15 paired healthy controls were compared (Figure 2). Methylation assay data profiling showed evidence of differential promoter methylation levels between patients with CD and healthy controls in which all three regions of the *SOCS3* gene were more highly methylated in the active inflamed mucosa (unmethylated DNA=0.00048 \pm 0.0011) than in mucosa of healthy controls (unmethylated DNA=0.07 \pm 0.142) (p <0.000, Figure 2). Qualitative methylation data analysis of each individual showed a significant difference in the frequency of promoter methylation status between rectal inflammatory mucosa (12/15, 80%), and paired normal terminal ileum (2/15, 13.3%). The relative risk of disease indicated that individuals who had an aberrant promoter methylation profiling, were at high risk of the development of CD (p =0.009, OR=26.00, CI=22.85-337.72).




Figure 1. The figure shows the fluorescent melting peaks for the promoter CpG island of the *SOCS3* gene. DNA melt curve analysis of changes in CpG methylation status was analyzed based on amplification of unmethylated bisulfite-treated DNA from colorectal biopsies of CD cases.




Figure 2. Comparison of changes in CpG island methylation status at the *SOCS3* gene promoter region in colorectal tissue specimens of patients with ulcerative colitis and healthy control. Error bars mean \pm SD (p <0.05).

Discussion

Chronic inflammation including IBD is tightly related to increased levels of DNA methylation^{15,16}. Although there have been studies reporting increased DNA methylation in colonic mucosa of IBD disease, the most recently published evidence in this field show

the association between patterns of DNA methylation of colon mucosa and UC-associated dysplasia or carcinoma incidence without considering the mucosal inflammatory status¹⁵. In the present study, we have examined and compared the alterations in CpG island methylation status at the *SOCS3* gene promoter region in colorectal tissue specimens on the disease susceptibility. Our data demonstrated that methylation levels of *SOCS3* gene-associated CpG islands were substantially higher in colorectal mucosa of patients with CD than in mucosa from healthy controls.

Despite the importance of the currently known genetic predispositions of multiple genes in the pathogenesis of IBD, an increasing number of observations reveal that abnormal epigenetic changes are considered as or even more important for IBD pathogenesis than genetic predisposition. Twin studies reveal that epigenetic modifications play an important etiologic role so that monozygotic twin on colectomy samples, one of who had CD methylation differences were observed in several genes that reveals the importance of the role of epigenetics. This observation suggests that prenatal developmental programming of colonic mucosal and immunological maturation imprints/programs postnatal microbiota shifts in the intestinal mucosa DNA methylation at different sites in a gene has quite different outcome in transcriptional activity^{2,17}.

Cytokines are secreted proteins central for coordination of the initiation, maintenance, and termination of all types of immune responses, including host responses to infection, inflammation. A tight control of cytokine release and of responses to cytokines is required for the defense against infections, the prevention of infection-associated immunopathology, and the correct development of immune cell populations. Several cytokines and growth factors utilize multiple intracellular signaling pathways such as the JAK-STAT pathway to transmit their information into the cell nucleus¹⁸.

As mentioned, the SOCS proteins negatively regulate JAK-STAT signal transduction by forming a negative feedback loop that modulates cytokine-induced signaling through different mechanisms^{19,20}. The expression of *SOCS3* gene is induced by phosphorylated STAT3 in response to cytokines that signal through gp130-related cytokine receptors, which in turn leads to attenuate or terminate the activation of STAT3 through targeting the receptor associated JAK kinases for degradation⁹.

IL-6/STAT3 was an important pathways for the progress of IBD²¹. Expression of *SOCS3*, which is a direct target gene of STAT3 and mediates suppression of IL-6 signaling through the ubiquitination and degradation of signaling intermediates²². Mice with a deletion of *SOCS3* in hematopoietic cells (*Socs3^{fl/fl} vav cre*) have been shown to develop a severe inflammatory disease during adult life in response to pro-inflammatory cytokines including IL-6¹⁸. Accordingly, ade-

noviral-delivered *SOCS3* reduced joint inflammation in mice with arthritis *via* inhibition of IL-6 signaling¹⁸.

SOCS3 protein is an important regulator of cytokine signaling and consequently the immune response. Since *SOCS3* regulates the function of a variety of cytokine associated with IBD, it plays a critical role in the development of these diseases². Studies in different mouse models have proven the critical importance of *SOCS3* in restraining inflammation and allowing optimal levels of protective immune responses against infections¹⁸.

SOCS3 promoter methylation is a major regulator of *SOCS3* expression in the colonic epithelial compartment. Increased methylation in inflamed CD may result from the upregulation of DNMT1 expression. It thus seems that *SOCS3* promoter methylation is an important mechanism controlling IL-6-dependent *SOCS3* expression in colon cancer cultures²². Data suggest that IL-6-dependent DNMT1 induction mediates *SOCS3* downregulation, in turn releasing IL-6 signaling towards STAT3 from its negative regulation. A prediction from this hypothesis is that inhibition of DNMT1 should restore IL-6-dependent induction of *SOCS3* and counteract methylation of *SOCS3* promoter. Concomitant with the loss of *SOCS3*, STAT3 activation increased, providing a rational explanation as to why *SOCS3* loss is related to an increased propensity to develop UC-CRC. Thus, we proposed that the loss of *SOCS3* expression represents a critical event, permissive for the development of UC-CRC, but the molecular mechanism as to why chronic inflammation would provoke the loss of *SOCS3* expression remained obscure at most²².

Interestingly, when analyzing methylation status of *SOCS3* gene, we observed very low levels of unmethylated DNA in both groups. The maximum inhibitory activity of *SOCS1* and *SOCS3* proteins toward the activation of STAT1 can be achieved by very low levels of *SOCS* protein expression, suggesting that *SOCS* proteins have a high affinity toward Jaks *in vivo*²³.

Conclusion

In summary, this case-control study detected new profiles of CpG hypermethylation of *SOCS3* gene that were associated with the susceptibility to CD development. Our study revealed a high negative correlation between promoter methylation and basal *SOCS3* levels. We report that the hypermethylation status of *SOCS3* in colorectal tissue specimens of CD patients is significantly higher compared to healthy controls. Our findings suggest that the CpG hypermethylation at promoter region of gene might influence *SOCS3* gene expression and are fundamentally related to the etiology of UC. Our data provide an important insight into the STAT/SOCS axis by representing that epigenetic modifications in the *SOCS3* gene regulation are the basis for hyperactivity of the inflammation-related IL6-

STAT3 in UC patients. Further studies using *in vivo* and *ex vivo* systems will be needed to confirm our findings and even propose that how this knowledge may open novel avenues for the rational treatment of IBD.

Acknowledgement

This study was supported by a grant from Tehran University of Medical Sciences (93-02-30-25932).

Ethics Approval

The study protocol was approved by the Ethics Committees of Tehran University of Medicine science. Written informed consent was obtained from all individuals before sampling.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Bojic D, Markovic S. Terminal ileitis is not always Crohn's disease. *Ann Gastroenterol* 2011;24(4):271-5.
- Li Y, de Haar C, Peppelenbosch MP, van der Woude CJ. SOCS3 in immune regulation of inflammatory bowel disease and inflammatory bowel disease-related cancer. *Cytokine Growth Factor Rev* 2012;23(3):127-38.
- Xavier R, Podolsky D. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448(7152):427-34.
- Haag LM, Siegmund B. Intestinal Microbiota and the Innate Immune System-A Crosstalk in Crohn's Disease Pathogenesis. *Front Immunol* 2014;6:489.
- Lin Z, Hegarty J, Cappel J, Yu W, Chen X, Faber P, et al. Identification of disease-associated DNA methylation in intestinal tissues from patients with inflammatory bowel disease. *Clinical Genet* 2011;80(1):59-67.
- Wilson AG. Epigenetic regulation of gene expression in the inflammatory response and relevance to common diseases. *J Periodontol* 2008;79(8S):1514-9.
- Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012;491(7422):119-24.
- Ventham NT, Kennedy NA, Nimmo ER, Satsangi J. Beyond gene discovery in inflammatory bowel disease: the emerging role of epigenetics. *Gastroenterology* 2013;145(2):293-308.
- Egwuagu CE. STAT3 in CD4+ T helper cell differentiation and inflammatory diseases. *Cytokine* 2009;47(3):149-56.
- Yoshimura A. Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci* 2006;97(6):439-47.
- Carow B, Rottenberg ME. SOCS3, a major regulator of infection and inflammation. *Front Immunol* 2014;5:58.
- Cheng X, Zhang X, Su J, Zhang Y, Zhou W, Zhou J, et al. miR-19b downregulates intestinal SOCS3 to reduce intestinal inflammation in Crohn's disease. *Sci Rep* 2015;5:10397.
- Li Y, de Haar C, Chen M, Deuring J, Gerrits MM, Smits R, et al. Disease-related expression of the IL6/STAT3/SOCS3 signalling pathway in ulcerative colitis and ulcerative colitis-related carcinogenesis. *Gut* 2010;59(2):227-35.
- Xu AT, Li Y, Zhao D, Shen J, Xu XT, Qiao YQ, et al. High suppressor of cytokine signaling-3 expression impairs STAT3-dependent protective effects of interleukin-22 in ulcerative colitis in remission. *Inflamm Bowel Dis* 2015;21(2):241-50.
- Saito S, Kato J, Hiraoka S, Horii J, Suzuki H, Higashi R, et al. DNA methylation of colon mucosa in ulcerative colitis patients: correlation with inflammatory status. *Inflamm Bowel Dis* 2011;17(9):1955-65.
- Low D, Mizoguchi A, Mizoguchi E. DNA methylation in inflammatory bowel disease and beyond. *World J Gastroenterol* 2013;19(32):5238-49.
- Kellermayer R. Epigenetics and the developmental origins of inflammatory bowel diseases. *Can J Gastroenterol* 2012;26(12):909-15.
- Rottenberg ME, Carow B. SOCS3, a major regulator of infection and inflammation. *Front Immunol* 2014;5:58.
- Krebs DL, Hilton DJ. SOCS proteins: negative regulators of cytokine signaling. *Stem Cells* 2001;19(5):378-87.
- Li Y. The role of SOCS3 signaling in ulcerative colitis and ulcerative colitis-related carcinogenesis [thesis]. [Rotterdam, the Netherlands]. 2012. 154 p.
- Chen YY, Ma ZB, Xu HY, Shi LJ, Li DY, Sun LY, et al. IL-6/STAT3/SOCS3 signaling pathway playing a regulatory role in ulcerative colitis carcinogenesis. *Int J Clin Exp Med* 2015;8(8):12009-17.
- Li Y, Deuring J, Peppelenbosch MP, Kuipers EJ, de Haar C, van der Woude CJ. IL-6-induced DNMT1 activity mediates SOCS3 promoter hypermethylation in ulcerative colitis-related colorectal cancer. *Carcinogenesis* 2012;33(10):1889-96.
- Song MM, Shuai K. The suppressor of cytokine signaling (SOCS) 1 and SOCS3 but not SOCS2 proteins inhibit interferon-mediated antiviral and antiproliferative activities. *J Biol Chem* 1998;273(52):35056-62.