



Investigation of Expression Profile of Placenta-specific 1 (PLAC1) in Acute Myeloid and Lymphoid Leukemias

Parastou Gholami¹, Hossein Asgarian-Omran², Marjan Yaghmaie³, Jafar Mahmudian⁴, Shirin Kianersi⁵, Sina Salari⁵, Ehsan Zabolli⁶, Mahmood Jeddi-Tehrani⁴, Amir-Hassan Zarnani^{7,8} and Mahdi Shabani^{1*}

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

3. Hematology-Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran

4. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

5. HSCT Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

6. Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran

7. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

8. Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Abstract

Background: Placenta-specific 1 (PLAC1) is one of the cancer-testis-placenta antigens that has no expression in normal tissue except placenta trophoblast and testicular germ cells, but is overexpressed in a variety of solid tumors. There is a lack of studies on the expression of PLAC1 in leukemia. We investigated expression of *PLAC1* in Acute Myeloid Leukemia (AML) and Acute Lymphoblastic Leukemia (ALL).

Methods: In this study, we investigated expression pattern of *PLAC1* gene in peripheral blood and bone marrow mononuclear cells of newly-diagnosed patients with AML (n=31) and ALL (n=31) using quantitative real-time PCR. Normal subjects (n=17) were considered as control. The PLAC1 protein expression in the samples were also detected using western blotting.

Results: Our data demonstrated that *PLAC1* transcripts had 2.7 and 2.9 fold-change increase in AML and ALL, respectively, compared to normal samples. *PLAC1* transcript expression was totally negative in all studied normal subjects. Level of *PLAC1* mRNA expression in ALL statistically increased compared to normal samples (p=0.038). However, relative mRNA expression of *PLAC1* in AML was not significant in comparison to normal subjects (p=0.848). Furthermore, relative mRNA expression of *PLAC1* in AML subtypes was not statistically significant (p=0.756). *PLAC1* gene expression showed no difference in demographical clinical and para-clinical parameters. Western blotting confirmed expression of PLAC1 in the ALL and AML samples.

Conclusion: Considering *PLAC1* expression profile in acute leukemia, PLAC1 could be a potential marker in leukemia which needs complementary studies in the future.

Keywords: Acute lymphoblastic leukemia, Acute Myeloid Leukemia, Biomarker, Expression profile, Leukemia, PLAC1

To cite this article: Gholami P, Asgarian-Omran H, Yaghmaie M, Mahmudian J, Kianersi Sh, Salari S, et al. Investigation of Expression Profile of Placenta-specific 1 (PLAC1) in Acute Myeloid and Lymphoid Leukemias. Avicenna J Med Biotech 2023;15(3):167-172.

* Corresponding author:
Mahdi Shabani, Ph.D.,
Department of Immunology,
School of Medicine, Shahid
Beheshti University of Medical
Sciences, Tehran, Iran
Tel/Fax: +98 21 22439970
E-mail:
msshabani@sbmu.ac.ir,
msshabani@yahoo.com
Received: 24 Sept 2022
Accepted: 26 Apr 2023

Introduction

Acute Myeloid Leukemia (AML) is a heterogeneous disorder of hematopoietic stem cells that mainly affects middle-aged people (mean age of diagnosis is 69 years). AML is seen with uncontrolled clonal proliferation of myeloid progenitor cells leading to defects in normal hematopoiesis¹. It is the most common acute leukemia that accounting for about 80% of cases in this

group. In the United States, the incidence of AML is 3-5 cases per 100000 population and over 20000 per year². Acute Lymphoblastic Leukemia (ALL) is a clonal malignancy characterized by the accumulation of lymphocyte blasts that do not differentiate into adult lymphocytes³. ALL is the most common cancer among children and the most important cause of cancer death

in people under 20 years of age⁴. The disease appears to begin with genetic mutations in hematopoietic stem cells. These mutations accumulate in stem cells to eventually lead to three functional disorders, including increased proliferation, cessation of blast differentiation, and decreased apoptotic death⁵. There is no tumor marker with a wide expression pattern in the acute leukemia that could be applicable in general diagnosis or targeting leukemic cells, thus scientists are now searching for such biomarker.

Placenta-specific 1 (*PLAC1*) is an X-linked (Xq26) gene with a conserved structure in mice and humans that is involved in placental evolution^{6,7}. It has six exons located in an area of about 200 Kbp, with only one protein-containing transcript in humans (NP-068568.1, ENS T00000359237)⁷. *PLAC1* has two promoters, P1 and P2, which are about 100 Kbp apart, but all 639 bp that contain the gene coding sequence are in the exon 6. Therefore, the *PLAC1* protein is the product of the last and the largest exon of *PLAC1* gene⁸. Keratinocyte Growth Factor (KGF) increases the expression of *PLAC1* transcript by 2-3 times. This factor plays a role in regulating placental function and differentiation in mammals⁹. In healthy individuals, the expression of *PLAC1* is mainly limited to placental tissue and its lower levels of mRNA expression have been reported in the testes^{10,11} and cerebellum^{11,12}.

In humans, *PLAC1* protein is found in placenta with relatively constant expression during pregnancy^{9,13}. The *PLAC1* protein is found in the apical syncytiotrophoblast^{7,13} and has been shown to bind to F-actin at the membrane and apical syncytial trophoblast levels. Studies have shown that *PLAC1* protein is associated with membranes and is predominantly found in microvillous brush border membranes⁷. *In vitro* studies have shown that silencing *PLAC1* gene using siRNA in MCF-7 and BT-544 cell lines reduces motility, migration, and invasion, as well cell cycle arrest leading to a reduction of about 80-90% in cell proliferation. Decreased *PLAC1* expression was associated with decreased in cyclin D1 expression and AKT kinase phosphorylation¹⁴. Importantly, *PLAC1* expression in both mRNA and protein levels were detected in a variety of malignancies including hepatocellular carcinoma¹⁵, ovarian cancer¹⁶, prostate cancer¹⁷, melanoma¹⁸, gastric adenocarcinoma¹⁹ and breast cancer²⁰. There is a lack of study on the expression of *PLAC1* in leukemia such as AML and ALL. Based on popularity of this type of cancers, here we investigated the expression pattern of *PLAC1* in the newly-diagnosed patients with AML and ALL.

Materials and Methods

Samples collection

Sixty two EDTA-anticoagulated Peripheral Blood (PB) and Bone Marrow (BM) samples were obtained from AML (BM=13, and PB=18) and ALL (BM=16, and PB=15) patients that had referred to Blood and On-

colony Department of Imam Khomeini Hospital (Mazandaran University of Medical Sciences), Hematology Research Center of Oncology and Stem Cell Transplantation (Tehran University of Medical Sciences) or Taleghani Hospital (Shahid Beheshti University of Medical Sciences) during October 2018 to September 2019 (Table 1). Blood samples were also taken from 17 normal healthy individuals (the mean age: 35.6 years; SD: 12.9 and 64.7% male) for the control group. A consent letter was taken from all patients and control group. This study was approved by the Ethical Committee of Shahid Beheshti University of Medical Sciences (IR.SBMU.MSP.REC.1398.328).

RNA Extraction and cDNA preparation

Mononuclear cells were immediately isolated by Ficoll density centrifugation, and washed twice with Phosphate Buffered Saline (PBS, pH=7.2, 0.15 M). The separated cells were lysed in Trizol reagent (Yekta Tajhiz Azma, Iran). Total RNA extraction was performed following the manufacturer's instructions. Single-stranded cDNA preparation was performed from 1 µg total RNA of each sample using reverse transcriptase (Thermo Scientific, Lithuania) and random primer (Cybergene, Sweden).

Quantitative RT-PCR

We used the SYBR green (Yekta Tajhiz Azma, Iran) to determine *PLAC1* expression levels. The specific primers were used as follows: *PLAC1* sense, 5'-G TGAGCACAAAGCCACATTTC-3' and antisense, 5'-GCAGCCAATCAGATAATGAACC-3'; *GAPDH* sense, 5'-GAAGGTGAAGGTCGGAGTC-3' and antisense, 5'-GAAGATGGTGTATGGGATTTC-3' (Bioneer,

Table 1. Demographic and laboratory findings of studied acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL)

Characteristic	Results	
	ALL group	AML group
Age, mean+SD	20.8+16.7	48.7+19.1
Male	19 (73.1%)	17 (54.8%)
Female	7 (26.9%)	14 (45.2%)
Sample	BM: 16 PB: 15	BM: 13 PB: 18
WBC ($\times 10^3/\mu\text{l}$), mean+SD	46.2+62.6	38.6+55.6
PLT ($\times 10^3/\mu\text{l}$), mean+SD	347+23.6	545+32.9
Hb (gr/dl), mean+SD	8.8+2.0	8.1+1.3
PB Blast (%)	26.3	36.4
BM Blast (%)	67.3	58.2
Subtype (FAB/ immunophenotype)	B-ALL: 26 (83.9%) T-ALL: 5 (16.1%)	M0: 2 (6.5%) M1: 7 (22.6%) M2: 8 (25.8%) M3: 3 (9.7%) M4: 5 (16.1%) M5: 4 (12.9%) M6: 1 (3.2%)

Korea). The quantitative PCR (qPCR) assays were conducted on a Corbett rotor gene 6000 (QIAGEN, Germany). qPCR mainly included enzyme activation (15 min at 95°C), followed by denaturation (15 s, 95°C), annealing, extension and fluorescence acquiring (60 s, 60°C) for 40 cycles. The amplicons' size of *PLAC1* and *GAPDH* genes were 118 bp and 226 bp, respectively. qPCR results were analyzed and expressed relative to threshold cycle (CT) values and then converted to fold-change. The *PLAC1* gene expression was normalized using *GAPDH* as house-keeping internal control and relative gene expression data presented as fold-change compared to normal group based on $2^{-\Delta\Delta CT}$ method.

Western blotting

Separated mononuclear cells (1×10^6) were lysed in 50 μ l sample buffer containing 150 mM Tris Ph=6.8, 1.2% SDS, 200 mM 2-ME, 0.2% bromophenol blue and 33% glycerol on ice. The lysates were boiled for 10 min and then centrifuged at $17700 \times g$ for 20 min at 4°C. Thirty μ l of cell lysates and 35 ng recombinant human *PLAC1* (rh*PLAC1*) (produced in Dr. Zarnani's lab) were run on a 14% SDS-PAGE gel at 100 V for 2 hrs and then transferred to PVDF membrane at 200 mA for one hr. The existence of protein bands was visualized by Ponceau S staining. The membrane was blocked with PBS-0.1% Tween 20 (PBS-T) containing 5% skimmed milk solution overnight without shaking at 4°C. The membrane was then immunoprobed with rabbit anti-rh*PLAC1* antibody (produced in Dr. Zarnani's lab) at 2 μ g/ml in PBS-T containing 1% skimmed milk solution for 1.5 hr with shaking at RT followed by goat anti-rabbit IgG-HRP (1:6000) (Bio-Rad, Hercules, CA, USA) in PBS-T containing 1% skimmed milk solution for one hr with shaking at RT. After extensive washing, membrane was developed with ECL. Anti- β -actin rabbit monoclonal antibody (Sigma, St. Louis, USA) was used for visualization of β -actin after re-probing of the membrane.

Statistical analysis

IBM SPSS statistics 26 (SPSS Inc., Chicago, IL, USA), and the REST software was used for the statistical analysis. The results of data analysis are reported based on p value and in the form of tables and graphs. In a quantitative study, the Kolmogrov-Smirnov test was used to determine whether the data distribution was normal or abnormal. Because the distribution of our study data was abnormal, non-parametric statistical tests such as Kruskal Wallis and Man-Whitney U tests were used to compare the means between several groups and the two groups, respectively, and to determine the correlation between quantitative parameters performed by Spearman Correlation Test. The results with $p < 0.05$ were considered statistically significant.

Results

Relative expression analysis of *PLAC1* by real time PCR

Specific amplification of the *GAPDH* and *PLAC1* genes were confirmed by agarose gel electrophoresis. The amplicons' size of *PLAC1* and *GAPDH* genes were compatible with the expected sizes as 118 bp and 226 bp, respectively (Figures 1A and B). The amplicon and melting curves of the *GAPDH* and *PLAC1* was also confirmed with their specific amplification (Figure 2). Furthermore, *PLAC1* PCR product of one AML sample was confirmed by Sanger sequencing (ABI 3500 genetic analyzer, Thermofisher Scientific, Waltham, MA, USA; data not shown). In addition, the efficiency of *PLAC1* (96%) and *GAPDH* (98%) amplification was validated using MDA-MB-231 cells, as positive control.

Our data demonstrated that *PLAC1* transcripts had 2.9 fold-change increase in ALL compared to normal samples. Further analysis pointed out that the level of *PLAC1* mRNA expression in both PB and BM mononuclear cells of ALL patients was significantly higher compared to normal samples ($p=0.038$) (Figure 3). Interestingly, *PLAC1* transcript expression compared to normal control group was statistically significant just in B-ALL but not in T-ALL subtype ($p=0.029$) (Figure 3). Furthermore, *PLAC1* expression was significantly different just in BM and PB samples of B-ALL ($p=0.04$), but not in T-ALL.

Furthermore, *PLAC1* transcripts had 2.7 fold-change increase in AML compared to normal samples. But despite of ALL samples, relative mRNA expression of *PLAC1* in AML samples was not significantly different compared to normal samples ($p=0.848$) (Figure 3). Furthermore, relative mRNA expression of *PLAC1* in AML subtypes was not statistically significant ($p=0.756$). In comparison to normal PB samples, *PLAC1* expression was significantly higher in PB ($p=0.011$)




Figure 1. Agarose gel electrophoresis of real-time PCR products of *PLAC1* (A) and *GAPDH* (B).

Lanes 1: marker 100 bp, 2: AML12-BM, 3: AML7-PB, 4: ALL1-BM, 5: ALL23-PB, 6: Normal 4.

Placental-specific 1 Expression Profile in ALL and AML




Figure 2. The real time PCR amplification curve and melting curve of *PLAC1* and *GAPDH*.




Figure 3. Comparison of *PLAC1* mRNA expression in leukemic and normal groups.

and BM blasts ($p=0.029$) of AML and ALL patients. Analysis of *PLAC1* expression showed that *PLAC1* had no difference in demographical clinical and para-clinical subgroups.

PLAC1 protein expression by Western blotting

PLAC1 expression at protein level in 5 samples was checked in immunoblotting using a rabbit polyclonal antibody specific to human *PLAC1* recombinant protein (produced in Dr. Zarnani's laboratory). Based on the immunoblotting findings, the expression of *PLAC1* protein was consistent with the expression of *PLAC1*




Figure 4. Representative data on *PLAC1* protein expression pattern in AML and ALL patients. *PLAC1* and β -actin proteins bands laid on 21 and 42 *kDa* in SDS-PAGE 10%. Lanes 1: AML21-PB, 2: AML23-PB, 3: ALL25-PB, 4: ALL27-PB, 5: ALL19-BM, 6: human placenta, 7: *PLAC1* recombinant protein.

gene, and samples with positive *PLAC1* transcript had *PLAC1* protein expression, too. Representative Western blot results with anti-*PLAC1* and β -actin antibodies are presented in figure 4, as expected *PLAC1* and β -actin showed 24 and 42 *kDa* sizes, respectively.

Discussion

Due to the lack of access to a suitable biomarker that can be used as a diagnostic marker in all AML or ALL patients, we investigated the expression of *PLAC1* transcript in these leukemias. In this study the expression of *PLAC1* transcript is detected in ALL and AML to a significant extent, which requires more examination on a larger number of samples. In addition to quantitative analysis of *PLAC1* real-time PCR, we arbitrarily considered samples with $CT \leq 34$ as positive and the cases with $CT \geq 35$ as negative for *PLAC1* mRNA

expression. According to this criterion, *PLAC1* expression was positive in 67.7% of ALL and 48.3% of AML patients. In this regard, *PLAC1* gene expression was negative in all included normal samples. Furthermore, *PLAC1* expression was detected in 69.2 and 60% of B-ALL and T-ALL, respectively. Although, the sample size of different subgroups of B-ALL was very small, interestingly Pre-B ALL subtype had the highest *PLAC1* expression (6 out of 9; 65.2%). This finding was also found in AML-M1 with 71.4% (5 out of 7 cases) as the highest frequency of *PLAC1* positive samples among AML FAB subtypes. In addition, the expression of *PLAC1* transcript was examined in normal healthy individuals and none of the normal individuals had *PLAC1* transcript in PB samples, which is a confirmation of the lack of expression of *PLAC1* molecule in normal tissues except placenta and testis¹⁰.

Identification of genes that are expressed in tumors and have no expression in normal tissues is a prerequisite for the study of cancer immunotherapy. Identification of these genes will also help advance diagnosis and treatment, including tumor immunotherapy. *PLAC1* is a molecule of the cancer testis family of antigens that is expressed in placental and testicular tissue but is not expressed in normal tissues, according to published studies¹⁰. Xue-Yuan Dong *et al* showed that *PLAC1* is highly expressed in hepatocellular carcinoma tissue and anti-*PLAC1* antibodies have been found in the serum of patients which indicated its immunogenicity¹⁵. Nana E. Tchabo *et al* showed that *PLAC1* and DPPA2 were among the testicular cancer antigens (CTA) that were expressed on 21 and 31% ovarian epithelial cancer, respectively¹⁶.

In another study Roya Ghods *et al* examined the pattern of expression of the *PLAC1* gene in advanced prostate adenocarcinoma. They reported that in prostate cancer, as the expression of *PLAC1* increased, the expression of (prostate specific antigen) PSA decreased¹⁸. In another study by Roya Ghods *et al* on melanoma, 100% of *PLAC1* expression on melanoma tissues was observed¹⁹. Fangfang Liu *et al* proposed *PLAC1* as a potential biomarker in the diagnosis and treatment of gastric adenocarcinoma²⁰. In a study on breast cancer by Hongyan Yuan *et al*, *PLAC1* was detected as a biomarker in the serum of 40-60% of breast cancer patients²¹. Yongfei Li and colleagues showed that following the interaction of *PLAC1* with Furin, the intracellular domain of Notch1 transcription factor, called NICD, is activated and inhibits PTEN suppressor tumor expression, a pathway leads to tumor progression, proliferation and metastasis. Therefore, high expression of *PLAC1* in breast cancer is associated with poor prognosis, migration and tumor invasion²². Qiongshu Li and colleagues investigated the targeting of *PLAC1* in breast cancer using genetically engineered T cells against *PLAC1* (CAR-T-*PLAC1* cells)²³. All reported studies indicated the importance of *PLAC1* in biology of tumors and the potential applications of this mole-

cule in diagnosis and therapy of *PLAC1* positive cancers.

Conclusion

In conclusion, due to the lack of *PLAC1* expression in normal samples and its expression in ALL and AML patient samples, *PLAC1* can be considered as a tumor-specific antigen or tumor-dependent antigen, which seems to be expressed in some ALL and AML patients. Additional studies are suggested in more samples in each leukemia group and with better homogeneity, for stronger results.

Statement of Ethics

The Shahid Beheshti University of Medical Sciences research committee approved the study (IR.SBMU.MSP.REC.1398.328). All experiments or tests performed involving human subjects were conducted according to the institute ethical standards and the Declaration of Helsinki. Samples of 1-2 ml of aspirated bone marrow and 3-5 ml of peripheral blood were collected in vials containing EDTA after obtaining the signature of the patient or the patient's guardian on an informed consent form as approved by Shahid Beheshti University of Medical Sciences Research Committee.

Data Availability Statement

The dataset used and analyzed during the current study are available from the corresponding author upon reasonable request.

Acknowledgement

This work is financially supported by Shahid Beheshti University of Medical Sciences (grant number: 15082). This work is part of Parastou Gholami's thesis. We appreciate Ms. Sedighe Vafaei for letting us have her valuable experience on setting up Western blot.

Conflict of Interest

All authors declare that there is no conflict of interest.

References

1. Sharma P, Pollyea DA. Shutting Down Acute Myeloid Leukemia and Myelodysplastic Syndrome with BCL-2 Family Protein Inhibition. *Curr Hematol Malig Rep* 2018;13(4):256-64.
2. De Kouchkovsky I, Abdul-Hay M. 'Acute myeloid leukemia: a comprehensive review and 2016 update'. *Blood Cancer J* 2016 1;6(7):e441.
3. Cobaleda C, Sanchez-Garcia I. B-cell acute lymphoblastic leukaemia: towards understanding its cellular origin. *Bioessays* 2009;31(6):600-9.
4. Hunger SP, Mullighan CG. Acute Lymphoblastic Leukemia in Children. *N Engl J Med* 2015;373(16):1541-52.
5. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000 Jan 7;100(1):57-70.

Placental-specific 1 Expression Profile in ALL and AML

6. Cocchia M, Huber R, Pantano S, Chen EY, Ma P, Forabosco A, et al. PLAC1, an Xq26 gene with placenta-specific expression. *Genomics* 2000 ;68(3):305-12.
7. Fant M, Barerra-Saldana H, Dubinsky W, Poindexter B, Bick R. The PLAC1 protein localizes to membranous compartments in the apical region of the syncytiotrophoblast. *Mol Reprod Dev* 2007;74(7):922-9.
8. Chen Y, Moradin A, Schlessinger D, Nagaraja R. RXRalpha and LXR activate two promoters in placenta- and tumor-specific expression of PLAC1. *Placenta* 2011; 32(11):877-84.
9. Fant M, Weisoly DL, Cocchia M, Huber R, Khan S, Lunt T, et al. PLAC1, a trophoblast-specific gene, is expressed throughout pregnancy in the human placenta and modulated by keratinocyte growth factor. *Mol Reprod Dev* 2002;63(4):430-6.
10. Silva WA, Jr., Gnjatich S, Ritter E, Chua R, Cohen T, Hsu M, et al. PLAC1, a trophoblast-specific cell surface protein, is expressed in a range of human tumors and elicits spontaneous antibody responses. *Cancer Immun* 2007;7: 18.
11. Fant M, Farina A, Nagaraja R, Schlessinger D. PLAC1 (Placenta-specific 1): a novel, X-linked gene with roles in reproductive and cancer biology. *Prenat Diagn* 2010; 30(6):497-502.
12. Wang X, Baddoo MC, Yin Q. The placental specific gene, PLAC1, is induced by the Epstein-Barr virus and is expressed in human tumor cells. *Virology* 2014;11:107.
13. Massabba E, Parveen S, Weisoly DL, Nelson DM, Smith SD, Fant M. PLAC1 expression increases during trophoblast differentiation: evidence for regulatory interactions with the fibroblast growth factor-7 (FGF-7) axis. *Mol Reprod Dev* 2005;71(3):299-304.
14. Koslowski M, Sahin U, Mitnacht-Kraus R, Seitz G, Huber C, Tureci O. A placenta-specific gene ectopically activated in many human cancers is essentially involved in malignant cell processes. *Cancer Res* 2007;67(19): 9528-34.
15. Dong XY, Peng JR, Ye YJ, Chen HS, Zhang LJ, Pang XW, et al. Plac1 is a tumor-specific antigen capable of eliciting spontaneous antibody responses in human cancer patients. *Int J Cancer* 2008;122(9):2038-43.
16. Tchabo NE, Mhaweche-Fauceglia P, Caballero OL, Villella J, Beck AF, Miliotto AJ, et al. Expression and serum immunoreactivity of developmentally restricted differentiation antigens in epithelial ovarian cancer. *Cancer Immun* 2009;9:6.
17. Ghods R, Ghahremani MH, Madjd Z, Asgari M, Abolhasani M, Tavasoli S, et al. High placenta-specific 1/low prostate-specific antigen expression pattern in high-grade prostate adenocarcinoma. *Cancer Immunol Immunother* 2014;63(12):1319-27.
18. Ghods R, Ghahremani MH, Darzi M, Mahmoudi AR, Yeganeh O, Bayat AA, et al. Immunohistochemical characterization of novel murine monoclonal antibodies against human placenta-specific 1. *Biotechnol Appl Biochem* 2014;61(3):363-9.
19. Mahmoudi AR, Ghods R, Rakhshan A, Madjd Z, Bolouri MR, Mahmoudian J, et al. Discovery of a potential biomarker for immunotherapy of melanoma: PLAC1 as an emerging target. *Immunopharmacology Immunotoxicol* 2020;42(6):604-13.
20. Liu F, Shen D, Kang X, Zhang C, Song Q. New tumour antigen PLAC1/CP1, a potentially useful prognostic marker and immunotherapy target for gastric adenocarcinoma. *J Clin Pathol* 2015;68(11):913-6.
21. Yuan H, Chen V, Boisvert M, Isaacs C, Glazer RI. PLAC1 as a serum biomarker for breast cancer. *PLoS One* 2018;13(2):e0192106.
22. Li Y, Chu J, Li J, Feng W, Yang F, Wang Y, et al. Cancer/testis antigen-Plac1 promotes invasion and metastasis of breast cancer through Furin/NICD/PTEN signaling pathway. *Mol Oncol* 2018 ;12(8):1233-48.
23. Li Q, Liu M, Wu M, Zhou X, Wang S, Hu Y, et al. PLAC1-specific TCR-engineered T cells mediate antigen-specific antitumor effects in breast cancer. *Oncol Lett* 2018;15(4):5924-32.