



Antitumor Activities of Green Tea by Up-regulation of miR-181a Expression in LNCaP Cells Using 3D Cell Culture Model

Fatemeh Safari ^{1*}, Narjes Rayat Azad ¹, Ali Alizadeh Ezdiny ², Safura Pakizehkar ³,
Zeinab Khazaei Koozpar ⁴, and Najmeh Ranji ²

1. Department of Biology, Faculty of Science, University of Guilan, Rasht, Iran

2. Department of Biology, Faculty of Basic Sciences, Rasht Branch, Islamic Azad University, Rasht, Iran

3. Cellular and Molecular Endocrine Research Center (CMERC), Research Institute for Endocrine Science, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Cell and Molecular Biology, Faculty of Biological Sciences, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

Abstract

Background: Prostate Cancer (PCa) is the major reason for the high mortality rates among men worldwide. In fact, current therapeutic approaches are not successful. It appears that discovering more effective methods considering several parameters such as availability, low cost, and no toxicity to normal cells is one of the biggest challenges for interested researchers. Green tea (extracted from the plant *Camellia sinensis*) with high level of polyphenolic compounds and as the most globally consumed beverage has attracted considerable interest. MicroRNAs (or miRNAs) were considered as novel tools in cancer therapy which modulate various biological events in cell by regulation of gene expression. The aim of the current study was to evaluate the antitumor activity of green tea in LNCaP cells through up-regulation of miR-181a expression.

Methods: First, LNCaP cells were cultured and by using quantitative real time PCR (qRT-PCR) and western blot methods, the expression levels of Bax and BCL2 were analyzed. Next, a 3D cell culture model was applied to evaluate the expression of miRNA-181a in LNCaP cells.

Results: It was shown that green tea induced cellular apoptosis. The high number of apoptotic nuclei was also shown by using DAPI staining. The inhibition of tumor growth was revealed by analyzing the size and number of spheroids. Also, up-regulation of miR-181a expression in LNCaP cells was revealed after treatment with green tea.

Conclusion: Our results are helpful to design antitumor regimens based on consumption of green tea through up-regulation of miRNA-181a expression and induction of apoptosis.

Avicenna J Med Biotech 2022; 14(1): 89-94

Keyword: Apoptosis, Cell culture techniques, Green tea, Proto-oncogene proteins c-bcl-2, Up-regulation

* **Corresponding author:**
Fatemeh Safari, Ph.D.,
Department of Biology, Faculty
of Science, University of Guilan,
Rasht, Iran
Tel: +98 133 3333647
E-mail:
fsafari@guilan.ac.ir
Received: 21 Apr 2021
Accepted: 7 Jun 2021

Introduction

Cancer is a multifactorial disease with high death rates in human. Among different cancer types, Prostate Cancer (PCa) is one of the leading causes of death among men worldwide ¹. Hormone therapy, surgery, radiation, and chemotherapy are current approaches for treatment of PCa patients. Resistance to anticancer drugs and their side effects are well recognized as major reasons for the failure of cancer treatment and therefore, the identification of new tools and specific platforms with the lowest side effects especially from

natural products is urgently needed. In herbal therapy, it seems that plants may serve as potent cost effective chemotherapeutic agents with less toxicity to normal mammalian tissues ². In this regard, protective effects of green tea as an herbal medicinal product have attracted considerable interest. Green tea is one of the most popular beverages worldwide which is produced from the leaves of the *Camellia sinensis* plant. Green tea contains a group of polyphenolic compounds including four main catechins of (–)-epicatechin (EC),

(-)-epigallocatechin (EGC), (-)-epicatechin-3-gallate (ECG), and (-)-epigallocatechin-3-gallate (EGCG)^{3,4}. Healthy effects of green tea on many human diseases have been documented⁵⁻⁸. Moreover, the antitumor activities of green tea on different types of human cancers have been discovered⁹⁻¹².

MicroRNAs (miRNAs) are new promising targets in cancer therapy. miRNAs are single-stranded non-coding RNAs involved in various cellular pathways, negatively or positively regulating their target genes^{13,14}. Therefore, by using miRNAs, new therapeutic approaches and diagnostic tools to treat cancer were employed¹⁵. In PCa, the implication of several miRNAs in modulating a variety of biological processes including proliferation, migration, invasion, and apoptosis was shown¹⁶⁻¹⁹. miR-181a belongs to the miR-181 family. It was reported that miR-181a can act as an oncogenic miRNA or tumor suppressor miRNA and thereby, the exact role of miR-181a in tumorigenesis is not clearly understood²⁰⁻²⁷. It seems that the role of miR-181a in tumorigenesis is tumor-specific. On the other hand, several reports also showed the inhibitory effects of purified green tea catechins on the cancer cells by the dysregulation of miRNAs²⁸⁻³¹.

In this study, an attempt was made to study the anti-cancer effects of green tea against LNCaP human prostate cancer cells through changing miR-181a expression in 3D cell culture model. By using DAPI staining, quantitative real-time PCR (qRT-PCR) and western blot, the induction of apoptosis was shown. Moreover, the antitumor effects of green tea on LNCaP cells through miR-181a were evaluated in 3D cell culture model. Based on our results, it was found that green tea may be a potential candidate and a novel natural product in the treatment of prostate cancer patients through up-regulation of miR-181a expression.

Materials and Methods

Sample preparation and extraction

The fresh leaves of related plants were collected from Lahijan city, Guilan province, Iran in March 2018. It was considered as *Camellia sinensis* in the Department of Pharmacognosy at Tehran University of Medical Sciences and the herbarium was registered as THE-6561. Plant extraction was performed as previously described¹². Briefly, the fresh leaves were dried at room temperature and darkness for two weeks, and the powder was obtained. The powder was soaked in 50% pure ethanol (Merck, Germany) +50% water for three days. The extract was filtered and stored at -20°C.

Cell line and culture

Lymph node carcinoma of the prostate (LNCaP) was provided from Pasture Institute (Tehran, Iran), grown in DMEM medium, and supplemented with 10% Fetal Bovine Serum (FBS; Bioidea BI201, Iran), 100 µg/ml penicillin G/streptomycin and 1% L-glutamine.

MTT assay

The effect of green tea extract on the viability of LNCaP cells was measured through 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay (CatNo:BI1017, Bioidea, Iran) based on the ability of live cells to cleave the tetrazolium ring into a molecule by absorbance at 490 nm based on manufacturer's instructions. For this purpose, LNCaP cells were plated in 96-well culture plates (5×10³ cells/well), incubated for 24 hr, and treated with green tea extract. After 48 hr at 37 °C, the media were removed, 100 µl of MTT reagent (1 mg/ml) was added into each well, and the cells were incubated at 37 °C for 4 hr. Next, the MTT solution was removed and 50 µl of dimethyl sulfoxide (DMSO) was added into each well to dissolve formazan crystals. Finally, the plates were gently shaken for 10 min and read with an ELISA plate reader (BioTek, USA)¹². In this study, etoposide and deionized water were used as positive and negative controls, respectively.

DAPI staining assay

DAPI (4,6-Diamidino-2-phenylindole dihydrochloride) staining assay was used to determine the changes in the chromatin of LNCaP cells after treatment with green tea extract for 24 hr. DAPI was purchased from Sigma-Aldrich, USA. Briefly, LNCaP cells were seeded in the six-well plates (5×10⁴ cells per well) containing 12 mm cover slips, and consequently treated with green tea extract for 24 hr. Then, the cells were fixed with 3.7% paraformaldehyde, permeabilized in 0.5% (w/v) Triton X-100 and 1% BSA (w/v) for 5 min, washed in Phosphate-Buffered Saline (PBS), and stained through DAPI. All images were taken by using Eclipse Ti-E inverted fluorescent microscope (Nikon, Japan)¹².

Antibodies, SDS-PAGE, and western blot

Anti-β-Actin Antibody (C4) (Santa Cruz Biotechnology, USA), Anti-Bax Antibody (B-9) (Santa Cruz Biotechnology, USA), Bcl-2 Antibody (N-19) (Santa Cruz Biotechnology, USA) were used as primary antibodies for immunoblotting and western blot was performed as previously described³².

RNA and miRNA extraction, cDNA synthesis, and quantitative real-time PCR (qRT-PCR)

In order to perform quantitative real-time RT-PCR analysis, LNCaP cells were lysed after 48 hr along with culturing green tea extract. In addition, total RNA was extracted by using 500 µl of Trizol® reagent based on the manufacturer-provided protocol (Invitrogen Life Technologies, USA), followed by reverse transcription into cDNA by considering manufacturer's protocol (ReveretAid M-Mulv reverse transcriptase kit, Thermo Fisher Scientific, USA). Further, real-time RT-PCR was implemented to amplify cDNA using SYBR Green dye universal master mix (Bioron GmbH, Germany) in Light Cycler 480 (Roche Diagnostics, USA) using the primers for GAPDH-F: 5'-CAA GGT CAT CCA TGA

CAA CTTTG-3', R:5'-GTCCACCACCCTGTTGCTG TAG-3', *Bax*-F: 5'-GTCGCCCTTTTCTACTTTGCC -3', R: 5'-CTCCCGCCACAAAGATGGTCA-3', *BCL*-2F: 5'-CCCCTCGTCCAAGAATGCAA-3', and R: 5'-TCTCCCGGTTATCGTACCCTG-3' for forty cycles. Data represent average copy number normalized to the *GAPDH* housekeeping gene. Also, the primers were synthesized by Pishgam Biotech Co. in Iran. A reaction similar to the above but with deionized water instead of cDNA was set as a negative control. Thermal conditions of the PCR consisted of primary denaturation at 94 °C for 2 min, 45 cycles of denaturation at 94 °C for 30 s, annealing at 59 °C for 30 s, and amplification at 72 °C for 30 s¹² (miR-181a-F: 5'-ACCAACATTCAA CGCTGTC-3'; universal reverse primer: 5'-GAGCA GGGTCCGAGGT-3'). miR-181a and universal reverse primers were purchased from Bon Yakhteh in Iran. The delta Ct values of the samples were classified, and the expression levels were studied using 2- $\Delta\Delta$ Ct (Fold change) values.

Hanging drop formation

The hanging drop method was performed to create a 3D cell culture model and form spheroid³³. In this regard, LNCaP cells were cultured when they had 90% confluence, trypsinized and counted. In addition, ten 20- μ l drops involving 20×10^3 cells were pipetted into the lid of tissue culture dish (60 mm) and 5 ml of PBS was placed in its bottom. The experiment was conducted for the control (cells+medium) and sample (cells+medium containing green tea extract). Furthermore, the number and size of spheroids was determined after three days. Briefly, spheroid formation was monitored using a phase contrast inverted microscope (INV100, BEL Engineering, Italy) and spheroid number was calculated³².

Statistical analysis

The data were analyzed by SPSS v22 (Chicago, USA) and graphs were drawn using Graphpad Prism 7 software. Additionally, the data are expressed as means \pm standard deviation (SD). Further, the experiments were performed three times and two-tailed Student's t test was used to compare the two groups. Finally, p-value less than 0.05 was considered statistically significant¹².

Results

The role of green tea in inducing apoptosis in LNCaP cells

The anti-apoptotic activities of green tea against LNCaP cells were investigated by using MTT assay. To do so, LNCaP cell line was cultured and then treated with different concentrations of green tea extract (100-1000 μ g/ml) for 48 hr (Figure 1A). Next, the morphology of LNCaP cells was analyzed before and after treatment with green tea extract (Figure 1B). To further determine the effect of green tea extract on apoptotic cell death, the cellular nuclear morphology changes of LNCaP cell line were examined by DAPI staining and nuclear fragmentation was clearly shown (Figure 1C). The number of apoptotic nuclei in the cells of the control and sample was compared (Figure 1D). By using quantitative real-time PCR and western blot, the expression of Bax and BCL2 was finally evaluated (Figures 1E, 1F, and 1G). Based on the results, it was found that green tea induces the cellular apoptosis in LNCaP cells.

Antitumor effects of green tea on LNCaP cells via up regulation of miR-181a using 3D cell culture model

A 3D cell culture system is a good reflection of the *in vivo* cell behaviors. By using hanging drop method,




Figure 1. A) The effects of green tea on LNCaP cell viability in different concentrations using MTT assay. B) Comparison of morphological changes of LNCaP before and after treatment with green tea. The experiments were performed three times (original microscope magnification, 20X, scale bar, 10 μ m). C, D) DAPI staining of LNCaP cells under treatment with green tea (sample) compared with control. The experiments were performed three times (original microscope magnification, 40X, scale bar, 10 μ m). E, F) Relative fold-changes in gene expression levels of Bax and BCL2 were shown. Data represent mean \pm SD of three independent experiments. * p<0.05 was considered statistically significant. G) The expression of Bax and BCL2 proteins using western blot was shown. Actin used as an internal control (TCL: Total Cell Lysate).

Antitumor Activities of Green Tea by Up-regulation of miR-181a Expression in LNCaP Cells Using 3D Cell Culture Model




Figure 2. A) Schematic model of spheroid formation using hanging drop technique. B, C) Relative fold-changes in gene expression levels of Bax and BCL2 in LNCaP cells after treatment with green tea (five days) were shown. Data represent mean±SD of three independent experiments. * $p < 0.05$ was considered statistically significant. D, E) Medium containing green tea inhibits size and number of spheroids (magnification: 20x, scale bar: 100 μm ; three independent experiments were done. * $p < 0.05$ was considered statistically significant). F) Relative fold-changes in gene expression levels of miR-181a in LNCaP cells during treatment with green tea were shown. Data represent mean±SD of three independent experiments. * $p < 0.05$ was considered statistically significant.

spheroids are formed in droplets³³. As explained in Materials and Methods section, LNCaP cells were cultured, trypsinized, and counted. Ten drops containing 20×10^3 cells were pipetted into the bottom of the tissue culture dish lid and PBS was placed in its bottom (Figure 2A). After spheroid formation (around 3 days), the expression of Bax and BCL2 were evaluated (Figures 2B and 2C). Next, the size and number of spheroids were analyzed which represent the smaller size and number in the sample (Cells+medium containing green tea) compared to the control (Cells+medium) (Figures 2D and 2E). Finally, the expression of miR-181a in the sample and control group was evaluated. To do so, two days after spheroid formation, LNCaP cells were harvested and the expression of miR-181a was evaluated by qRT-PCR method (Figure 2F). Taken together, these results suggested that green tea has therapeutic effects on LNCaP cells using 3D cell culture model and miR-181a expression was elevated in LNCaP cancer cells after green tea treatment.

Discussion

PCa is considered as one of the most important causes of cancer-associated death in men worldwide. PCa therapy remains a challenge for interested researchers. Moreover, correlation between dietary factors and diseases has been reported. Interestingly, miRNAs have been reported to be potential biomarkers for diagnosis and treatment of cancers due to their management of cellular dysregulations and deregulated expression in a variety of human cancer types. In the present study, the purpose was to evaluate the *in vitro*

antitumor activities of green tea in LNCaP cells. In this regard, 2D and 3D cell culture models were employed for performing our experiments. It was found that green tea inhibited the growth and induced apoptosis in LNCaP cells. Also, it was shown that miR-181a was up-regulated in LNCaP cells after treatment with green tea using 3D cell culture model. Consistent with our finding that miR-181a may act as a tumor suppressor, Shen *et al* showed that the overexpression of miR-181a could significantly inhibit cell proliferation and induce G1 cell cycle arrest in PCa³⁴. Also, Huang *et al* showed that the overexpression of miRNAs members suppressed Non-Small Cell Lung Cancer (NSCLC) proliferation, migration, and invasion and induced cellular apoptosis through BCL2 targeting³⁵. In Glioblastoma, it was found that miR-181a acts as a tumor suppressor by inhibition of cell growth, invasion, and by induction of cell apoptosis²⁶. In another study, it was found that overexpression of miR-181a in K562CML cells inhibits cell growth and induces cellular apoptosis and differentiation. Moreover, it was revealed that overexpression of miR-181a enhances the chemotherapeutic sensitivity of K562 CML cells to imatinib³⁶. In cutaneous squamous cell carcinoma, miR-181a reduces cell viability and suppresses cell apoptosis via down-regulating KRAS³⁷. In breast cancer, it was demonstrated that miR-181a-5p inhibits migration and angiogenesis of cancer cells through down-regulation of matrix metalloproteinase-14³⁸.

In contrast with tumor suppressor roles of miR-181a, the oncogenic roles of miR-181a were shown by several studies. In this regard, it was found that miR-

181a enhanced cell viability and diminished apoptosis by targeting Kruppel-Like Factor 6 (KLF6) in clear renal cell carcinoma³⁹. Tong *et al* reported that miR-181a was up-regulated in prostate cancer tissues compared with adjacent normal tissues¹⁹. Also, miR-181a activates the Epithelial-Mesenchymal Transition (EMT) process in prostate cancer cells through inhibition of TGIF2 (A repressor of the Smad pathway)⁴⁰. Also, the expression level of plasma miR-181 in pancreatic cancer patients was elevated⁴¹ and it was shown that miR-181a promotes pancreatic cancer invasion and progression *via* targeting the tumor suppressor genes, phosphatase and tensin homolog (*PTEN*) and mitogen-activated protein kinase kinase 4 (*MAP2K4*)⁴². Oncogenic role of miR-181a through suppression of WIF-1 (Wnt inhibitory factor-1) in colorectal carcinoma was also shown⁴³. It seems that tumor suppressor or oncogenic roles of miR-181a may be dependent on the cellular and tumor context.

Conclusion

In the present study, it was found that green tea has therapeutic influences on LNCaP cells by induction of apoptosis, suppression of tumor growth, and up regulation of miR-181a expression. However, the targeting of miR-181a in LNCaP cells treated with green tea is unknown. miRNAs play a broad range of roles in cellular processes and they are currently considered as a promising tool in cancer therapy. Therefore, more studies are required to identify the exact molecular mechanisms of miR-181a in prostate cancer after treatment with green tea.

Conflict of Interest

The authors declare that there is no conflict of interest to disclose.

Funding Statement

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA cancer J Clin* 2018;68(6):394-424.
- Jeong WS, Kim IW, Hu R, Kong AN. Modulatory properties of various natural chemopreventive agents on the activation of NF-kappaB signaling pathway. *Pharm Res* 2004;21:661-70.
- Jigisha A, Nishant R, Navin K, Pankaj G. Green tea: A magical herb with miraculous outcomes. *Int Res J Pharm* 2012;3(5):139-48.
- Cabrera C, Artacho R, Gimenez R. Beneficial effects of green tea—A review. *J Am Coll Nutr* 2006;25:79-99.
- Lowe GM, Gana K, Rahman K. Dietary supplementation with green tea extract promotes enhanced human leukocyte activity. *J Complement Integr Med* 2015;12(4):277-82.
- Boschmann M, Thielecke F. The effects of epigallocatechin-3-gallate on thermogenesis and fat oxidation in obese men: A pilot study. *J Am Coll Nutr* 2007;26(4):389S-395S.
- Basu A, Sanchez K, Leyva MJ, Wu M, Betts NM, Aston CE, et al. Green tea supplementation affects body weight, lipids, and lipid peroxidation in obese subjects with metabolic syndrome. *J Am Coll Nutr* 2010;29(1):31-40.
- Ide K, Yamada H, Takuma N, Park M, Wakamiya N, Nakase J, et al. Green tea consumption affects cognitive dysfunction in the elderly: A pilot study. *Nutrients* 2014;6(10):4032-42.
- Hwang JT, Ha J, Park IJ, Lee SK, Baik HW, Kim YM, et al. Apoptotic effect of EGCG in HT-29 colon cancer cells via AMPK signal pathway. *Cancer Lett* 2007;247(1):115-21.
- Nakachi K, Matsuyama S, Miyake S, Suganuma M, Imai K. Preventive effects of drinking green tea on cancer and cardiovascular disease: epidemiological evidence for multiple targeting prevention. *Biofactors* 2000;13(1-4):49-54.
- Shankar S, Ganapathy S, Hingorani SR, Srivastava RK. EGCG inhibits growth, invasion, angiogenesis and metastasis of pancreatic cancer. *Front Biosci* 2008;13:440-52.
- Safari F, Rabieepor M, Jamalomididi F, Baghaeifar Z, Khodaei L. Evaluation of anti-cancer and pro-apoptotic activities of Iranian green tea extract (IGTE) against A549, PC3 and MCF-7 cancer cell lines. *Int J Basic Sci Med* 2019;4(3):113-8.
- Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009;136(2):215-33.
- He L, Hannon G. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 2004;5(7):522-31.
- Esquela-Kerscher A, Slack FJ. Oncomirs-microRNAs with a role in cancer. *Nat Rev Cancer* 2006;6(4):259-69.
- Cai C, Chen QB, Han ZD, Zhang YQ, He HC, Chen JH, et al. miR-195 inhibits tumor progression by targeting RPS6KB1 in human prostate cancer. *Clin Cancer Res* 2015;21(21):4922-34.
- Liang J, Li Y, Daniels G, Sfanos K, De Marzo A, Wei J, et al. Lef1 targeting EMT in prostate cancer invasion is regulated by miR-34a. *Mol Cancer Res* 2015;13(4):681-8.
- Wang M, Ren D, Guo W, Wang Z, Huang S, Du H, et al. Loss of miR-100 enhances migration, invasion, epithelial-mesenchymal transition and stemness properties in prostate cancer cells through targeting Argonaute 2. *Int J Oncol* 2014;45(1):362-72.
- Tong SJ, Liu J, Wang X, Qu LX. microRNA-181 promotes prostate cancer cell proliferation by regulating DAX-1 expression. *Exp Ther Med* 2014;8(4):1296-300.
- Kim CH, Kim HK, Rettig RL, Kim J, Lee ET, Apreli-

- kova O, et al. miRNA signature associated with outcome of gastric cancer patients following chemotherapy. *BMC Med Genomics* 2011;4:79.
21. Ueda T, Volinia S, Okumura H, Shimizu M, Taccioli C, Rossi S, et al: Relation between microRNA expression and progression and prognosis of gastric cancer: A microRNA expression analysis. *Lancet Oncol* 2010;11(2):136-46.
 22. Meng F, Glaser SS, Francis H, DeMorrow S, Han Y, Passarini JD, et al: Functional analysis of microRNAs in human hepatocellular cancer stem cells. *J Cell Mol Med* 2012;16(1):160-73.
 23. Taylor MA, Sossey-Alaoui K, Thompson CL, Danielpour D, Schiemann WP. TGF- β upregulates miR-181a expression to promote breast cancer metastasis. *J Clin Invest* 2013;123(1):150-63.
 24. Shin KH, Bae SD, Hong HS, Kim RH, Kang MK, Park NH. miR-181a shows tumor suppressive effect against oral squamous cell carcinoma cells by downregulating K-ras. *Biochem Biophys Res Commun* 2011;404(4):896-902.
 25. Lin F, Li Y, Yan S, Liu S, Qian W, Shen D, et al. MicroRNA-181a inhibits tumor proliferation, invasiveness, and metastasis and is downregulated in gastric cancer. *Oncol Res* 2015;22(2):75-84.
 26. Shi L, Cheng Z, Zhang J, Li R, Zhao P, Fu Z, et al. hsa-mir-181a and hsa-mir-181b function as tumor suppressors in human glioma cells. *Brain Res* 2008;1236:185-93.
 27. Marton S, Garcia MR, Robello C, Persson H, Trajtenberg F, Pritsch O, et al. Small RNAs analysis in CLL reveals a deregulation of miRNA expression and novel miRNA candidates of putative relevance in CLL pathogenesis. *Leukemia* 2008;22(2):330-8.
 28. Wang H, Bian S, Yang CS. Green tea polyphenol EGCG suppresses lung cancer cell growth through upregulation miR-210 expression caused by stabilizing HIF-1 α . *Carcinogenesis* 2011;32(12):1881-9.
 29. Yamada S, Tsukamoto S, Huang Y, Makio A, Kumazoe M, Yamashita S, et al. Epigallocatechin-3-O-gallate up-regulates microRNA let-7b expression by activating 67-kDa laminin receptor signaling in melanoma cells. *Sci Rep* 2016;6:19225.
 30. Mokhtari H, Yaghmaei B, Sirati-Sabet M, Jafari N, Mardomi A, Abediankenari S, et al. Epigallocatechin-3-gallate enhances the efficacy of microRNA-34 mimic and microRNA-93 inhibitor co-transfection in prostate cancer cell line. *Iran J Allergy Asthma Immunol* 2020;19 (6):612-23.
 31. Lin CH, Wang HH, Chen TH, Chiang MC, Hung PH, Chen YJ. Involvement of microRNA-296 in the inhibitory effect of epigallocatechin gallate against the migratory properties of anoikis-resistant nasopharyngeal carcinoma cells. *Cancers* 2020;12(4):973.
 32. Rahmani Z, Safari F. Evaluating the in vitro therapeutic effects of human amniotic mesenchymal stromal cells on MiaPaca2 pancreatic cancer cells using 2D and 3D cell culture model. *Tissue Cell* 2020;23:68:101479.
 33. Yilmaz O, Sakaraya S. Is "Hanging Drop" a useful method to form spheroids of Jimt, Mcf-7, T-47d, Bt-474 that are breast cancer cell lines? *Single Cell Biol* 2018;7(1):1000170.
 34. Shen H, Weng XD, Liu XH, Yang D, Wang L, Guo J, et al. miR-181a-5p is downregulated and inhibits proliferation and the cell cycle in prostate cancer. *Int J Clin Exp Pathol* 2018;11(8):3969-76.
 35. Huang P, Ye B, Yang Y, Shi J, Zhao H. MicroRNA-181 functions as a tumor suppressor in non-small cell lung cancer (NSCLC) by targeting Bcl-2. *Tumor Biol* 2015;36:3381-7.
 36. Wang G, Zhao R, Zhao X, Chen XI, Wang D, Jin Y, et al. MicroRNA-181a enhances the chemotherapeutic sensitivity of chronic myeloid leukemia to imatinib. *Oncol Lett* 2015;10(5):2835-41.
 37. Neu J, Dziunycz PJ, Dzung A, Lefort K, Falke M, Denzler R, et al. miR-181a decelerates proliferation in cutaneous squamous cell carcinoma by targeting the proto-oncogene KRAS. *PLoS One* 2017;12(9):e0185028.
 38. Li Y, Kuscus C, Banach A, Zhang Q, Pulkoski-Gross A, Kim D, et al. miR-181a-5p inhibits cancer cell migration and angiogenesis via downregulation of matrix metalloproteinase-14. *Cancer Res* 2015;75(13):2674-85.
 39. Lei Z, Ma X, Li H, Zhang Y, Gao Y, Fan Y, et al. Up-regulation of miR-181a in clear cell renal cell carcinoma is associated with lower KLF6 expression, enhanced cell proliferation, accelerated cell cycle transition, and diminished apoptosis. *Urol Oncol* 2018;36(3):93.e23-93.e37.
 40. Zhiping C, Shijun T, Linhui W, Yapei W, Lianxi Q, Qiang D. MiR-181a promotes epithelial to mesenchymal transition of prostate cancer cells by targeting TGIF2. *Eur Rev Med Pharmacol Sci* 2017;21(21):4835-43.
 41. Liu J, Gao J, Du Y, Li Z, Ren Y, Gu J, et al. Combination of plasma microRNAs with serum CA19-9 for early detection of pancreatic cancer. *Int J Cancer* 2012;131(3):683-91.
 42. Liu J, Xu D, Wang Q, Zheng D, Jiang X, Xu L. LPS induced miR-181a promotes pancreatic cancer cell migration via targeting PTEN and MAP2K4. *Dig. Dis. Sci* 2014;59(7):1452-60.
 43. Ji D, Chen Z, Li M, Zhan T, Yao Y, Zhang Z, et al. MicroRNA-181a promotes tumor growth and liver metastasis in colorectal cancer by targeting the tumor suppressor WIF-1. *Mol Cancer* 2014;13:86.