

Investigation of Integron-Associated Resistance Gene Cassettes in Urinary Isolates of *Klebsiella pneumoniae* in Yasuj, Southwestern Iran During 2015-16

Fariba Jahanbin¹, Masoud Marashifard², Sanaz Jamshidi¹, Maryam Zamanzadeh¹, Masumeh Dehshiri³, Seyed Ali Asghar Malek Hosseini⁴, and Seyed Sajjad Khoramrooz^{5,6*}

1. Department of Basic Sciences, Islamic Azad University, Yasuj Branch, Yasuj, Iran

2. Treatment Management of Social Security Organization of Kohgiluyeh and Boyer-Ahmad Province, Yasuj, Iran

3. Cellular and Molecular Research Center, Yasuj University of Medical Sciences, Yasuj, Iran

4. Student Research Committee, Yasuj University of Medical Sciences, Yasuj, Iran

5. Medicinal Plants Research Center, Yasuj University of Medical Sciences, Yasuj, Iran

6. Department of Microbiology, Faculty of Medicine, Yasuj University of Medical Sciences, Yasuj, Iran

Abstract

Background: Growing antibiotic resistance among urinary opportunistic pathogens such as *Klebsiella pneumoniae* (*K. pneumoniae*) has created a worrisome condition in the treatment of the Urinary Tract Infections (UTIs) in recent years. Integrons play a significant role in the dissemination of antibiotic resistance genes. The present study was conducted to investigate class 1-3 integrons and the corresponding resistance gene cassettes in urinary *K. pneumoniae* isolates.

Methods: In this study, from December 2015 to September 2016, a total of 196 *K. pneumoniae* isolates were collected from the patients with UTI referred to medical diagnostic laboratories in Yasuj, Southwestern Iran. Antibiotic susceptibility patterns of isolates were determined using 12 antibiotics by the disc diffusion method. Polymerase Chain Reaction (PCR) was used for detection of integron genes (*int1*, *int2*, and *int3*). The variable regions of integrons were amplified by PCR and sequenced to identify the corresponding gene cassettes.

Results: Thirty-nine different antibiotic resistance profiles were observed among *K. pneumoniae* isolates. Only 12.2% of *K. pneumoniae* isolates were found to harbor the *int1* gene. While 17 (60.7%) out of 28 Multidrug Resistance (MDR) *K. pneumoniae* isolates carried the *int1* gene, only 4.2% of non-MDR isolates harbored *int1* gene. Totally 7 different gene cassette arrays were found in the *int1* gene of *K. pneumoniae* isolates. The *aadA1* was the most prominent gene cassette. Also, high frequency of *dfrA* containing gene cassettes was observed.

Conclusion: Continuous monitoring and characterization of integrons and their associated gene cassettes could be helpful in controlling the rising rate of antibiotic resistance.

Avicenna J Med Biotech 2020; 12(2): 124-131

Keywords: Antibiotic resistance, Integrons, Iran, *Klebsiella pneumoniae*

Introduction

Urinary Tract Infection (UTI) is one of the most prominent infectious diseases in both community and healthcare setting. *Escherichia coli* (*E. coli*) is considered as the most prominent uropathogen which accounts for 75-90% of all UTIs in both inpatients and outpatients followed by *Klebsiella pneumoniae* (*K. pneumoniae*)^{1,2}. The increasing resistance rate to antibiotics among UTI causing organisms makes the empiric treatment of these infections very challenging³. Increasing antibiotic usage, as well as horizontal transfer

of antibiotic resistance genes located on various types of mobile DNA elements as plasmids, transposons and gene cassettes in integron, has facilitated the development of Multidrug Resistance (MDR) in *Enterobacteriaceae* family⁴. Integrons are genetic elements that play a significant role in the transmission of multidrug resistance genes in several gram-negative bacteria⁵. An integron is structurally composed of three genetic elements; *integrase* gene (*intI*) which is responsible for site specific recombination of mobile gene cassettes,

* Corresponding author:
Seyed Sajjad Khoramrooz, Ph.D.,
Medicinal Plants Research
Center, Yasuj University of
Medical Sciences, Yasuj, Iran
Tel/Fax: +98 743 323 5153
E-mail:
Khoramrooz@gmail.com,
masoud.marashifard@gmail.com
Received: 28 Aug 2019
Accepted: 25 Nov 2019

attachment site (*attI*) and the promoter (Pc) ⁶. Gene cassettes are discrete genetic elements consisting of the single Open Reading Frame (ORF) and a recombination site (a 59 bp element) known as the *attC* site ⁷. The integrase mediates the integration of circular gene cassettes between *attI* and *attC* sites ⁸. Integrons are classified into several classes, based on the amino acid sequence homology of *intI* gene, among them class 1, 2 and 3 are usually recovered from clinical isolates ^{9,10}. Class1 integron has been reported in different studies as the most frequent class identified in clinical isolates ⁹. Class1 integron contains variable regions of gene cassettes that are sometimes absent in the structure of integron, flanked by two highly conserved regions; 5'-conserved segment (5'-CS) and 3'-conserved segment (3'-CS) ⁶. Gene cassettes are promoterless variable regions of integrons, which encode antibiotic resistance phenotype, located between *attC* and *attI* region. Their expression depends on the integron promoter (Pc) which relies on the 5'-CS in the case of class1 integron. Different arrays of cassettes have been reported and most arrays had two or three gene cassettes ¹¹.

At least 130 different gene cassettes including diverse resistance genes have been identified, mainly conferring resistance to different classes of antibiotics including aminoglycosides, β -lactams, chloramphenicol, trimethoprim, erythromycin, and rifampicin ^{11,12}. The most prominent gene cassettes within class 1 integron among *Enterobacteriaceae* family are *aadA* and *dfrA*, which confer resistance to streptomycin and trimethoprim, respectively ¹³. Although some studies have reported the prevalence of class 1-3 integrons and gene cassettes in *E. coli*, *Pseudomonas aeruginosa* and *Acinetobacter baumannii* isolates in Iran ¹⁴⁻¹⁷, there are limited data regarding the distribution of gene cassettes in *K. pneumoniae* isolates from Iran ¹⁸. Therefore, in the present study, an attempt was made to investigate the gene cassettes in addition to the prevalence of class 1-3 integrons and antibiotic resistance patterns in a series of clinical isolates of *K. pneumoniae* in Yasuj, Southwestern Iran.

Materials and Methods

Bacterial isolation

From December 2015 to September 2016, 196 non-repetitive isolates of *K. pneumoniae* were collected from the urine samples of patients with UTI who were referred to medical diagnostic laboratories in Yasuj,

Southwestern Iran. Each urine sample was cultured on EMB and Blood Agar (Merck, Germany) and incubated at 37°C for 24 hr. Using conventional biochemical tests on culture media such as Methyl red-Voges Proskauer (MR-VP), Triple Sugar Iron (TSI), Sulfide Indole Motility (SIM), Simmons Citrate, and Urea Agar (Merck, Germany), identification of *K. pneumoniae* isolates was performed ¹⁹. Verified isolates of *K. pneumoniae* were stored at -20°C in TSB medium (Merck, Germany) with 15% glycerol for next steps. This study was approved by the ethical committee of Yasuj University of Medical Sciences.

Antibiotic susceptibility testing (AST)

Susceptibility of *K. pneumoniae* isolates was determined by the disk diffusion method according to the Clinical and Laboratory Standards Institute (CLSI) guidelines ²⁰. Twelve antibiotic disks (MAST, UK) including amoxicillin/clavulanic acid (AMC; 30 μ g), cephalothin (CEP; 30 μ g), ceftazidime (CAZ; 30 μ g), imipenem (IMP; 10 μ g), chloramphenicol (CLR; 30 μ g), nitrofurantoin (NI; 300 μ g), tetracycline (TET; 30 μ g), gentamycin (GEN; 10 μ g), amikacin (AMI; 30 μ g), ciprofloxacin (CIPR; 5 μ g), nalidixic acid (NAL; 30 μ g), and trimethoprim-sulfamethoxazole (SXT; 25 mg) were used. *K. pneumoniae* ATCC 700603 was used as the control for antibiotic resistance. MDR was outlined as acquired non-susceptibility to a minimum of one agent in 3 or more antimicrobial classes ²¹.

DNA extraction

The boiling method with some modification was used to extract the genome of bacteria ²². In short, 0.5 McFarland bacterial suspension was prepared first, and then 300 μ l of it was transferred to the 1.5 ml microtube containing sterile distilled water. The suspension within the microtube was homogenized with the vortex and then boiled for 10 min at 100°C in a water bath. In the next step, microtube was centrifuged for 10 min at 14000 g and the supernatant containing genome was transferred to the 0.5 ml microtube and kept at 18°C until the next step.

PCR for detection of integron genes

Polymerase Chain Reaction (PCR) was carried out for detection of *integron genes* (*intI1*, *intI2*, and *intI3*) using the primers described in table 1 ^{23,24}. Each reaction mixture in a final volume of 25 μ l contained 12.5 μ l of 2X Master mix (Amplicon, Denmark), 0.3 μ l of each primer (20 pM/ μ l), 2.5 μ l of DNA template and

Table 1. Primers used for the detection of class 1-3 integrons and variable region of class 1 integrons

Target genes	Oligonucleotide sequences of primers (5' to 3')	Expected size (bp)	Reference
<i>intI1</i>	F: CCTCCCGCACGATGATC R: TCCACGCATCGTCAGGC	280	Kraft <i>et al</i> (1986)
<i>intI2</i>	F: TTATTGCTGGGATTAGGC R: ACGGCTACCCTCTGTTATC	233	Goldstein <i>et al</i> (2001)
<i>intI3</i>	F: AGTGGGTGGCGAATGAGTG R: TGTTCTTGATCGGCAGGTTG	600	Goldstein <i>et al</i> (2001)
Variable region of class 1 integrons	5'-CS: GGCATCCAAGCAGCAAG 3'-CS: AAGCAGACTTGACCTGA	variable	Levesque <i>et al</i> (1995)

9.4 μ l of distilled water.

The planned conditions in the thermocycler (Bio-Rad, T100, USA) for amplification of each of the genes examined were as follows; initial denaturation at 94°C for 4 min; 35 thermal cycles of denaturation at 94°C for 40 s, annealing at 58°C (For *intI1* genes) or 51°C (For *intI2* and *intI3* genes) for 30 s, and extension at 72°C for 40 s; and a post-PCR final incubation at 72°C for five minutes. The amplified PCR products were electrophoresed on 1.5% agarose gel containing 0.5 μ g/ml ethidium bromide and visualized under a UV transilluminator (Major science, Taiwan).

Characterization of gene cassettes inserted in the variable regions of class 1 integrons

PCR was carried out for amplification of variable regions of class 1 integron similar to the planned conditions for the detection of *intI1* gene, except that the annealing temperature was 60°C using primers described in table 1²⁵. For each integron-positive isolate, some of the PCR product was electrophoresed and, after ensuring that the expected fragment was present, the remaining PCR product was sent to Macrogen company (Seoul, South Korea) for sequencing.

Obtained sequences were submitted in the NCBI database and blasted by online BLAST search (<http://www.ncbi.nlm.nih.gov/BLAST/>).

Statistical analysis

Data were analyzed with SPSS software (Version 15, Chicago, IL, USA). The chi-square and Fisher's exact tests were used for determining the association between presence of *intI* genes and antibiotic resistance status. A $p < 0.05$ was regarded statistically significant.

Results

Antibiotic susceptibility test results

Among 196 studied isolates of *K. pneumoniae*, 137 isolates (69.9%) were collected from females and the remaining 59 isolates (30.1%) were obtained from male patients. The results of the AST showed that resistance to amoxicillin is highest (38.8%). Resistance rate to other antibiotics was as follows; they were cephalothin (32.1%), nitrofurantoin (22.4%), ceftazidime (12/8%) trimethoprim-sulfamethoxazole (9.7%), tetracycline (8.7%), gentamycin (5.1%), nalidixic acid (3.6%), chloramphenicol (3.6%), ciprofloxacin (2.6%), and amikacin (1%). There was not any imipenem-resistant isolate. Twenty-eight (14.3%) isolates were MDR and 39 different antibiotic resistance profiles were observed (Table 2).

Detection of integrons and characterization of gene cassettes

Among the 196 *K. pneumoniae* isolates, class 1 in-

Table 2. *intI1* gene presence status in *K. pneumoniae* isolates according to antibiotic susceptibility pattern

Antibiotics	<i>intI1</i> gene presence	Antibiotic resistance pattern			p-value
		Susceptible (%)	Intermediate (%)	Resistant (%)	
SXT	<i>intI1+</i> (24)	7 (29.2)	0	17 (70.8)	<0.001
	<i>intI1-</i> (172)	169 (98.3)	1 (0.6)	2 (1.2)	
TET	<i>intI1+</i> (24)	7 (29.2)	6 (25)	11 (45.8)	<0.001
	<i>intI1-</i> (172)	162 (94.2)	4 (2.3)	6 (3.5)	
NI	<i>intI1+</i> (24)	9 (37.5)	8 (33.3)	7 (29.2)	0.642
	<i>intI1-</i> (172)	79 (45.9)	56 (32.6)	37 (21.5)	
GEN	<i>intI1+</i> (24)	19 (79.2)	1 (0.5)	4 (16.7)	0.003
	<i>intI1-</i> (172)	166 (96.5)	0	6 (3.5)	
AMI	<i>intI1+</i> (24)	24 (100)	0	0	1
	<i>intI1-</i> (172)	168 (97.7)	2 (1.2)	2 (1.2)	
CIP	<i>intI1+</i> (24)	17 (70.8)	3 (12.5)	4 (16.7)	<0.001
	<i>intI1-</i> (172)	169 (98.3)	2 (1.2)	1 (0.6)	
NAL	<i>intI1+</i> (24)	18 (75)	2 (8.3)	4 (16.7)	<0.001
	<i>intI1-</i> (172)	168 (97.7)	1 (0.6)	3 (1.7)	
CLR	<i>intI1+</i> (24)	17 (70.8)	2 (8.3)	5 (20.8)	<0.001
	<i>intI1-</i> (172)	169 (98.3)	1 (0.6)	2 (1.2)	
AMC	<i>intI1+</i> (24)	1 (4.2)	2 (8.3)	21 (87.5)	<0.001
	<i>intI1-</i> (172)	50 (29.1)	67 (39)	55 (32)	
CEP	<i>intI1+</i> (24)	6 (25)	2 (8.3)	16 (66.7)	0.001
	<i>intI1-</i> (172)	100 (58.1)	25 (14.5)	47 (27.3)	
CAZ	<i>intI1+</i> (24)	11 (45.8)	4 (16.7)	9 (37.5)	<0.001
	<i>intI1-</i> (172)	154 (89.5)	2 (1.2)	16 (9.3)	

SXT=Trimethoprim-Sulfamethoxazole, TET=Tetracycline, NI=Nitrofurantoin, GEN=Gentamycin, AMI=Amikacin, CIP=Ciprofloxacin, NAL=Nalidixic acid, CLR=Chloramphenicol, AMC=Amoxicillin/Clavulanic acid, CEP=Cephalothin, CAZ=Ceftazidime.

Table 3. Frequency of *intI1* gene-positive isolates in each of the resistance profiles

No.	Resistance phenotypes	No. of isolates	No. of <i>intI1</i> gene-positive isolates
1	No resistance (0)	68	0
2	SXT(1)	1	1
3	AMC(1)	20	1
4	CEP(1)	12	0
5	CAZ(1)	3	0
6	NI(1)	25	0
7	TET(1)	4	2
8	AMI(1)	1	0
9	SXT, AMC(2)	2	2
10	AMC, CEP(2)	15	0
11	AMC, CAZ(2)	1	1
12	AMC, NI(2)	5	0
13	CEP, CAZ(2)	2	0
14	CEP, NI(2)	3	0
15	CAZ, NI(2)	1	0
16	SXT, AMC, CEP(3)	2	2
17	SXT, AMC, CLR(3)	2	1
18	AMC, CEP, TET(3)	1	1
19	AMC, CEP, GEN(3)	1	0
20	CAZ, AMC, CEP(3) (non MDR)	5	0
21	NAL, AMC, CEP, TET(4)	1	0
22	SXT, AMC, CEP, CAZ(4)	1	1
23	SXT, AMC, CEP, GEN(4)	2	2
24	AMC, CEP, CAZ, NI(4)	1	0
25	AMC, CEP, CAZ, GEN(4)	1	0
26	AMC, CEP, NI, TET(4)	1	1
27	AMC, CEP, TET, GEN(4)	1	0
28	CIPR, NAL, AMC, CEP, GEN(5)	1	0
29	NAL, AMC, CEP, CAZ, NI(5)	1	0
30	SXT, AMC, CEP, CAZ, TET(5)	1	1
31	AMC, CEP, CAZ, NI, TET(5)	2	1
32	AMC, CEP, CAZ, GEN, AMI(5)	1	0
33	SXT, AMC, CEP, CAZ, CLR, TET(6)	1	1
34	SXT, AMC, CEP, CAZ, NI, GEN(6)	1	1
35	SXT, AMC, CEP, CLR, TET, GEN(6)	1	0
36	CIPR, NAL, SXT, AMC, CEP, NI, TET(7)	2	2
37	SXT, AMC, CEP, CAZ, CLR, TET, GEN(7)	1	1
38	CIPR, NAL, SXT, AMC, CEP, CAZ, CLR, NI(8)	1	1
39	CIPR, NAL, SXT, AMC, CEP, CAZ, CLR, NI, TET(9)	1	1

tegron (*intI1*) was identified in 24 (12.2%) isolates while *intI2* and *intI3* genes were not found in any of the studied *K. pneumoniae* isolates. In table 2, the frequency of *intI1* gene-positive isolates in each of the resistance profiles is shown.

Significant differences were observed between susceptibility patterns of trimethoprim-sulfamethoxazole, tetracycline, amoxicillin/clavulanic acid, ceftazidime, ciprofloxacin, chloramphenicol, nalidixic acid, cephalothin, gentamycin and presence of class1 integron in *K. pneumonia* isolates (Table 3). While 17 out of 28 (60.7%) MDR *K. pneumoniae* isolates harbored the *intI1* gene, only 4.2% of non-MDR isolates carried *intI1* gene (p<0.001).

Eighteen out of 24 (75%) *intI1* positive *K. pneumoniae* isolates were found to have 7 different gene cassette arrays including the *dfrA5*, *dfrA25*, *dfrA7*, *aadA1*, *dfrA17-aadA5*, *dfrA1-orfC* and *aadB-cat-bla_{OXA10}-aadA1* (Table 4, Figure 1). In the remaining 6 *intI1* po-

sitive *K. pneumoniae* isolates, variable region was not detected.

Among seven different cassette arrays found in *K. pneumoniae*, *aadA1* was the most frequent gene cassette detected in 9 (37.5%) out of 24 *intI1* positive isolates. *dfrA5*, *dfrA25*, *dfrA7*, *dfrA1-orfC* and *aadB-cat-bla_{OXA10}-aadA1* gene cassette arrays were found in one isolate. Four isolates were found to be positive for *dfrA17-aadA5* gene cassette array.

Discussion

Similar to previous reports from Iran and other countries, all of the studied *K. pneumoniae* isolates were found to be susceptible to imipenem²⁶⁻³⁰. Also, in our previous study, only 1% of urinary *E. coli* isolates were imipenem resistant²². The low and prudent administration of imipenem by physicians in Iran and consequently reducing selection pressure of this antibiotic on bacteria has made it the most reliable and

Investigation of Integron-Associated Resistance Gene Cassettes

Table 4. Gene cassette arrays found in class 1 integrons and related resistance profiles

Gene cassette arrays	Approximate amplicon size	No. of isolates (%)	Resistance profiles
<i>dfrA5</i>	700 bp	1 (4.2)	AMC, CAZ
<i>dfrA25</i>	700 bp	1 (4.2)	AMC, CEP, TET
<i>dfrA7</i>	750 bp	1 (4.2)	CIPR, NAL, SXT, AMC, CEP, CAZ, CLR, NI, TET
<i>aadA1</i>	1000 bp	9 (37.5)	AMC TET (2 case) SXT, AMC SXT, AMC, CEP (2 case) AMC, CEP, NI, TET SXT, AMC, CEP, CAZ, NI, GEN CIPR, NAL, SXT, AMC, CEP, NI, TET
<i>dfrA1-orfC</i>	1300 bp	1 (4.2)	SXT, AMC, CLR SXT, AMC
<i>dfrA17-aadA5</i>	1650 bp	4 (16.7)	SXT, AMC, CEP, CAZ SXT, AMC, CEP, GEN CIPR, NAL, SXT, AMC, CEP, CAZ, CLR, NI, TET
<i>aadB-cat-bla_{OXA10}-aadA1</i>	3000 bp	1 (4.2)	SXT, AMC, CEP, CAZ, TET

most effective antibiotic against bacteria such as *E. coli* and *K. pneumoniae*. In the present study, the rate of resistance to trimethoprim-sulfamethoxazole (9.7%), tetracycline (8.7%), gentamycin (5.1%), nalidixic acid (3.6%), chloramphenicol (3.6%), ciprofloxacin (2.6%) and amikacin (1%) all were less than 10%. These relatively low rates were not observed in other studies from Iran and other countries in recent years. The resistance rate of urinary *K. pneumoniae* isolates in a study by Akram *et al* in India was reported as co-trimoxazole 53%, tetracycline 53%, gentamycin 53%, ciprofloxacin 47%, and amikacin 35%³¹. Also, in another study in India by Mariya and Hatkar, the rate of resistance to co-trimoxazole and amikacin was 59.73% and 31.95%, respectively²⁹. In another study in Iran, the resistance rate of *K. pneumoniae* isolates was reported as co-trimoxazole 95.3%, tetracycline 64.7%, gentamycin 76%, chloramphenicol 57.3%, ciprofloxacin 43.3%, and amikacin 50.7%³². Differences in the pattern of antibiotic resistance in different countries and even in different regions of a country may depend on different factors such as rate of access and use of different antibiotics, population, climatic conditions and public health levels³³.

In the present study, the highest rate of antibiotic resistance was observed for amoxicillin/clavulanic acid in 38.8% of *K. pneumoniae* isolates. Unfortunately, in Iran, the arbitrary use of this antibiotic among people has a very high prevalence that increases the selective pressure of the antibiotic on bacteria leading to resistance mechanisms.

In addition to amoxicillin/clavulanic acid, the resistance rate of our isolates was found to be higher than 10% for only three other antibiotics including cephalothin (32.1%), nitrofurantoin (22.4%) and ceftazidime (12/8%).

In the present study, class1 integron was detected in 12.2% of *K. pneumoniae*. This frequency is much lower than that reported by Ahangarzadeh Rezaee *et al* in




Figure 1. PCR assay for identification of some gene cassette arrays among *K. pneumoniae* isolates. Lane M: DNA marker, lane 1: *aadA1* (1000 bp), lane 2: *dfrA7* (750 bp), lane 3 and 5: *dfrA17-aadA5* (1650 bp), lane 4: *aadB-cat-bla_{OXA10}-aadA1* (3000 bp), lane 6: *dfrA1-orfC* (1300 bp).

northwest Iran (78%)³². Considering that in the study of Ahangarzadeh Rezaee *et al*, 99.3% of isolates were MDR and in our study only 14.3% of isolates were MDR, this high difference between the presence of the integron in these two studies was not surprising. The above comparison may illustrate well the association between the emergence of MDR bacteria and the presence of integron. Also, Li *et al* and Lina *et al* reported a higher prevalence of class I integron compared to the current study^{4,26}. The prevalence of class I integron in their studies was 51.1% and 54%, respectively. On the other hand, in another study in Iran by Seyed Javadi *et al*, the prevalence of class I integrons among *K. pneumoniae* isolates (13.3%) was very close to our study³⁴.

Similar to other studies class 2 and class 3 integrons were not found in any of our *K. pneumoniae* isolates,

while Ahangarzadeh Rezaee *et al* reported class 2 integrons in 13.4 % of their isolates^{25,32,35,36}.

Significant differences were observed between the presence of the *intI1* gene and susceptibility patterns of all antibiotics except amikacin and nitrofurantoin. Similar to our study, Li *et al* did not find a significant association between resistance to nitrofurantoin and the presence of class1 integron⁴.

In the present study, 7 different cassette arrays were found in 75% of *intI1* positive *K. pneumoniae* isolates. Among them, *aadA1* was the most frequent gene cassette detected in 37.5% of 24 *intI1* positive isolates. The *aadA* type gene cassettes confer resistance to aminoglycosides such as streptomycin that was widely used for treating UTIs in the early years³⁷. Since streptomycin usage has been limited to some specific human diseases, its frequent application in agriculture and food animals, especially in livestock production, may bring development of resistance *via* acquiring resistance gene cassettes³⁷. Hence, this gene cassette can remain in animal pathogenic strains of *E. coli* and be disseminated to other strains by horizontal gene transfer mechanisms.

The high prevalence rate of gene cassette arrays with *dfr* genes in the current study, which codes for Dihydrofolate Reductase (DHFR) conferring resistance to trimethoprim, can reflect the wide use of trimethoprim in the treatment of urinary tract infections in recent years.

Similar to findings of previous studies^{4,19}, in the present work, 25% of *K. pneumoniae* isolates were found to carry class1 integron without gene cassettes. In a study from China performed on *K. pneumoniae* isolates, *dfrA1-orfC* was the most predominant cassette array among 10 different identified cassette arrays among which *dfrA5*, *aadA1* and *dfrA1-orfC* gene cassettes were also detected in our study⁴. Also, Salimzand *et al* identified only 3 gene cassettes or cassette arrays in *K. pneumoniae* isolates including *arr-5*, *aacA4-orfD*, and *dfrA17-aadA5*, among which the latter cassette was found in our study²⁷.

In the present study, all isolates with the *dfrA7*, *dfrA17-aadA5*, and *dfrA1-orfC* gene cassette arrays exhibited a resistance phenotype to trimethoprim sulfamethoxazole. Two isolates with *dfrA25* and *dfrA5* cassettes, contrary to our expectation, have not shown resistance to trimethoprim sulfamethoxazole. The reason for this can be found in the presence of weak promoters or the presence of mutations in the region between the two specific sequences of promoters³⁸.

The *cat* gene that codes for Chloramphenicol Acetyl Transferase (CAT) conferring resistance to chloramphenicol was found in *aadB-cat-bla_{OXA10}-aadA1* cassette array. Although this gene cassette array has already been reported from China³⁹, identification of this gene cassette array in *K. pneumoniae* isolates was reported for the first time in Iran. Surprisingly, in the current study, this cassette array was found in one chlo-

ramphenicol sensitive isolate and despite the presence of *cat* cassette, a large distance between the promoter and the corresponding gene cassette may affect the transcription of *cat* gene in this isolate³⁸. *bla_{OXA10}* gene found in *aadB-cat-bla_{OXA10}-aadA1* cassette array encodes for β -lactamase enzymes conferring resistance to beta-lactam antibiotics such as amoxicillin and ampicillin⁴⁰.

Conclusion

In the present study, the proportional prevalence of integrons and the antibiotic resistance among *K. pneumoniae* isolates clearly shows the importance of integrons in dissemination of antibiotic resistance. *aadA1* was found to be the most prominent gene cassette among 7 different cassette arrays identified in *K. pneumoniae* isolates. Continuous monitoring and characterization of integrons and their associated gene cassettes could be helpful in controlling the rate of antibiotic resistance by planning to take preventive measures to hinder the spread of resistant strains.

Acknowledgement

This study has been supported by Deputy of Research and Technology, Yasuj University of Medical Sciences.

Conflict of Interest

The authors declare no conflicts of interest.

References

1. El-Najjar NG, Farah MJ, Hashwa FA, Tokajian ST. Antibiotic resistance patterns and sequencing of class I integron from uropathogenic Escherichia coli in Lebanon. Lett Appl Microbiol 2010;51(4):456-461.
2. Salem MM, Magdy M, Alhosiny IM. Distribution of classes 1 and 2 integrons among multi drug resistant E. coli isolated from hospitalized patients with urinary tract infection in Cairo, Egypt. Aust J Basic Appl Sci 2010; 4(3):398-407.
3. Márquez C, Labbate M, Raymondo C, Fernández J, Gestal AM, Holley M, et al. Urinary tract infections in a South American population: dynamic spread of class 1 integrons and multidrug resistance by homologous and site-specific recombination. J Clin Microbiol 2008;46(10):3417-3425.
4. Li B, Hu Y, Wang Q, Yi Y, Woo PC, Jing H, et al. Structural diversity of class 1 integrons and their as-associated gene cassettes in Klebsiella pneumoniae isolates from a hospital in China. PloS One 2013;8(9):e75805.
5. Japoni A, Gudarzi M, Farshad S, Basiri E, Ziyaeyan M, Alborzi A, et al. Assay for integrons and pattern of antibiotic resistance in clinical Escherichia coli strains by PCR-RFLP in Southern Iran. Jpn J Infect Dis 2008;61(1):85-88.
6. Domingues S, da Silva GJ, Nielsen KM. Global dissemination patterns of common gene cassette arrays in class 1 integrons. Microbiology 2015;161(7):1313-1337.

7. Gillings MR. Integrons: past, present, and future. *Microbiol Mol Biol Rev* 2014;78(2):257-277.
8. Stokes HW, O'gorman DB, Recchia GD, Parsekhian M, Hall RM. Structure and function of 59-base element recombination sites associated with mobile gene cassettes. *Mol Microbiol* 1997;26(4):731-745.
9. Deng Y, Bao X, Ji L, Chen L, Liu J, Miao J, et al. Resistance integrons: class 1, 2 and 3 integrons. *Ann Clin Microbiol Antimicrob* 2015;14:45.
10. Partridge SR. Analysis of antibiotic resistance regions in Gram-negative bacteria. *FEMS Microbiol Rev* 2011;35(5):820-855.
11. Partridge SR, Tsafnat G, Coiera E, Iredell JR. Gene cassettes and cassette arrays in mobile resistance integrons. *FEMS Microbiol Rev* 2009;33(4):757-784.
12. Mazel D. Integrons: agents of bacterial evolution. *Nat Rev Microbiol* 2006;4(8):608-620.
13. DeLappe N, O'Halloran F, Fanning S, Corbett-Feeney G, Cheasty T, Cormican M. Antimicrobial resistance and genetic diversity of *Shigella sonnei* isolates from western Ireland, an area of low incidence of infection. *J Clin Microbiol* 2003;41(5):1919-1924.
14. Zeighami H, Haghi F, Masumian N, Hemmati F, Samei A, Naderi G. Distribution of integrons and gene cassettes among uropathogenic and diarrheagenic *Escherichia coli* isolates in Iran. *Microb Drug Resist* 2015;21(4):435-440.
15. Akrami F, Shahandashti EF, Yahyapour Y, Sadeghi M, Khafri S, Pournajaf A, et al. Integron types, gene cassettes and antimicrobial resistance profile of *Acinetobacter baumannii* isolated from BAL samples in Babol, north of Iran. *Microb Pathog* 2017;109:35-38.
16. Haghi F, Keramati N, Hemmati F, Zeighami H. Distribution of integrons and gene cassettes among metallo- β -lactamase producing *Pseudomonas aeruginosa* clinical isolates. *Infect Epidemiol Microbiol* 2017;3(2):36-40.
17. Aky A, Lorestani RC, Rostamian M, Elahi A, Baakhshii S, Aliabadi M, et al. The relationship of class I integron gene cassettes and the multidrug-resistance in extended-spectrum β -lactamase producing isolates of *Escherichia coli*. *Pediatr Infect Dis J* 2019;7(3).e87961.
18. Firoozeh F, Mahluji Z, Khorshidi A, Zibaei M. Molecular characterization of class 1, 2 and 3 integrons in clinical multi-drug resistant *Klebsiella pneumoniae* isolates. *Antimicrob Resist Infect Control* 2019;8:59.
19. Mahon CR, Lehman DC, Manuselis G. Textbook of diagnostic microbiology-E-Book. 6th ed. USA: Elsevier health sciences; 2014. 1024 p.
20. Clinical and Laboratory Standards Institute (2015) Performance standards for antimicrobial susceptibility testing. In: Twenty-Fifth informational supplement, Wayne, PA: CLSI: M100- S25.
21. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012;18(3):268-281.
22. Khoramrooz SS, Sharifi A, Yazdanpanah M, Malek Hosseini SA, Emaneini M, Gharibpour F, et al. High frequency of class 1 integrons in *Escherichia coli* isolated from patients with urinary tract infections in Yasuj, Iran. *Iran Red Crescent Med J* 2016;18(1):e26399.
23. Kraft CA, Timbury MC, Platt DJ. Distribution and genetic location of Tn7 in trimethoprim-resistant *Escherichia coli*. *J Med Microbiol* 1986;22(2):125-131.
24. Goldstein C, Lee MD, Sanchez S, Hudson C, Phillips B, Register B, et al. Incidence of class 1 and 2 integrases in clinical and commensal bacteria from livestock, companion animals, and exotics. *Antimicrob Agents Chemother* 2001;45(3):723-726.
25. Levesque C, Piche L, Larose C, Roy PH. PCR mapping of integrons reveals several novel combinations of resistance genes. *Antimicrob Agents Chemother* 1995;39(1):185-191.
26. Lina TT, Rahman SR, Gomes DJ. Multiple-antibiotic resistance mediated by plasmids and integrons in uropathogenic *Escherichia coli* and *Klebsiella pneumoniae*. *Bangladesh J Microbiol* 2007;24(1):19-23.
27. Salimizand H, Shahcheraghi F, Kalantar E, Badmasti F, Mousavi SF. Molecular characterization of class 1 integrons and gene cassettes in multidrug resistant (MDR) *Klebsiella* spp. isolated from hospitalized and outpatients in Iran, 2009. *Iran J Microbiol* 2013;5(1):48-55.
28. Moini AS, Soltani B, Ardakani AT, Moravveji A, Erami M, Rezaei MH, et al. Multidrug-resistant *Escherichia coli* and *Klebsiella pneumoniae* isolated from patients in Kashan, Iran. *Jundishapur J Microbiol* 2015;8(10):e27517.
29. Mariya S, Hatkar SS. Antimicrobial susceptibility profile of urinary isolates of *Escherichia coli* and *Klebsiella pneumoniae*. *Int J Health Sci Res* 2015;5(2):169-172.
30. Najjuka CF, Kateete DP, Kajumbula HM, Joloba ML, Essack SY. Antimicrobial susceptibility profiles of *Escherichia coli* and *Klebsiella pneumoniae* isolated from outpatients in urban and rural districts of Uganda. *BMC Res Notes* 2016;9(1):235.
31. Akram M, Shahid M, Khan AU. Etiology and antibiotic resistance patterns of community-acquired urinary tract infections in JNMC Hospital Aligarh, India. *Ann Clin Microbiol Antimicrob* 2007;6(1):4.
32. Ahangarzadeh Rezaee M, Langarizadeh N, Aghazadeh M. First report of class 1 and class 2 integrons in multidrug-resistant *Klebsiella pneumoniae* isolates from northwest Iran. *Jpn J Infect Dis* 2012;65(3):256-259.
33. Hendriksen RS, Munk P, Njage P, van Bunnik B, McNally L, Lukjancenko O, et al. Global monitoring of antimicrobial resistance based on metagenomics analyses of urban sewage. *Nat Commun* 2019;10(1):1124.
34. Seyedjavadi S, Eslami G, Goudarzi M, Goudarzi H, Fallah F. Integrons and multidrug resistance among *E. coli* and *Klebsiella pneumoniae* isolated from children with urinary tract infections. *Health Med* 2013;7(1):243-249.
35. Derakhshan S, Peerayeh SN, Fallah F, Bakhshi B, Rahbar M, Ashrafi A. Detection of class 1, 2, and 3 integrons among *Klebsiella pneumoniae* isolated from children in Tehran hospitals. *Arch Pediatr Infect Dis* 2014;2(1):164-168.

36. Haddadi A, Mohammadi R, Harzandi N. Prevalence of integrons as the carrier of multidrug resistance genes among clinical isolates of klebsiella. *J Med Bacteriol* 2019;8(3, 4):23-30.
37. Yan H, Li L, Zong M, Alam MJ, Shinoda S, Shi L. Occurrence and characteristics of class 1 and 2 integrons in clinical bacterial isolates from patients in South China. *J Health Sci* 2010;56(4):442-450.
38. Cocchi S, Grasselli E, Gutacker M, Benagli C, Convert M, Piffaretti JC. Distribution and characterization of integrons in *Escherichia coli* strains of animal and human origin. *FEMS Immunol Med Microbiol* 2007;50(1):126-132.
39. Gu B, Pan S, Wang T, Zhao W, Mei Y, Huang P, et al. Novel cassette arrays of integrons in clinical strains of Enterobacteriaceae in China. *Int J Antimicrob Agents* 2008;32(6):529-633.
40. Sivri N, Sandalli C, Ozgumus OB, Colakoglu F, Dogan D. Antibiotic resistance profiles of enteric bacteria isolated from Kucukcekmece Lagoon (Istanbul-Turkey). *Turk J Fish Aquat Sci* 2012;12(3):699-707.