

## Anti-Quorum Sensing and Anti-Biofilm Activity of Ginger (*Zingiber officinale*) Rhizomes against Multidrug-Resistant Clinical Isolates of *Pseudomonas aeruginosa*

Pankaj Kumar Sagar <sup>1</sup>, Poonam Sharma <sup>2</sup> and Rambir Singh <sup>3\*</sup>

1. Department of Microbiology, Bundelkhand University, Jhansi-284128, Uttar Pradesh, India

2. Department of Zoology, Indira Gandhi National Tribal University (A Central University), Amarkantak-484886, Madhya Pradesh, India

3. Department of Horticulture, Aromatic and Medicinal Plants, Mizoram University (A Central University), Aizawl-796004, Mizoram, India

### Abstract

**Background:** The aim of this study was to determination of Anti-Quorum Sensing (AQS) and anti-biofilm potential of the methanol extract of ginger (*Zingiber officinale*) rhizomes against multidrug-resistant clinical isolates of *Pseudomonas aeruginosa* (*P. aeruginosa*).

**Methods:** The AQS activity of ginger was determined against *Chromobacterium violaceum* (*C. violaceum*) ATCC 12472 (CV12472), a biosensor strain, in qualitative manner using the agar well diffusion method. The violacein pigment inhibition was assessed to confirm AQS activity of ginger. The AQS potential of sub-minimum Inhibitory Concentrations (sub-MICs) of the ginger extract was determined by targeting different QS regulated virulence factors, including swarming motility (using swarm diameter measurement method), pyocyanin pigment (using chloroform extraction method), Exopolysaccharide (EPS) (using phenol-sulphuric acid method), and biofilm formation (using microtiter plate assay), against clinical isolates (CIs 2, 3, and 4) and standard reference strain of *P. aeruginosa* (PA01).

**Results:** The AQS activity of methanol extract of ginger was confirmed against *C. violaceum* (CV12472) as inhibition of violacein pigment formation without effecting the growth of CIs and PA01 of *P. aeruginosa*. The ginger extract exhibited concentration-dependent inhibition of virulence factors and biofilm formation. The maximum reduction was found in swarming motility, pyocyanin, EPS and biofilm formation against PA01 (51.38%), CI3 (57.91%), PA01 (63.29%) and CI2 (64.37%), respectively at 1/2 MIC of ginger extract.

**Conclusion:** The results of present study revealed the effective AQS and anti-biofilm potential of *Zingiber officinale* rhizome methanol extract at a reduced dose (sub-MICs). The extract may be explored as an agent of antimicrobial compounds having AQS and anti-biofilm activity for controlling microbial infection and also for reducing the chances of emergence of resistance in *P. aeruginosa*.

**Keywords:** *Chromobacterium violaceum*, Ginger, *Pseudomonas aeruginosa*, Pyocyanin, Rhizome  
**To cite this article:** Kumar Sagar P, Sharma P, Singh R. Anti-Quorum Sensing and Anti-Biofilm Activity of Ginger (*Zingiber officinale*) Rhizomes against Multidrug-Resistant Clinical Isolates of *Pseudomonas aeruginosa*. Avicenna J Med Biotech 2024;16(1):49-56.

\* Corresponding author:  
Rambir Singh, Ph.D., Department of Horticulture, Aromatic and Medicinal Plants, Mizoram University (A Central University), Aizawl-796004, Mizoram, India  
Tel: +91 9473583251  
E-mail: sehrawat\_r@yahoo.com  
Received: 9 Jun 2023  
Accepted: 19 Aug 2023

### Introduction

Continuous increasing resistance against conventional antibiotics in pathogenic micro-organism has become a major obstacle in the treatment of microbial infections. In the present scenario, along with the new therapeutic agents, antimicrobial mechanism and alternate drug target should also be focused for the effective solution of Multidrug-Resistant (MDR) microbial in-

fections. *Pseudomonas aeruginosa* (*P. aeruginosa*) is a gram-negative opportunistic bacterium that exhibits MDR pattern and a cause of nosocomial infection, urinary tract infection, burn wound infection, eye infection, respiratory infections and also infecting cystic fibrosis patients, Acquired Immunodeficiency Syndrome (AIDS), and pulmonary diseases <sup>1-4</sup>. Antibiotics,

are most commonly used for the treatment of infections caused by *P. aeruginosa*<sup>5</sup>, generally depending on growth inhibition mechanism. Quorum Sensing (QS), a cell to cell communication gene regulatory mechanism, targets regulation of various virulence factors and biofilm formation rather than growth inhibition or killing bacteria in several gram-negative bacteria such as *P. aeruginosa*<sup>6-8</sup>. In *P. aeruginosa*, the two major QS systems are lasI-lasR and rhlI-rhlR which are inter-related and functions in a hierarchical manner<sup>9,10</sup>. lasI-lasR and rhlI-rhlR quorum sensing systems involve auto inducer signal molecule, N-(3-oxododecanoyl) homoserine lactone (OdDHL) and N-butanoyl homoserine lactone (BHL), respectively<sup>11,12</sup>. *Pseudomonas* quinolone signal (PQS), (2-heptyl-hydroxy-4-quinolone), a third intercellular signal has also been investigated in *P. aeruginosa*<sup>13</sup>.

Various virulence factors such as swarming motility, Exopolysaccharide (EPS), pigments (pyoverdine, pyocyanin), alginate, LasA protease, LasB elastase and biofilm formation in *P. aeruginosa* are regulated by QS mechanism<sup>14,15</sup>. Bacterial biofilm is a group or communities of cells that is encased in self secreted extracellular polymeric substance consisting of proteins, polysaccharides and nucleic acids attached to a surface and involved in pathogenesis of infection and drug resistance<sup>16</sup>. Biofilm formation in MDR *P. aeruginosa* may reduce the effect of antibiotics and enhance the chances of emergence of drug resistance<sup>6</sup>. The treatment of infection with higher dose, overdose or prolonged uses of antibiotics is one of the major causes of drug resistance and biofilm formation. Comparatively the required doses of antibiotics or natural plant material for growth inhibition (antibacterial activity) are generally higher than required for interruption in QS cascade; therefore, regulation of virulence factors and biofilm formation by AQS mechanism at lower doses may be an effective solution of drug resistance and biofilm formation. The complications related to antibiotic therapy such as serious side effect in patients and chances of increasing drug resistance force the scientific community to investigate and explore the alternatives of antibiotics<sup>17-19</sup>.

The medicinal plant extracts/phytoconstituents exhibit antibacterial and Anti-Quorum Sensing (AQS) activity may be explored as an effective alternative of antibiotic therapy. Ginger (*Zingiber officinale* Roscoe.) family Zingiberaceae, has been used in traditional system of medicine for management of many diseases. It has been widely used for the management of nausea, diabetes, fever, cough, loss of appetite, indigestion, abdominal pain, intestinal colic, inflammation, infertility insomnia, and urinary tract infections and also exhibit antibacterial, anticoagulation, antipyretic, anti-obesity, anti-tumorigenic, AQS and anti-biofilm activity<sup>20-25</sup>. The major phytochemicals with diverse bioactivity including gingerol, shogaol and paradol as phenolic compounds; zingiberene and bisabolene as vola-

tile sesquiterpenes; and curcumene and citral as monoterpenoids<sup>26</sup>, may be an effective alternative of synthetic compound or antibiotics against antibiotic resistant micro-organism<sup>27,28</sup>. We have already determined the antibacterial activity and Minimum Inhibitory Concentrations (MIC) of methanol extract of ginger against reference PA01 and CIs 2,3,4 in our previous reported study<sup>29</sup>. In the present research work the AQS and anti-biofilm potential of methanol extract of ginger were investigated at sub-MICs against multidrug-resistant clinical isolates of *P. aeruginosa*.

## Materials and Methods

### Plant material collection and extract preparation

The fresh ginger (*Zingiber officinale*) rhizomes were collected and authenticated from Central Council for Research in Ayurvedic Sciences-Regional Ayurveda Research Institute, Jhansi (Uttar Pradesh), India. The air dried and powdered rhizomes were subjected for extraction with methanol (100%)<sup>29</sup>. Briefly 10 gm of rhizome powder were soaked in methanol solvent and mixed for 3 successive days at room temperature. The methanol extract was filtered through Whatman filter paper and centrifuged at 3000 rpm for 10 min. Supernatant was concentrated with rotary evaporator. The methanol extract was aseptically stored in sterile air tight bottle at 4°C for further experiment.

### Collection of clinical isolates and reference strains

Clinical isolates of *P. aeruginosa* were collected from Department of Microbiology, Sarojini Naidu Medical College, Agra (Uttar Pradesh), India. Reference strain (PA01) of *P. aeruginosa* and biosensor strain, *Chromobacterium violaceum* (*C. violaceum*) ATCC 12472 (CV12472) were collected from Department of Agricultural Microbiology, Aligarh Muslim University, Aligarh (Uttar Pradesh), India. *P. aeruginosa* were sub-cultured and maintained on Luria-Bertani (LB) broth at 37°C. *C. violaceum* (CV12472) was maintained on nutrient broth at 28°C.

### Screening of antibacterial activity and determination of MIC

In the previous step of our research study, agar well diffusion method was used for screening of antibacterial activity of crude ginger methanolic extract against Clinical Isolates (CIs) and reference strain of *P. aeruginosa* and 96 well microdilution method was used to determine MICs<sup>29</sup> as per the guideline of Clinical and Laboratory Standard Institute<sup>30</sup>.

### Screening of preliminary AQS activity

The preliminary AQS activity of ginger extract was confirmed qualitatively as inhibition in quorum sensing controlled violacein pigment formation in biosensor CV12472 using standard agar well diffusion method<sup>31</sup> with slight modification. Mueller Hinton agar plate was prepared with 100 µl of freshly grown and aseptically diluted bacterial suspension (inoculum size of 2.5×10<sup>6</sup> CFU/ml). The wells of approximate 6-8 mm diameter

were made sterile cub borer and bottom of wells were sealed using soft agar. 50  $\mu$ l of ginger methanolic extract (500 mg/ml) was aseptically poured in wells and plate was incubated at 28°C for 18-20 hr. A well for negative control was made with DMSO (dimethyl sulfoxide). The diameter of inhibition zone around the well representing violacein pigment inhibition (mm) was measured to determine the AQS activity<sup>32</sup>.

#### Determination of AQS and anti-biofilm activity against *P. aeruginosa*

**Assay for swarming motility:** The effect of ginger extract (at sub- MICs) on swarming motility was determined using the swarm plate method. The swarm plate was prepared using sterile 8 gm/l of nutrient broth added with 5 gm/l of glucose and 0.5% of bacto agar was aseptically mixed to solidify the media. The 50  $\mu$ l of sub-MICs of ginger extract was aseptically added in the swarm agar. The agar was poured in sterile plates and plates were kept at room temperature to solidify<sup>33</sup>. The swarm plates were point inoculated with freshly grown *P. aeruginosa* culture and incubated at 37°C for 18-24 hr. Swarm diameter (mm) (turbid zone form the point of inoculation) with treated and untreated *P. aeruginosa* culture (as control) was noted to calculate percentage inhibition in swarming motility<sup>32</sup>.

#### Assay for pyocyanin formation

The chloroform extraction method<sup>34</sup> was used to determine the effect of ginger extract on pyocyanin pigment formation. Freshly grown bacterial culture (5 ml) were mixed with 150  $\mu$ l of sub-MICs of ginger extract and incubated at 37°C for 18-24 hr. A total volume of 3 ml of chloroform were added to the culture supernatant and mixed vigorously. The chloroform layer was transferred in sterile tube and further 1 ml of 0.2 M HCl (hydrochloric acid) was added and solution was centrifuged (8,000 rpm for 10-12 min at room temperature). Absorbance of HCL layer (pink to deep red colour) in treated and untreated culture (as control) was noted at 520 nm to calculate percentage inhibition in pyocyanin formation<sup>32</sup>.

#### Extraction and estimation of EPS

The bacterial isolates were grown in LB broth containing 150  $\mu$ l of sub-MICs of ginger extract at 37°C for 18-24 hr and subjected for centrifugation after incubation. Supernatant was filtered and transferred aseptically in fresh sterile tubes. Three volume of 100% chilled ethanol was added and the tubes were incubated at 4°C for 24 hr to precipitate extraction and quantification of (EPS)<sup>35</sup>. The phenol (5%)-sulphuric acid (concentrated)<sup>36</sup> was further used for quantification of EPS by measuring the sugar. Percentage inhibition in EPS was calculated by measuring the absorbance of treated and untreated culture (as control) at 490 nm<sup>32</sup>.

#### Assay for biofilm formation

The anti-biofilm potential of ginger extract was determined using microtitre plate assay<sup>37</sup> with slight

modification. Biofilm formation in the wells of microtitre plate were allowed in the presence of ginger extract (at different sub-MICs) at 37°C for 18-24 hr. After incubation, the wells were drained properly and washed three times with sterile Phosphate-Buffered Saline (PBS). The biofilm in the wells was stained with 150  $\mu$ l of crystal violet solution (0.1%) and plate was incubated at room temperature for 5-10 min. After incubation, the wells were washed with sterile PBS to remove excess dye and plate was kept for air drying at room temperature. The dye taken by biofilm cells was solubilized with 150  $\mu$ l of glacial acetic acid (33% v/v). Absorbance of treated and untreated culture (as control) was recorded at 570 nm to calculate percentage inhibition in biofilm formation<sup>32</sup>.

#### Statistical analysis

The experiments were performed in triplicate and the mean  $\pm$  standard deviation was calculated. The data were further statistically analyzed with One way ANOVA and level of significance were expressed as P value (p<0.001, p<0.005, p< 0.01 and p<0.05 were denoted with \*\*\*\*, \*\*\*, \*\* and \* respectively) when inhibition in PA01 was compared to clinical isolates. Comparison between inhibitions by sub-MICs with in strain/clinical isolates was made and based on Tukey's HSD post hoc test and the mean difference was expressed as significant at the 0.05 level. All the statistical analyses were performed in IBM SPSS statistics 20 (version 20.0. Armonk, NY: IBM Corporation).

## Results

The preliminary AQS potential of ginger extract was confirmed qualitatively as inhibition in violacein pigment formation using agar well diffusion method against *C. violaceum* CV12472 (Figure 1). As an extension of our previous reported study, the sub-MICs (1/2,1/4,1/8 MIC) of previously determined MIC of ginger extract were made (Table 1) and subjected to determine anti-quorum and anti-biofilm activity against CIs and reference PA01 of *P. aeruginosa*. The effect of ginger extract on growth of CIs and PA01 of *P. aeruginosa* at selected sub-MICs were determined using spectroscopic growth curve analysis and observed that




Figure 1. Inhibition in violacein pigment as anti-quorum sensing activity of crude methanolic extract of ginger against *C. violaceum* CV12472. A) methanol extract of ginger; B) negative control.

Table 1. MIC and sub-MICs of ginger extract against clinical isolates (CI) 2, 3, 4, and PA01 of *P. aeruginosa*

| Strain/clinical isolates (CIs) | MIC and sub-MICs (mg/ml) of methanol extract of <i>Zingiber officinale</i> rhizomes |          |         |         |
|--------------------------------|---|----------|---------|---------|
|                                | MIC   | sub-MICs |         |         |
|                                |   | 1/2 MIC  | 1/4 MIC | 1/8 MIC |
| CI 2                           | 15.62   | 7.81     | 3.90    | 1.95    |
| CI 3                           | 31.25   | 15.62    | 7.81    | 3.90    |
| CI 4                           | 31.25   | 15.62    | 7.81    | 3.90    |
| PA01                           | 7.81  | 3.90     | 1.95    | 0.97    |

growth of tested isolates were not affected by mentioned sub-MICs (data not shown). The concentration dependent reduction in swarming motility was observed in tested clinical isolates and PA01 by ginger extract. The maximum reduction in swarming motility was found against PA01 (51.38%) at 1/2 MIC. It was approximately similar against CI2, CI3 and CI4 at 1/2 MIC. The reduction in all tested isolates were slightly similar at 1/4 MIC (32.04-33.8%) and at 1/8 MIC (17.79-20.6 %) which showed that maximum reduction was possible at 1/2 MIC and approximately similar at the concentration of less than 1/2 MIC (Figure 2A). The ginger extract reduced the pyocyanin pigment

formation against the tested isolates within the range (47.61-57.92%) at 1/2 MIC, (28.44-36.16%) at 1/4 MIC and (19.92-23.70%) at 1/8 MIC and maximum against CI3 (57.92%) (Figure 2B). The dose dependent reduction (%) in EPS was observed and the maximum reduction was found against PA01 (63.29%) followed by CI3 (43.66%) and CI2 (41.22%) (Figure 2C). The ginger showed maximum reduction in biofilm formation against CI2 (64.37%), followed by PA01 (60.36%), CI3 (58.89%) and CI4 (51.60%) at 1/2 MIC, while maximum reduction at 1/4 MIC was found against PA01 (46.28%) followed by CI2 (44.23%), CI4 (33.13%) and CI3 (28.61%) (Figure 2D).




Figure 2. Inhibition of the formation of various virulence factors (A), Inhibition in swarming motility (B), Inhibition in pyocyanin formation (C) Inhibition in exopolysaccharide (EPS), and (D) Biofilm Inhibition. Each bar represents the mean value of three independent replicates, and the error bar shows the standard deviation. \*\*\*\*p<0.001, \*\*\*p<0.005, \*\*p<0.01, \*p<0.05, NS: not significant when the reference PA01 strain was compared to CI 2, 3, and 4. Values sharing a common letter (a, b, c) were not significant (at p<0.05) when comparison was made between inhibitions by sub-MICs with in strain/clinical isolates.




Figure 3. Inhibition (%) of quorum-sensing-regulated virulence factors and biofilm formation in the (A) reference strain PA01, (B) CI 2, (C) CI 3, and (D) CI 4 of *P. aeruginosa* by sub-MICs of the ginger extract. Each bar represents the mean value of three independent replicates, and the error bar shows the standard deviation.

## Discussion

Traditionally plant component/extract has been used for the treatment of microbial infections. From the past several decades' growth inhibition mechanism is focused as one of the major targets for controlling microbial infections but in present scenario scientific community established that targeting QS mechanism through AQS approach may be a better solution for reduction in drug resistance and safe treatment of microbial infection at lower doses of antibiotics/plant component. In the present study, AQS and anti-biofilm potential of ginger was evaluated against MDR *P. aeruginosa* and observed the dose dependent reduction in the various virulence factors including swarming motility, pyocyanin pigment, exopolysaccharides and biofilm. Preliminary AQS activity of ginger extract was investigated and confirmed qualitatively as inhibition of violacein pigment formation against a biosensor strain, CV12472 (Figure 1) which showed similarity with previously reported findings with different plants including *Eucalyptus globulus*<sup>32</sup>, *Syzygium aromaticum*<sup>38,39</sup>, *Terminalia bellerica*<sup>40</sup>, *Xylopiya aethiopica* and *Monodora myristica*<sup>41</sup>, *Acacia nilotica*, *Cinnamomum verum* and *Punica grantum*<sup>39</sup>. The growth of *P. aeruginosa* was not affected by the tested sub-MICs of ginger extract which was in accordance of previous study reported Kim and Park in 2013. Pyocyanin pigment, associated in cystic fibrosis *P. aeruginosa* infections, is a well characterized QS regulated viru-

lence factor. The concentration dependent reduction (%) in pyocyanin pigment formation was found with ginger extract against all tested CIs and reference PA01 (Figure 2B). Previously similar reduction in pyocyanin pigment formation with plant extracts was reported<sup>40,42</sup>. The components of biofilm matrix keep the microbial cell together closely and serve as a barrier against antibacterial compounds to enhance its protection and drug resistance ability<sup>42</sup>. The biofilm formation is a complex and multifactor dependent process in which EPS and swarming motility plays a key role for its formation in *P. aeruginosa*<sup>43,44</sup>. Bacteria having ability of biofilm formation generally exhibit antibiotic resistance due to the formation of EPS as an essential component of biofilm maturation<sup>45</sup>. Therefore, it was hypothesized that the regulation in biofilm formation may be achieved by controlling the EPS formation and swarming motility through interference in quorum sensing mechanism using plant extracts and consequently bacterial drug resistance may be regulated. Concentration dependent inhibition (%) in swarming motility was observed against CIs and reference strain PA01 (Figure 2A); similar reduction with plant extract has been previously reported<sup>31,39</sup>.

In the present study concentration-based reduction (%) in EPS (Figure 2C) (for controlling biofilm formation) along with the reduction (%) in biofilm (Figure 2D) were found with ginger at sub-MICs in both the CIs and reference PA01. The effectiveness of ginger

extract for reduction in biofilm formation (%) against *P. aeruginosa* reference strain was also reported<sup>23,46</sup>. Yahya *et al* in 2013 reported the reduction in biofilm formation using different concentration of ginger ethanolic extract against *P. aeruginosa*<sup>47</sup>. Miari *et al* in 2020 reported the anti-biofilm activity of ginger ethanol extract at 1, 3 and 5% and found 42.74% inhibition at 5%, while biofilm inhibitory activity was not observed at rest of any tested concentration below 5% (1% and 3%) against reference strain of *P. aeruginosa* which showed the minimum anti-biofilm inhibitory concentration (5%) without affecting growth<sup>48</sup>. Similarly, we also observed the maximum biofilm inhibition (%) without affecting growth at concentration of 1/2 MIC (Table 1) and below which (at 1/4 MIC and 1/8 MIC) percentage reduction in biofilm was found comparatively low. Kim and Park in 2013 reported the 39-56% reduction in biofilm formation in *P. aeruginosa* (PA14) using different concentration of ginger extract (1, 5, and 10%). Reduction in EPS formation by 1% ginger extract in *P. aeruginosa* (PA14) has been also reported by Kim and Park in 2013. Akbari *et al* in 2023 reported the inhibition in biofilm formation with aqueous ethanolic extract of ginger against *P. aeruginosa* depending on the concentration of extract<sup>49</sup>. Results of present study showed that the limit in reduction (%) of tested virulence factors and biofilm formation and required doses of ginger extract for observed reduction were isolates/strain dependent. The present study suggests that formation of biofilm may be regulated by controlling the EPS and swarming motility as both are the important factors for biofilm formation. Inhibition in different QS regulated virulence factors and biofilm formation was observed with plant extract in previously reported research studies<sup>32,38,39,42,50-52</sup>.

The present study revealed the potential of ginger for reduction of QS regulated virulence factors and biofilm formation (Figures 3A-3D), which may be an effective approach for controlling bacterial infection as well as drug resistance. Although AQS and anti-biofilm potential of ginger extract has been revealed through the present study yet molecular investigation is required to determine the actual site of action for QSI mechanism.

### Conclusion

The findings of present study suggest that the methanol extract of ginger rhizome, has an effective AQS and anti-biofilm activity at low dose (sub-MIC) for inhibition of various QS regulated virulence factors and biofilm formation; therefore, having potential for regulation in bacterial pathogenesis. Ginger may be explored as a weapon for the safe and effective treatment of infections caused by MDR *P. aeruginosa* as well as controlling the chances of emergence of drug resistance.

### Acknowledgement

The authors thank to the Central Council for Research in Ayurvedic Sciences-Regional Ayurveda Research Institute, Jhansi (Uttar Pradesh), India for providing and authenticating the ginger (*Zingiber officinale*) rhizomes used in this research work. The authors gratefully thank the Department of Microbiology, Sarojini Naidu Medical College, Agra (Uttar Pradesh), India for providing the CIs of *P. aeruginosa* and the Department of Agricultural Microbiology, Aligarh Muslim University, Aligarh (Uttar Pradesh), India for providing *Chromobacterium violaceum* ATCC 12472 (CV12472) and the reference strain (PA01) for this study. For providing instrument/equipment facility for experiments, authors thank the Innovation centre, Bundelkhand University, Jhansi (Uttar Pradesh), India.

Funding: Nil.

### Conflict of Interest

There are no conflicts of interest.

### References

- Boyle EM, Ainsworth JR, Levin AV, Campbell AN, Watkinson M. Ophthalmic *Pseudomonas* infection in infancy. Arch Dis Child Fetal Neonatal Ed 2001;85(2): F139-40.
- Chastre J, Fragon J-Y. Ventilator-associated pneumonia. Am J Respir Crit Care Med 2002;165(7):867-903.
- Rossolini GM, Mantengoli E. Treatment and control of severe infections caused by multiresistant *Pseudomonas aeruginosa*. Clin Microbiol Infect 2005;11(Suppl 4):17-32.
- Gomez MI, Prince A. Opportunistic infections in lung diseases; *Pseudomonas* infections in cystic fibrosis. Curr Opin Pharmacol 2007;7(3):244-251.
- Bassetti M, Vena A, Croxatto A, Righi E, Guery B. How to manage *Pseudomonas aeruginosa* infections. Drugs Context 2018;7:212527.
- Adonizio A, Kong KF, Mathee K. Inhibition of quorum sensing-controlled virulence factor production in *Pseudomonas aeruginosa* by South Florida plant extracts. Antimicrob Agents Chemother 2008;52(1):198-203.
- Hentzer M, Givskov M, Parsek MR. Targeting quorum sensing for treatment of chronic bacterial biofilm infections. Lab Med 2002a;33:295-306.
- Dandekar AA, Chugani S, Greenberg EP. Bacterial quorum sensing and metabolic incentives to cooperate. Science 2012;338(6104):264-6.
- Schuster M, Greenberg EP. A network of networks: quorum sensing gene regulation in *Pseudomonas aeruginosa*. Int J Med Microbiol 2006;296(2-3):73-81.
- de Kievit TR. Quorum sensing in *Pseudomonas aeruginosa* biofilms. Environ Microbiol 2009;11(2):279-88.
- Favre-Bonté S, Köhler T, Van Delden C. Biofilm formation by *Pseudomonas aeruginosa*: role of the C4-HSL cell-to-cell signal and inhibition by azithromycin. J Antimicrob Chemother 2003;52(4):598-604.

12. Venturi V. Regulation of quorum sensing in *Pseudomonas*. *FEMS Microbiol Rev* 2006;30(2):274-91.
13. Pesci EC, Milbank JB, Pearson JP, McKnight S, Kende SA, Greenberg EP, et al. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* 1999;96(20):11229-11234.
14. Whitehead NA, Barnard AM, Slater H, Simpson NJ, Salmond GP. Quorum sensing in Gram-negative bacteria. *FEMS Microbiol Rev* 2001;25(4):365-404.
15. Lyczak JB, Cannon CL, Pier GB. Lung infections associated with cystic fibrosis. *Clin Microbiol Rev* 2002;15(2):194-222.
16. Vetrivel A, Ramasamy M, Vetrivel P, Natchimuthu S, Arunachalam S, Kim GS, et al. *Pseudomonas aeruginosa* biofilm formation and its control. *Biologics* 2021;1(3): 312-36.
17. Livermore DM. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare?, *Clin Infect Dis* 2002;34 (5):634-40.
18. Flamm RK, Weaver MK, Thornsberry C, Jones ME, Karlowsky JA, Sahm DF. Factors associated with relative rates of antibiotic resistance in *Pseudomonas aeruginosa* isolates tested in clinical laboratories in the United States from 1999 to 2002. *Antimicrob Agents Chemother* 2004;48(7):2431-6.
19. Fankham A, Kuete V, Voukeng I, Kuiate J, Page J. Antibacterial activities of selected Cameroonian species and their synergistic effects with antibiotics against multidrug resistant phenotypes. *BMC Complem Alter Med* 2011;11:104.
20. Nurtjahja-Tjendraputra E, Ammit AJ, Roufogalis BD, Tran VH, Duke CC. Effective anti-platelet and COX-1 enzyme inhibitors from pungent constituents of ginger. *Thromb Res* 2003;111(4-5):259-65.
21. Chrubasik S, Pittler MH, Roufogalis BD. *Zingiberis rhizoma*: A comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine* 2005;12(9):684-701.
22. Vattem DA, Mihalik K, Crixell SH, McLean RJ. Dietary phytochemicals as quorum sensing inhibitors. *Fitoterapia* 2007;78(4):302-10.
23. Kim HS, Park HD. Ginger extract inhibits biofilm formation by *Pseudomonas aeruginosa* PA14. *PLoS One* 2013;8(9):e76106.
24. Asfour HZ. Anti-quorum sensing natural compounds. *J Microsc Ultrastruct* 2018;6(1):1-10.
25. Unuofin JO, Masuku NP, Paimo OK, Lebelo SL. Ginger from Farmyard to Town: Nutritional and Pharmacological Applications. *Front Pharmacol* 2021;12:779352.
26. Ali BH, Blunden G, Tanira MO, Nemmar A. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food Chem Toxicol.* 2008;46(2):409-20.
27. Wail EA, Emad MA. Antibacterial activity of Ginger (*Zingiber Officinale* Rosc.) rhizome: A mini review. *Int J Pharmacogn Chinese Med* 2018;2(4):000142.
28. Beristain-Bauza DCS, Hernández-Carranza P, Soledad Cid-Pérez T, Ávila-Sosa R, Israel Ruiz-López I, Ochoa-Velasco CE. Antimicrobial activity of ginger (*Zingiber Officinale*) and its application in food products. *Food Reviews International* 2019;35:407-26.
29. Sagar PK, Sharma P, Singh R. Antibacterial efficacy of different combinations of clove, eucalyptus, ginger, and selected antibiotics against clinical isolates of *Pseudomonas aeruginosa*. *Ayu* 2020;41(2):123-9.
30. Clinical and Laboratory Standards Institute. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: approved standard. 9th ed. Wayne: CLSI; 2012.
31. Zahin M, Hasan S, Aqil F, Khan MS, Husain FM, Ahmad I. Screening of certain medicinal plants from India for their anti-quorum sensing activity. *Indian J Exp Biol* 2010;48(12):1219-24.
32. Sagar PK, Sharma P, Singh R. Inhibition of quorum sensing regulated virulence factors and biofilm formation by eucalyptus globulus against multidrug-resistant *Pseudomonas aeruginosa*. *J Pharmacopuncture.* 2022;25(1): 37-45.
33. Bala A, Kumar R, Harjai K. Inhibition of quorum sensing in *Pseudomonas aeruginosa* by azithromycin and its effectiveness in urinary tract infections. *J Med Microbiol* 2011;60 (Pt 3):300-06.
34. Essar DW, Eberly L, Hadero A, Crawford IP. Identification and characterization of genes for a second anthranilate synthase in *Pseudomonas aeruginosa*: interchangeability of the two anthranilate synthases and evolutionary implications. *J Bacteriol* 1990;172(2):884-900.
35. Huston AL, Methe B, Deming JW. Purification, characterization, and sequencing of an extracellular cold-active aminopeptidase produced by marine psychrophile *Collwellia psychrerythraea* strain 34H. *Appl Environ Microbiol* 2004;70(6):3321-8.
36. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric method for determination of sugars and related substances. *Anal Chem* 1956;28(3):350-6.
37. O'Toole GA, Kolter R. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* 1998;28(3):449-61.
38. Sagar PK, Singh R. Anti-quorum sensing activity of clove (*Syzygium aromaticum*) bud extract and its combined efficacy with ceftazidime against multidrug-resistant *Pseudomonas aeruginosa*. *Research Journal of Biotechnology* 2021;16(3):75-82.
39. Samreen, Qais FA, Ahmad I. Anti-quorum sensing and biofilm inhibitory effect of some medicinal plants against gram-negative bacterial pathogens: in vitro and in silico investigations. *Heliyon* 2022;8(10):e11113.
40. Sankar Ganesh P, Ravishankar Rai V. Attenuation of quorum-sensing-dependent virulence factors and biofilm formation by medicinal plants against antibiotic resistant *Pseudomonas aeruginosa*. *J Tradit Complement Med* 2017;8(1):170-7.
41. Tamfu AN, Ceylan O, Kucukaydin S, Ozturk M, Duru ME, Dinica RM. Antibiofilm and enzyme inhibitory potentials of two annonaceous food spices, african pepper

- (*Xylopia aethiopica*) and african nutmeg (*Monodora myristica*). *Foods* 2020;9(12):1768.
42. El-Sayed NR, Samir R, Jamil M Abdel-Hafez L, Ramadan MA. Olive leaf extract modulates quorum sensing genes and biofilm formation in multi-drug resistant *Pseudomonas aeruginosa*. *Antibiotics (Basel)* 2020;9(9):526.
  43. Watnick PI, Kolter R. Steps in the development of a *Vibrio cholerae* El Tor biofilm. *Mol Microbiol* 1999;34(3):586-95.
  44. Köhler T, Curty LK, Barja F, van Delden C, Pechère JC. Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. *J Bacteriol* 2000;182(21):5990-6.
  45. Chen L, Wen YM. The role of bacterial biofilm in persistent infections and control strategies. *Int J Oral Sci* 2011;3(2):66-73.
  46. Nikolić M, Vasić S, Đurđević j, Stefanović O, Čomić L. Antibacterial and anti-biofilm activity of ginger (*Zingiber officinale* (roscoe)) ethanolic extract. *Kragujevac J Sci* 2014;36:129-36.
  47. Yahya MFZR, Saifuddin NFHA, Hamid UMA. *Zingiber officinale* ethanolic extract inhibits formation of *Pseudomonas aeruginosa* biofilm. *International Journal of Pharmacy and Biological Sciences* 2013;3(1):46-54.
  48. Miari M, Rasheed SS, Haidar Ahmad N, Itani D, Abou Fayad A, Matar GM. Natural products and polysorbates: Potential inhibitors of biofilm formation in *Pseudomonas aeruginosa*. *J Infect Dev Ctries* 2020;14(6):580-8.
  49. Akbari S, Didar Z, Vazifedoost M, Hajirostamloo B, Mohtashami M. Antibiofilm activity of ginger (*Zingiber officinale*) extracts in vitro and food model. *Journal of Food Processing and Preservation* 2023: Article ID 5134332.
  50. Bjarnsholt T, Jensen PØ, Rasmussen TB, Christophersen L, Calum H, Hentzer M, et al. Garlic blocks quorum sensing and promotes rapid clearing of pulmonary *Pseudomonas aeruginosa* infections. *Microbiology* 2005;151(Pt 12):3873-880.
  51. Khan MS, Zahin M, Hasan S, Husain FM, Ahmad I. Inhibition of quorum sensing regulated bacterial functions by plant essential oils with special reference to clove oil. *Lett Appl Microbiol* 2009;49(3):354-60.
  52. Husain FM, Ahmad I, Asif M, Tahseen Q. Influence of clove oil on certain quorum-sensing-regulated functions and biofilm of *Pseudomonas aeruginosa* and *Aeromonas hydrophila*. *J Biosci* 2013;38(5):835-44.