



Monocytes and Macrophages as Unique Cellular Compartments Governing Non-Alcoholic Fatty Liver Disease and Inflammation

Ghazale Hemmatian¹, Davoud Rostamzadeh², Kaveh Baghaei³ and Mahdi Shabani^{1*}

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Medicinal Plants Research Center, Yasuj University of Medical Sciences, Yasuj, Iran

3. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Abstract

Non-Alcoholic Fatty Liver Disease (NAFLD) is a spectrum of liver diseases from simple steatosis to the most severe form of hepatocellular carcinoma. Liver injuries resulting from various factors, including viral infections, alcohol consumption, and metabolic disorders, trigger the activation of resident immune cells and the recruitment of circulating immune cells to the liver. This chronic inflammatory environment leads to tissue damage and the progression of liver fibrosis. Macrophages are highly versatile immune cells that play a dual role in fibrosis: they contribute to the progression of fibrosis (M1 and Ly6c^{high} macrophages) and its resolution (M2 and Ly6c^{low} macrophages). M1 macrophages and those with high surface expression of Ly6C exhibit pro-inflammatory characteristics, while M2 macrophages and myeloid cells with low expression of Ly6C mitigate inflammation and inhibit fibrosis progression. Environmental stimuli influence the complex mechanisms hepatic macrophages regulate the fibrosis they encounter. Kupffer cells initiate the inflammatory cascade and recruit monocyte-derived macrophages, which modulate the propagation of fibrosis and promote fibrinolysis. Additionally, hepatic macrophages interact with other cell types through exosomes, facilitating the transfer of cellular components that influence the outcome of liver fibrosis. In this review, the critical role of macrophages in inflammation-induced fibrosis and tissue restoration is discussed.

Keywords: Carcinoma, Exosomes, Fibrinolysis, Hepatocellular, Kupffer cells, Liver cirrhosis, Liver neoplasms, Non-alcoholic fatty liver disease

To cite this article: Hemmatian Gh, Rostamzadeh D, Baghaei K, Shabani M. Monocytes and Macrophages as Unique Cellular Compartments Governing Non-Alcoholic Fatty Liver Disease and Inflammation. Avicenna J Med Biotech 2025;17(2):98-105.

Introduction

Non-Alcoholic Fatty Liver Disease (NAFLD) is a long-lasting liver disease characterized by the accumulation of huge amounts of fat in the liver which can lead to liver-related death, especially among steatonecrosis and fibrosis-afflicted patients. NAFLD may progress to a more severe liver condition termed fibrosis¹. Liver fibrosis is the accumulation of extracellular matrix in response to repeated liver damage. It takes place when the capacity of the liver to substitute hepatocytes is overwhelmed and fibrillar collagen is abundantly accumulated². Cellular mechanisms contributing to the progression of liver fibrosis include the activation of Hepatic Stellate Cells (HSCs) and their trans-differentiation into myofibroblast-like cells, which produce large amounts of extracellular collagen and facilitate fibrosis progression³.

Immune cells and liver fibrosis

Macrophages play contradictory roles in regulating liver fibrosis; they promote it through the production of IL-1 β and TNF- α by M1 macrophages, while M2 macrophages can reverse liver fibrosis⁴⁻⁷. Different subsets of T cells also influence the outcome of fibrosis: Th1 cells exhibit pro-inflammatory characteristics, Th2 cells release IL-13 and IL-4 to promote fibrogenesis, and Th17 cells produce IL-17, which activates HSCs and macrophages. In cases of chronic hepatic damage, despite the early apoptosis-inducing effects of Natural Killer (NK) cells on HSCs, these cells are ultimately considered to promote fibrosis. The changes in the number of immune cells in the context of liver fibrosis determine the persistence of inflammation. A decrease in NK cells and CD8⁺ cytotoxic T cells, along with an

* Corresponding author:
Mahdi Shabani, Ph.D.,
Department of Immunology,
School of Medicine, Shahid
Beheshti University of Medical
Sciences, Tehran, Iran
Tel: +98 21 22439970
Fax: +98 21 22439970
E-mail:
msshabani@sbmu.ac.ir
Received: 20 Jul 2024
Accepted: 30 Nov 2024

increase in CD4⁺ memory T cells, underscores the necessity of inflammation for the continuation of liver fibrosis⁸. Hepatocytes secrete IL-8, CXCL1, and CXCL2 to promote the recruitment of neutrophils, which is followed by increased myeloperoxidase activity, reactive oxygen species production, and the formation of neutrophil extracellular traps, all contributing to the progression of inflammation. Dendritic cells, which perform antigen presentation to naïve T cells, play paradoxical roles in NASH, they release TNF- α , MCP-1, and IL-6 during the initial phases of liver damage to promote inflammation, but in the regression phases of liver injury, they facilitate fibrosis regression. Natural Killer T (NKT) cells, a component of acquired immunity, drive fibrosis progression through the secretion of IFN- γ , IL-4, IL-15, VCAM-1, CXCL16, and osteopontin, as well as through an overactive Hedgehog pathway. B cells exhibit dual roles in regulating liver inflammation by presenting oxidative stress-derived epitopes to CD4⁺ helper T cells and releasing IL-10 to inhibit CD8⁺ T cell-mediated inflammation. However, they also produce anti-oxidative stress epitopes, such as IgG, to mitigate inflammation⁹. Dysregulation of matrix metalloproteinases, which promote matrix degradation, and tissue inhibitors of matrix metalloproteinases govern extracellular matrix remodeling, leading to the deposition of extracellular matrix components. Another contributing mechanism to liver inflammation and fibrosis is the disruption of immune tolerance mediated by regulatory T cells and dendritic cells, which instigate liver fibrosis¹⁰. Despite the complex involvement of the immune system in liver inflammation, this study aims to briefly discuss the roles of macrophages. Kupffer cells, the resident macrophages of the liver, are responsible for maintaining liver homeostasis through the phagocytosis of apoptotic cells and pathogens. They mediate the liver's homeostatic state through immune regulation *via* IL-10 production, phagocytosis of gut-derived invading substrates and facilitating hepatic repair and regeneration. The production of TGF- β and PDGF by Kupffer cells leads to HSC activation. Furthermore, Kupffer cell-derived IL-6 and TNF- α contribute to the persistence of inflammation. Additionally, Kupffer cells are activated through the sensing of apoptotic hepatocyte signaling and NF- κ B activation, promoting inflammation^{4,5,7,11-13}.

Liver macrophages, including Kupffer cells and monocyte-derived macrophages, constitute the hepatic macrophage pool. However, in response to chronic liver injury and inflammation, these macrophages adopt ambivalent functions, including those of M1 and Ly6C^{high} (lymphocyte antigen 6 complex, locus C1) monocytes. Following the cessation of the inflammatory response, macrophages acquire M2 and Ly6C^{low} phenotypes to promote tissue repair¹⁴. In mice, Ly6C^{high} monocytes, which correspond to CD14^{high} CD16^{low} human monocytes, are considered pro-inflammatory.

In contrast, Ly6C^{low} monocytes and their human counterparts, as CD14^{low} CD16^{high} monocytes, patrol along blood vessels and promote fibrosis regression due to their elevated expression of matrix metalloproteinases, phagocytosis-associated genes, and growth factors, extending beyond the M1 and M2 phenotypes¹⁵. Traditionally, macrophage polarization states were categorized as M1 or M2. M1 macrophages (classically activated macrophages) are pro-inflammatory and contribute to tissue injury, while M2 macrophages (non-classically activated macrophages) are involved in tissue remodeling and efferocytosis, thereby inhibiting inflammation¹⁶.

HSCs are non-parenchymal liver cells residing in the sub-endothelial space of Disse (also known as the perisinusoidal space) and interact with liver macrophages upon activation¹⁷. Following we provide an update on the critical role of macrophages in liver fibrosis, as well as their interactions with other liver cells.

NAFLD spectrum and prevalence

NAFLD is a spectrum of liver disorders, ranging from Non-Alcoholic Fatty Liver (NAFL), and Non-Alcoholic Steatohepatitis (NASH) to liver fibrosis, cirrhosis, and Hepatocellular Carcinoma (HCC). Due to the strong association between NAFLD and type 2 diabetes, hyperlipidemia, obesity, and hypertension, NAFLD is considered a metabolic syndrome¹⁸. NAFL is distinguished when the liver contains $\geq 5\%$ steatosis without the presence of fibrosis. NAFL is usually benign and without significant risk of mortality. NAFL can progress to NASH (non-alcoholic steatohepatitis), characterized by at least 5% hepatic steatosis, accompanied by hepatocyte ballooning and liver inflammation. NASH may be followed by more advanced complications including liver fibrosis, cirrhosis, and HCC. When defining NAFLD, it must be considered that people who consume abundant quantities of alcohol must be excluded from this spectrum^{19,20}.

NAFLD imposes a great health burden on societies worldwide. Based on a research study conducted by Ge *et al*, it was estimated that NAFLD prevalence would increase dramatically from 1990 to 2017 with the highest prevalence observed in many territories including East Asia and the Middle East²¹. Furthermore, NAFLD prevalence and mortality highly increased in the United States^{22,23}. These findings underscore the critical need for healthcare professionals to implement thorough measurements in managing this disease.

Inflammation initiation

Diverse innate and adaptive immune cells are implicated in liver fibrosis. The innate immune cells of the liver serve as the first line of defense against invading injuries. However, when chronic damage takes place, Endoplasmic Reticulum (ER) stress (unfolded protein response), Reactive Oxygen Species (ROS), and lipid overload especially free fatty acids, induce liver in-

inflammation initiation. A disruption occurring in the equilibrium of ROS production and antioxidant mechanisms causes molecular damage and leads to fibrosis development²⁴⁻²⁷. The main instigators of liver inflammation are attributed to several factors including Free Fatty Acids (FFAs) and bacterial-derived lipopolysaccharide that activate immune cells through interaction with Damage-Associated Molecular Patterns (DAMPs) and Pathogen-Associated Molecular Patterns (PAMPs), respectively leading to initiation of liver inflammation. Gut dysbiosis and adipose tissue dysfunction accelerate inflammation progression^{28,29}. Furthermore, leukocytes recruitment (macrophages, neutrophils, T cells) trigger tissue damage in NAFLD³⁰⁻³². Hepatocyte cell death is mediated by oxidative stress and FFAs deposition which are a result of mitochondrial dysfunction observed in NAFLD^{28,30}. In addition, an inequity between reactive nitrogen species and reactive oxygen species leads to fibrosis³³.

The liver resident macrophages termed Kupffer cells are the most abundant resident macrophages in this organ. They are hepatic sentinels against exogenous and endogenous danger signals. They are characterized as F4/80⁺, CD11b^{intermediate}, CD45⁺, and C-type lectin domain family 4, member f (CLEC4F) positive cells. They become activated after recognizing Lipopolysaccharide (LPS) by toll-like receptor 4 (TLR4) and transmitting signals through myeloid differentiation primary response protein 88 (MyD88) dependent or MyD88 independent pathways, thereby initiating the

inflammatory cascade and activation of HSCs^{14,24}. Macrophages acquire the pro-inflammatory phenotype by ROS production, phagocytosis, elevated expression of adhesion molecules, and cytokines to decrease danger signals. Macrophages' anti-inflammatory phenotype secrete TGF- β , and IL-10 and accomplish apoptotic cell phagocytosis to decrease inflammation. However, as a result of persistent inflammation, inflammatory monocytes secrete TGF- β to mediate the activation of fibroblasts. M2 macrophages are fibrogenic subsets driving fibrosis perpetuation via profibrotic mediators and matrix metalloproteinases prevention (Figure 1)³⁴⁻³⁶.

In addition, other metabolites such as adipokines, and cholesterol through scavenger receptors A and CD36, and DAMPs *e.g.*, ATP, mitochondrial DNA, and high mobility group box 1 originated from injured hepatocytes and free fatty acids *via* TLRs induce the activation of Kupffer cells^{37,38}. Activated Kupffer cells produce transforming growth factor- β 1 (TGF- β 1), platelet-derived growth factor (PDGF), interleukin-1 (IL-1), monocyte chemoattractant protein 1 (MCP1), IL-12 and tumor necrosis factor- α (TNF- α) essential for HSCs proliferation and recruitment. Kupffer cells gelatinases and chitinase expression also induce collagen type I production and HSCs activation, respectively. Kupffer cells-derived TGF- β 1 also stimulates the differentiation and activation of fibroblasts³⁹. In addition, activated Kupffer cells also serve as a source of IL-6 and inducible Nitric Oxide Synthase




Figure 1. Monocyte and macrophage of the liver in the progression and restoration of fibrosis.

Kupffer cells become activated after recognizing LPS, adipokines, cholesterol, DAMPs, and free fatty acids. Activated Kupffer cells produce TGF- β 1, PDGF, IL-1, MCP1, IL-12, and TNF- α essential for HSCs proliferation and recruitment. Kupffer cells-derived TGF- β 1 also stimulates the differentiation and activation of fibroblasts. The reciprocal interplay between hepatic macrophages and HSCs perpetuates fibrosis. Indeed, Kupffer cells assist HSCs activation, survival, and pro-fibrogenic phenotype maintenance through TGF- β , oncostatin M, PDGF, TNF- α , IL-1 β , TNFSF12A, and EGFR ligand secretion. In turn, activated HSCs release CCL2, hyaluronan, IL-6, and TGF- β which are essential for monocytes recruitment, activation, and polarization. Furthermore, Kupffer cells-derived CCL2 and CCL5 contribute to Ly6C^{high} monocytes infiltration to the liver that promotes hepatic inflammation. In normal conditions, Ly6C^{low} monocytes have been shown to contribute to fibrosis regression by matrix metalloproteinases MMP9 and MMP12 expression and induction of phagocytosis of cellular debris namely efferocytosis. M2 macrophages are stimulated by IL-13, IL-10, and IL-4. Matrix metalloproteinases expression by M2 macrophages facilitates wound healing. TGF- β , arginase-1 (Arg1), and IL-10 produced by M2 macrophages contribute to anti-inflammatory characteristics.

(iNOS) expression. Indeed, iNOS stimulates Nitric Oxide (NO) production which together with ROS produce Reactive Nitrogen Species (RNS) and stimulates the HSCs activation⁴⁰. The reciprocal interplay between hepatic macrophages and HSCs perpetuates fibrosis. Indeed, Kupffer cells assist HSCs activation, survival, and pro-fibrogenic phenotype maintenance through TGF- β , oncostatin M, PDGF, TNF- α , IL-1 β , TNF superfamily member 12 (TNFSF12A) and Epidermal Growth Factor Receptor (EGFR) ligand secretion. In turn, activated HSCs release C-C motif chemokine ligand 2 (CCL2), hyaluronan, IL-6, and TGF- β which are essential for monocytes recruitment, activation, and polarization. Activated HSCs also contribute to inflammation persistence through secreting CCL21 and IL-1 β . CCL21 helps to recruit immune cells to sites of inflammation. In addition, NLRP3 inflammasome activation leads to IL-1 β secretion that assists inflammation perpetuation^{41,42}. This positive feedback loop preserves hepatic inflammation and fibrosis⁴³. Furthermore, Kupffer cells-derived CCL2 and CCL5 contribute to Ly6c^{high} monocytes infiltration to the liver that promotes hepatic inflammation (Figure 1)¹³.

Inflammatory monocytes/macrophages in fibrosis progression

Kupffer cells TLR signaling, and activation cause upregulated levels of CCL2 or MCP1 production. These mediators induce the recruitment of Monocyte-derived Macrophages (MoMFs) which are characterized by Ly6c, C-X3-C motif chemokine receptor 1 (CX3CR1), C-C chemokine receptor 2 (CCR2), and CD11b expression and accomplish macrophage replenishment upon injury. Based on their Ly6c expression, Ly6c^{high} monocytes equal to CD14^{high} CD16^{low} human monocytes contribute to inflammation propagation^{44,45}. A research study conducted by Baeck *et al* deduced CCL2 as a chemokine indispensable for Ly6c^{high} pro-fibrogenic monocytes infiltration that hinders fibrosis resolution⁴⁶. This subset is crucial for determining fibrosis fate; because they are characterized as the main proliferative macrophage subset expressing iNOS⁴⁷.

Macrophages adopt different phenotypes depending on their microenvironmental stimuli. Interferon- γ (IFN- γ), LPS, and IL-12 induce M1 macrophages polarization which is characterized by the production of iNOS, TNF- α , IL-1 β , IL-6, IL-12, IL-8, and IL-15 and participate in pathogen elimination and inflammation propagation. These macrophages, also known as classically activated macrophages, are distinguished by their cell surface expression markers, including CD86, CD80, TLR4, TLR2, Interleukin-1 Receptor (IL-1R), and major histocompatibility complex class II (MHC-II). In addition, they secrete chemokines like CCL5, CCL2, CCL3, CCL4, C-X-C motif ligand 8 (CXCL8), CXCL9, CXCL10 and CXCL11. M1 macrophages also trigger HSC activation by secreting TGF- β and facilitating fibrosis progression⁴⁸⁻⁵¹.

Restorative monocytes/macrophages in fibrosis regression

Macrophages are highly plastic immune cells capable of both promoting and reversing liver inflammation and fibrosis. Ly6c^{high} monocytes can polarize to Ly6c^{low} monocytes equal to CD14^{low} CD16^{high} human monocytes that are implicated in fibrosis regression⁴⁴. These monocytes have been shown to contribute to fibrosis regression by matrix metalloproteinases MMP9 and MMP12 expression and induction of phagocytosis of cellular debris namely efferocytosis. Ly6c^{low} monocytes exhibit anti-fibrotic characteristics corroborated by the observation that they express CD47/Macrophage Migration Inhibitory Factor (MIF) and Insulin-like Growth Factor 1 (IGF1)¹⁵.

M2 macrophages also known as non-classically activated macrophages are stimulated by IL-13, IL-10, and IL-4. M2 macrophages express CD163, CD206, dectin-1, scavenger receptors and CCR2. Matrix metalloproteinases expression by M2 macrophages facilitates wound healing. TGF- β , arginase-1 (Arg1), and IL-10 produced by M2 macrophages contribute to anti-inflammatory characteristics⁴⁸. In addition, in a research study conducted by Wan and colleagues, it was reported that M2-polarized Kupffer cells induce M1 Kupffer cells apoptosis to inhibit inflammation⁵². In general, M2 macrophages are anti-inflammatory due to IL-13, IL-4, IL-10, and TGF- β secretion, and assist wound healing⁵⁰. Due to the high versatility of M2 macrophages, they are further subdivided into M2a, M2b, M2c, and M2d. M2a and M2b macrophages regulate immune responses, and M2c macrophages promote wound healing and tissue remodeling. In addition, this subgroup of macrophages suppresses immune responses. M2d macrophages are implicated in angiogenesis and assist tumor development⁴⁹. Due to the aforementioned dual paradoxical characteristics of macrophages in exacerbating and ameliorating liver inflammation and fibrosis, an approach that targets inflammatory macrophages or promotes the polarization of restorative macrophages holds great potential in mitigating the disease.

Macrophages therapeutic targeting

There have been several strategies proposed to mitigate liver fibrosis. The primary therapeutic approaches focus on reducing the activation of Kupffer cells through modulation of the gut-liver axis. These strategies include preventing monocyte recruitment via CCL2-CCR2 inhibition, directing macrophage polarization from a pro-inflammatory to a restorative phenotype, and the transplantation of autologous monocyte-derived macrophages as well as bone marrow-derived macrophages. All of these strategies aim to alleviate liver inflammation and fibrosis^{6,13,45,53,54}.

Macrophages are compelling targets for liver inflammation, as they play crucial roles in the progression and regression of liver fibrosis. Therefore, any agent that targets macrophages may have the potential

to alleviate liver fibrosis. A summary of some pharmacological interventions is presented in table 1.

Exosomes-mediated cellular crosstalk in liver fibrosis

The interaction between monocytes/macrophages and other cell types plays a crucial role in determining the fate of fibrosis. Liver sinusoidal endothelial cells express delta-like canonical Notch ligand 4 (DLL4), and the signaling pathways involving Notch, TGF- β , and LXR promote the trans differentiation of monocytes into Kupffer cells, which are recruited to the injured liver to expedite fibrosis progression. Furthermore, HSC-derived IL-1 β and TNF- α enhance the differentiation of monocytes into Kupffer cells. Macrophages release PDGF, TGF- β , and ROS, which induce HSC activation. In turn, HSCs release CCL2 to promote the infiltration and activation of macrophages. This creates a positive feedback loop in which macrophages support HSC survival, while HSCs inhibit macrophage phagocytosis, thereby perpetuating inflammation^{43,65}. Additionally, macrophages facilitate Endothelial-to-Mesenchymal Transition (EndMT) in endothelial cells, a process characterized by increased invasiveness and migration, cytoskeletal reorganization, and loss of cellular adhesion, all of which contribute to HSC activation. Endothelial cells upregulate Vascular Adhesion Protein-1 (VAP1) and the integrin $\alpha 4$ subunit VLA-4, further stimulating monocyte recruitment. Macrophages, in turn, mediate endothelial disruption

^{43,65}. Despite the complex interplay between macrophages and various immune cells that govern inflammation, the following section aims to elucidate macrophage-mediated communication through exosomes. Reason: Improved clarity, vocabulary, and technical accuracy while correcting grammatical and punctuation errors.

Due to the importance of secreted exosomes in cell-cell communications, in the following section, the effect of exosomes derived from either macrophages or other cellular sources on liver fibrosis outcome will be explained. Exosomes are types of extracellular vesicles ranging in size from 30-100 nm which contain multiple types of cargo including proteins, lipids, and nucleic acids, and assist intercellular communication. Exosome cargoes (e.g., miRNAs) influence the recipient cells and may stimulate or inhibit liver fibrosis⁶⁶⁻⁶⁸. Chen *et al* showed that exosomes derived from THP-1 (human leukemia monocytic cell line) macrophages treated with LPS altered miR-103-3p and could therefore stimulate HSCs activation and proliferation⁶⁹. Jiayi *et al* observed that LPS-mediated stimulation of THP-1 cells promotes exosomes secretion rich in miR-155-5p which elevates oxidative stress, collagen synthesis as well as migration and proliferation of HSCs⁷⁰. These studies corroborate macrophage-derived exosomes in the progression of hepatic fibrosis. In another study by Wan *et al*, they showed that miR-411-5p in M2 macro-

Table 1. Therapeutic agents targeting macrophages

Pharmacological agent	Type of impact on macrophages	Result	Reference
GR-MD-02	Direct	Macrophage Galectin 3 antagonist, Improvement in liver stiffness	(55)
Cenicriviroc	Direct	Monocyte CCR2/CCR5 antagonist, Improvement in liver fibrosis	(56)
Anti-CD163-IgG-dexamethasone	Direct	Corticoid delivery to macrophages, Diminution of hepatocyte ballooning, inflammation and fibrosis	(57)
Curcumin	Direct	NF- κ B inhibitor, Hepatic inflammation, fat accumulation, fibrosis, and insulin resistance decrease	(6)
1,25-dihydroxy-vitamin D3	Direct	NF- κ B inhibitor, Hepatic inflammation, fat accumulation, fibrosis, and insulin resistance decrease	(6)
Emricasan	Direct	Pan-caspase inhibitor	(58)
Antibiotics, probiotics, fecal microbiota transfer, sequestration of bile acids	Direct	Kupffer cells inhibitor	(6)
Liraglutide	Indirect	Glucagon-like peptide 1 agonist, NASH improvement	(59)
Pioglitazone	Indirect	Peroxisome proliferator-activated receptor γ agonist, Inflammation, hepatocyte ballooning, and fibrosis improvement	(60)
Obeticholic acid	Indirect	Farnesoid X receptor agonist, NASH resolution, improvement in liver fibrosis	(61)
Pentoxifylline	Indirect	Phosphodiesterase Inhibitor, Improvement in steatosis, inflammation, hepatocyte ballooning and fibrosis	(62)
Selonsertib	Indirect	Apoptosis signal-regulating kinase 1 antagonist, Fibrosis alleviation	(63)
Elafibranor	Indirect	Peroxisome proliferator-activated receptor α and δ agonist, NASH improvement	(64)
Pirfenidone	Indirect	Decreasing effects on TGF- β	(58)
Sorafenib	Indirect	Multi-kinase inhibitor inhibits the TGF- β /Smad3 pathway and targets the Raf/Mek/Erk pathway	(58)

phages-derived exosomes impeded HSC activation through Calmodulin-regulated spectrin-associated protein 1 (CAMSAP1) inhibition and the ensuing down-regulation of collagen and α -smooth muscle actin (α -SMA) in HSCs ⁷¹. Liu and coworkers reported that miR-192-5p expression in exosomes originated from lipotoxic hepatocytes stimulated pro-inflammatory M1 macrophages polarization *via* Rictor (rapamycin-insensitive companion of mammalian target of rapamycin)/Akt/FoxO1 (forkhead box transcription factor O1) pathway ⁷². It has been shown that lipids-mediated death receptor 5 induction in hepatocytes assists extracellular vesicles secretion which in turn instigates pro-inflammatory macrophage polarization ⁷³. Lysosomal dysfunction caused by cholesterol mediates hepatocytes exosomal miR-122-5p release that induces M1 macrophages polarization and increases inflammation ⁷⁴.

According to Chen and co-workers research study, LPS-induced macrophage activation and exosomal miR-500 release led to HSCs activation and proliferation in CCl₄ (carbon tetrachloride)-induced mice liver fibrosis model through MFN2 targeting ⁷⁵. Benbow and colleagues also confirmed that exosomes derived from activated HSCs stimulated macrophages' inflammatory TNF- α and IL-6 expression and migration ⁷⁶. Moreover, miR-148a in mesenchymal stem cells-derived exosomes facilitates macrophage phenotype switching from the pro-inflammatory M1 to anti-inflammatory M2 subset by Kruppel-Like Factor 6 (KLF6) and signal transducer and activator of transcription 3 (STAT3) inhibition transcription factors ⁷⁷. In conclusion, the source of exosome-producing cells in different micro-environments causes the alteration of miRNAs contents which induces fibrosis induction or restoration.

Conclusion

Macrophages are regarded as the most critical immune cells in the regulation of liver fibrosis. Different subsets of pro-inflammatory and anti-inflammatory macrophages play opposing roles in the progression and resolution of liver fibrosis. Therefore, strategies that inhibit the harmful effects of inflammatory monocytes, while promoting restorative monocytes or targeting their intercellular communication, could be promising avenues for alleviating the disease. However, further in-depth studies are necessary to identify the various monocyte subsets and the cellular pathways involved in fibrosis.

Acknowledgement

This work was financially supported by Shahid Beheshti University of Medical Sciences (grant number: 43008191).

Conflict of Interest

Authors declare no conflict of interest.

References

- Matteoni CA, Younossi ZM, Gramlich T, Boparai N, Liu YC, McCullough AJ. Nonalcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 1999;116(6):1413-9.
- Battaller R, Brenner DA. Liver fibrosis. *J Clin Invest* 2005;115(2):209-18.
- Yi Q, Yang J, Wu Y, Wang Y, Cao Q, Wen W. Immune microenvironment changes of liver cirrhosis: emerging role of mesenchymal stromal cells. *Front Immunol* 2023; 14:1204524.
- Li W, Chang N, Li L. Heterogeneity and function of kupffer cells in liver injury. *Front Immunol* 2022;13:940867.
- Li W, Yang Y, Yang L, Chang N, Li L. Monocyte-derived Kupffer cells dominate in the Kupffer cell pool during liver injury. *Cell Rep* 2023;42(10):113164.
- van der Heide D, Weiskirchen R, Bansal R. Therapeutic targeting of hepatic macrophages for the treatment of liver diseases. *Front Immunol* 2019;10:2852.
- Dixon LJ, Barnes M, Tang H, Pritchard MT, Nagy LE. Kupffer cells in the liver. *Compr Physiol* 2013;3(2):785.
- Sun R, Xiang Z, Wu B. T cells and liver fibrosis. *Potal Hypertension & Cirrhosis* 2022;1(02):125-32.
- Gregory SN, Perati SR, Brown ZJ. Alteration in immune function in patients with fatty liver disease. *Hepatoma Res* 2022;8(31):10-20517.
- Robinson MW, Harmon C, O'Farrelly C. Liver immunology and its role in inflammation and homeostasis. *Cell Mol Immunol* 2016;13(3):267-76.
- Chen J, Deng X, Liu Y, Tan Q, Huang G, Che Q, et al. Kupffer cells in non-alcoholic fatty liver disease: friend or foe? *Int J Biol Sci* 2020;16(13):2367.
- Ma YY, Yang MQ, He ZG, Wei Q, Li JY. The biological function of kupffer cells in liver disease. London: Intech-Open; 2017 May 10.
- Ju C, Tacke F. Hepatic macrophages in homeostasis and liver diseases: from pathogenesis to novel therapeutic strategies. *Cell Mol Immunol* 2016;13(3):316-27.
- Cheng D, Chai J, Wang H, Fu L, Peng S, Ni X. Hepatic macrophages: Key players in the development and progression of liver fibrosis. *Liver Int* 2021;41(10):2279-94.
- Ramachandran P, Pellicoro A, Vernon MA, Boulter L, Aucott RL, Ali A, et al. Differential Ly-6C expression identifies the recruited macrophage phenotype, which orchestrates the regression of murine liver fibrosis. *Proc Natl Acad Sci USA* 2012;109(46):E3186-E95.
- Alisi A, Carpino G, Oliveira FL, Panera N, Nobili V, Gaudio E. The role of tissue macrophage-mediated inflammation on NAFLD pathogenesis and its clinical implications. *Mediators Inflamm* 2017;2017(1):8162421.
- Gupta G, Khadem F, Uzonna JE. Role of hepatic stellate cell (HSC)-derived cytokines in hepatic inflammation and immunity. *Cytokine* 2019;124:154542.
- Younossi ZM, Henry L, Bush H, Mishra A. Clinical and economic burden of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *Clin Liver Dis* 2018;22(1): 1-10.

19. Sharma P, Arora A. Clinical presentation of alcoholic liver disease and non-alcoholic fatty liver disease: spectrum and diagnosis. *Transl Gastroenterol Hepatol* 2020;5.
20. De A, Duseja A. Natural history of simple steatosis or nonalcoholic fatty liver. *J Clin Exp Hepatol* 2020;10(3): 255-62.
21. Ge X, Zheng L, Wang M, Du Y, Jiang J. Prevalence trends in non-alcoholic fatty liver disease at the global, regional and national levels, 1990–2017: a population-based observational study. *BMJ Open* 2020;10(8): e036663.
22. Estes C, Razavi H, Loomba R, Younossi Z, Sanyal AJ. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 2018;67(1):123-33.
23. Younossi ZM, Stepanova M, Younossi Y, Golabi P, Mishra A, Rafiq N, Henry L. Epidemiology of chronic liver diseases in the USA in the past three decades. *Gut* 2020;69(3):564-8.
24. Hernandez-Gea V, Friedman SL. Pathogenesis of liver fibrosis. *Annu Rev Pathol* 2011;6:425-56.
25. Mooli RGR, Mukhi D, Ramakrishnan SK. Oxidative stress and redox signaling in the pathophysiology of liver diseases. *Compr Physiol* 2022;12(2):3167-92.
26. Zhang J, Guo J, Yang N, Huang Y, Hu T, Rao C. Endoplasmic reticulum stress-mediated cell death in liver injury. *Cell Death Dis* 2022;13(12):1051.
27. Wobser H, Dorn C, Weiss TS, Amann T, Bollheimer C, Büttner R, et al. Lipid accumulation in hepatocytes induces fibrogenic activation of hepatic stellate cells. *Cell Res* 2009;19(8):996-1005.
28. Song C, Long X, He J, Huang Y. Recent evaluation about inflammatory mechanisms in nonalcoholic fatty liver disease. *Front Pharmacol* 2023;14:1081334.
29. Ganz M, Szabo G. Immune and inflammatory pathways in NASH. *Hepatol Int* 2013;7 Suppl 2(Suppl 2):771-81.
30. Guo Z, Wu Q, Xie P, Wang J, Lv W. Immunomodulation in non-alcoholic fatty liver disease: exploring mechanisms and applications. *Front Immunol* 2024;15: 1336493.
31. Wang H, Mehal W, Nagy LE, Rotman Y. Immunological mechanisms and therapeutic targets of fatty liver diseases. *Cell Mol Immunol* 2021;18(1):73-91.
32. Peiseler M, Tacke F. Inflammatory mechanisms underlying nonalcoholic steatohepatitis and the transition to hepatocellular carcinoma. *Cancers (Basel)* 2021;13(4): 730.
33. Ortiz-Lopez N, Fuenzalida C, Dufeu MS, Pinto-Leon A, Escobar A, Poniachik J, et al. The immune response as a therapeutic target in non-alcoholic fatty liver disease. *Front Immunol* 2022;13:954869.
34. Wermuth PJ, Jimenez SA. The significance of macrophage polarization subtypes for animal models of tissue fibrosis and human fibrotic diseases. *Clin Transl Med* 2015;4:1-19.
35. Austermann J, Roth J, Barczyk-Kahlert K. The good and the bad: Monocytes' and macrophages' diverse functions in inflammation. *Cells* 2022;11(12):1979.
36. Ma W-T, Gao F, Gu K, Chen D-K. The role of monocytes and macrophages in autoimmune diseases: a comprehensive review. *Front Immunol* 2019;10:1140.
37. Kazankov K, Jørgensen SMD, Thomsen KL, Møller HJ, Vilstrup H, George J, et al. The role of macrophages in nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *Nat Rev Gastroenterol Hepatol* 2019;16(3): 145-59.
38. Wen Y, Lambrecht J, Ju C, Tacke F. Hepatic macrophages in liver homeostasis and diseases-diversity, plasticity and therapeutic opportunities. *Cell Mol Immunol* 2021;18(1):45-56.
39. Nguyen-Lefebvre AT, Horuzsko A. Kupffer cell metabolism and function. *J Enzymol Metab* 2015;1(1).
40. Iwakiri Y. Nitric oxide in liver fibrosis: The role of inducible nitric oxide synthase. *Clin Mol Hepatol* 2015;21 (4):319.
41. Garbuzenko DV. Pathophysiological mechanisms of hepatic stellate cells activation in liver fibrosis. *World J Clin Cases* 2022;10(12):3662.
42. Poulsen KL, Cajigas-Du Ross CK, Chaney JK, Nagy LE. Role of the chemokine system in liver fibrosis: a narrative review. *Dig Med Res* 2022;5:30.
43. Matsuda M, Seki E. Hepatic Stellate Cell-Macrophage Crosstalk in Liver Fibrosis and Carcinogenesis. *Semin Liver Dis* 2020 Aug;40(3):307-320.
44. Kaur KK, Allahbadia G, Singh M. How do we apply advances in knowledge of Hepatic Macrophages in treating Liver Diseases especially non alcoholic fatty liver disease (NAFLD), non alcoholic steatohepatitis (NASH), with the increasing incidence of Diabetes-A Systematic Review. *EC Endocrinology and Metabolic Research* published in2020.
45. Binatti E, Gerussi A, Barisani D, Invernizzi P. The role of macrophages in liver fibrosis: new therapeutic opportunities. *Int J Mol Sci* 2022;23(12):6649.
46. Baeck C, Wei X, Bartneck M, Fech V, Heymann F, Gassler N, et al. Pharmacological inhibition of the chemokine C-C motif chemokine ligand 2 (monocyte chemoattractant protein 1) accelerates liver fibrosis regression by suppressing Ly-6C+ macrophage infiltration in mice. *Hepatology* 2014;59(3):1060-72.
47. Pellicoro A, Ramachandran P, Iredale JP, Fallowfield JA. Liver fibrosis and repair: immune regulation of wound healing in a solid organ. *Nat Rev Immunol* 2014;14(3): 181-94.
48. Alisi A, Carpino G, Oliveira FL, Panera N, Nobili V, Gaudio E. The role of tissue macrophage-mediated inflammation on NAFLD pathogenesis and its clinical implications. *Mediators Inflamm* 2017;2017.
49. Wang C, Ma C, Gong L, Guo Y, Fu K, Zhang Y, et al. Macrophage polarization and its role in liver disease. *Front Immunol* 2021;12:803037.
50. Liang W, Huang X, Shi J. Macrophages serve as bidirectional regulators and potential therapeutic targets for liver fibrosis. *Cell Biochem Biophys* 2023;81(4):659-71.
51. Tugal D, Liao X, Jain MK. Transcriptional control of macrophage polarization. *Arterioscler Thromb Vasc Biol* 2013;33(6):1135-44.

52. Wan J, Benkdane M, Teixeira-Clerc F, Bonnafous S, Louvet A, Lafdil F, et al. M2 Kupffer cells promote M1 Kupffer cell apoptosis: a protective mechanism against alcoholic and nonalcoholic fatty liver disease. *Hepatology* 2014;59(1):130-42.
53. Ping D, Peng Y, Hu X, Liu C. Macrophage cytottherapy on liver cirrhosis. *Front Pharmacol* 2023;14:1265935.
54. Liu R, Scimeca M, Sun Q, Melino G, Mauriello A, Shao C, et al. Harnessing metabolism of hepatic macrophages to aid liver regeneration. *Cell Death Dis* 2023;14(8):574.
55. Harrison S, Marri S, Chalasani N, Kohli R, Aronstein W, Thompson G, et al. Randomised clinical study: GRMD-02, a galectin-3 inhibitor, vs. placebo in patients having non-alcoholic steatohepatitis with advanced fibrosis. *Aliment Pharmacol Ther* 2016;44(11-12):1183-98.
56. Friedman SL, Ratzliff V, Harrison SA, Abdelmalek MF, Aithal GP, Caballeria J, et al. A randomized, placebo-controlled trial of cenicriviroc for treatment of nonalcoholic steatohepatitis with fibrosis. *Hepatology* 2018;67(5):1754-67.
57. Svendsen P, Graversen JH, Etzerodt A, Hager H, Røge R, Grønbaek H, et al. Antibody-directed glucocorticoid targeting to CD163 in M2-type macrophages attenuates fructose-induced liver inflammatory changes. *Mol Ther Methods Clin Dev* 2017;4:50-61.
58. Shan L, Wang F, Zhai D, Meng X, Liu J, Lv X. New drugs for hepatic fibrosis. *Front Pharmacol* 2022;13: 874408.
59. Armstrong MJ, Gaunt P, Aithal GP, Barton D, Hull D, Parker R, et al. Liraglutide safety and efficacy in patients with non-alcoholic steatohepatitis (LEAN): a multicentre, double-blind, randomised, placebo-controlled phase 2 study. *Lancet* 2016;387(10019):679-90.
60. Boettcher E, Csako G, Pucino F, Wesley R, Loomba R. Meta-analysis: pioglitazone improves liver histology and fibrosis in patients with non-alcoholic steatohepatitis. *Aliment Pharmacol Ther* 2012;35(1):66-75.
61. Neuschwander-Tetri BA, Loomba R, Sanyal AJ, Lavine JE, Van Natta ML, Abdelmalek MF, et al. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 2015;385(9972):956-65.
62. Zeng T, Zhang C-L, Zhao X-L, Xie K-Q. Pentoxifylline for the treatment of nonalcoholic fatty liver disease: a meta-analysis of randomized double-blind, placebo-controlled studies. *Eur J Gastroenterol Hepatol* 2014;26(6): 646-53.
63. Loomba R, Lawitz E, Mantry PS, Jayakumar S, Caldwell SH, Arnold H, et al. The ASK1 inhibitor selonsertib in patients with nonalcoholic steatohepatitis: a randomized, phase 2 trial. *Hepatology* 2018;67(2):549-59.
64. Ratzliff V, Harrison SA, Francque S, Bedossa P, Leher P, Serfaty L, et al. Elafibranor, an agonist of the peroxisome proliferator-activated receptor- α and- δ , induces resolution of nonalcoholic steatohepatitis without fibrosis worsening. *Gastroenterology*. 2016;150(5):1147-59. e5.
65. Coelho I, Duarte N, Macedo MP, Penha-Gonçalves C. Insights into Macrophage/Monocyte-endothelial cell crosstalk in the liver: a role for trem-2. *J Clin Med* 2021;10(6):1248.
66. Wang C, Liu J, Yan Y, Tan Y. Role of exosomes in chronic liver disease development and their potential clinical applications. *J Immunol Res* 2022;2022.
67. Chen L, Brenner DA, Kisseleva T. Combatting fibrosis: exosome-based therapies in the regression of liver fibrosis. *Hepatol Commun* 2019;3(2):180-92.
68. Valencia K, Montuenga LM. Exosomes in liquid biopsy: the nanometric world in the pursuit of precision oncology. *Cancers (Basel)* 2021;13(9):2147.
69. Chen L, Yao X, Yao H, Ji Q, Ding G, Liu X. Exosomal miR-103-3p from LPS-activated THP-1 macrophage contributes to the activation of hepatic stellate cells. *FASEB J* 2020;34(4):5178-92.
70. Lin J, Lou A, Li X. [Lipopolysaccharide stimulates macrophages to secrete exosomes containing miR-155-5p to promote activation and migration of hepatic stellate cells]. *Nan Fang Yi Ke Da Xue Xue Bao* 2023 Jun 20;43(6):994-1001. Chinese.
71. Wan Z, Yang X, Liu X, Sun Y, Yu P, Xu F, Deng H. M2 macrophage-derived exosomal microRNA-411-5p impedes the activation of hepatic stellate cells by targeting CAMSAP1 in NASH model. *iScience*. 2022;25(7).
72. Liu X-L, Pan Q, Cao H-X, Xin F-Z, Zhao Z-H, Yang R-X, et al. Lipotoxic hepatocyte-derived exosomal miR-192-5p activates macrophages via Rictor/Akt/FoxO1 signaling in NAFLD. *Hepatology* 2020;72(2):454-69.
73. Hirsova P, Ibrahim SH, Krishnan A, Verma VK, Bronk SF, Werneburg NW, et al. Lipid-induced signaling causes release of inflammatory extracellular vesicles from hepatocytes. *Gastroenterology* 2016;150(4):956-67.
74. Zhao Z, Zhong L, Li P, He K, Qiu C, Zhao L, Gong J. Cholesterol impairs hepatocyte lysosomal function causing M1 polarization of macrophages via exosomal miR-122-5p. *Exp Cell Res* 2020;387(1):111738.
75. Chen L, Huang Y, Duan Z, Huang P, Yao H, Zhou Y, et al. Exosomal miR-500 derived from lipopolysaccharide-treated macrophage accelerates liver fibrosis by suppressing MFN2. *Front Cell Dev Biol* 2021;9:716209.
76. Benbow JH, Marrero E, McGee RM, Brandon-Warner E, Attal N, Feilen NA, et al. Hepatic stellate cell-derived exosomes modulate macrophage inflammatory response. *Exp Cell Res* 2021;405(1):112663.
77. Tian S, Zhou X, Zhang M, Cui L, Li B, Liu Y, et al. Mesenchymal stem cell-derived exosomes protect against liver fibrosis via delivering miR-148a to target KLF6/STAT3 pathway in macrophages. *Stem Cell Res Ther* 2022;13(1):330.