



Exploring the Role of hsa_circ_0052112 as a Potential Biomarker in Breast Cancer: Insights from Experimental and *In Silico* Analyses

Mahdi Alizadeh and Mahdiah Salimi *

Department of Medical Genetics, Institute of Medical Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

Abstract

Background: Circular RNAs (circRNAs) are important in tumorigenesis and cancer progression, highlighting their potential as biomarkers for diagnosis, prognosis, and treatment monitoring.

Methods: This study consists of experimental and *in silico* phases. In the experimental phase, the expression of hsa_circ_0052112 in tumor and blood samples from 40 breast cancer women was analyzed, compared to the control group using Sybr Green real-time RT-PCR followed by total RNA extraction and cDNA synthesis. Statistical analysis was performed using the *beta-actin* gene as a normalizer, compared to the normal control group as a fold change. In the *in silico* phase, interactions among circRNA, RNA-Binding Proteins (RBPs), and microRNAs (miRNAs) were investigated using Interactome Database. The miRcancer database was utilized to assess breast cancer-related miRNAs linked to hsa_circ_0052112. Target miRNAs were identified with TargetScan and filtered for relevance through DisGeNET. K-means clustering grouped genes by expression patterns, visualized in Cytoscape to illustrate circRNA-miRNA-mRNA relationships. Hub genes underwent pathway enrichment analysis using Reactome database to determine their functional significance.

Results: Data revealed a significant increase in hsa_circ_0052112 expression in both blood and tumour of breast cancer patients. This increase was especially pronounced in patients with estrogen, progesterone, and HER2 receptor positivity, as well as in advanced disease stages with lymph node involvement. Enrichment analysis of hub genes indicates their role in the PI3K/AKT signaling pathway.

Conclusion: hsa_circ_0052112 shows promise as a multifaceted biomarker for breast cancer, enhancing diagnosis and prognosis; while supporting personalized treatment strategies. Further clinical validation is necessary to confirm its utility.

Keywords: Breast neoplasms, Circular RNAs, hsa_circ_0052112, MicroRNAs, Prognosis

To cite this article: Alizadeh M, Salimi M. Exploring the Role of hsa_circ_0052112 as a Potential Biomarker in Breast Cancer: Insights from Experimental and *In Silico* Analyses. Avicenna J Med Biotech 2025;17(2):136-146.

* **Corresponding author:**
Mahdiah Salimi, Ph.D.,
Department of Medical Genetics,
Institute of Medical Biotechnology,
National Institute of Genetic
Engineering and Biotechnology
(NIGEB), Tehran, Iran
Tel: +98 9123341722, 44787382
Fax: +98 21 44787395
E-mail:
salimi@nigeb.ac.ir
Received: 5 Oct 2024
Accepted: 5 Mar 2025

Introduction

Heritable and dynamic changes occurring to the genome independently of DNA sequence are known as "epigenetics". Cancer was the first disease reported as an epigenetic changes-related disease ¹. The activation of oncogenes and/or the suppression of tumor suppressor genes are critical events always accompanied by epigenetic changes. These epigenetic changes include DNA methylation, micro-RNA and long non-coding RNA expression, and histone post-translational modifications ^{2,3}. Breast cancer development is a multistep and complex process that includes the synergistic crosstalk between epigenetic and genetic alterations,

which are influenced by an abundance of external and internal factors. Such factors include but are not limited to the cell's intrinsic nutrient supply, microenvironment, cellular stress, and external environmental exposures to endocrine disruptors or carcinogenic agents. The epigenetic changes in breast cancer development and progression are influenced by critical genes involved in apoptosis, proliferation, cell motility, invasion, and other processes ^{4,5}. Modification of epigenetic changes has been reported as a promising therapeutic strategy in breast cancer. Breast cancer is a heterogeneous type of malignancy that results from unique mo-

lecular changes in breast tissue ⁶.

Circular RNAs (circRNAs) are stable and evolutionarily conserved RNA regulators that can act as microRNA (miRNA) sponges, regulate alternative splicing mechanisms, and have an influential role in the expression of the encoded genes ⁷. CircRNAs have been shown to play essential roles in the cell and are one of the most crucial research focuses, particularly in cancer ⁸. Due to their significant roles in critical genetic pathways, particularly in oncology, two distinct therapeutic potentials have been identified for circRNAs: inhibiting carcinogenic circRNAs or restoring circRNAs that possess tumor suppressor functions ⁹. Since molecular phenotypes do not operate in isolation, but their interactions collectively carry out biological processes, it is essential to model these processes in living cells using networks ^{10,11}. It is anticipated that identifying circRNA-miRNA-mRNA connections will be crucial for explaining the molecular processes underlying numerous diseases, detecting biomarkers for early diagnosis, and expanding therapeutic options. A single miRNA can target hundreds of genes, while a single circRNA can act as a sponge for tens of miRNAs. Using bioinformatics data to simplify the circRNA-miRNA-mRNA interactions may shed light on *in vitro* and *in vivo* investigations ¹².

In this study, focus was on hsa_circ_0052112, a circular RNA derived from the *ZNF83* gene. This selection was based on circRNA expression profile analyses in breast cancer cells, which were conducted using a microarray assay in 2018. The expression levels of hsa_circ_0052112 in MCF-7 and MDA-MB-231 cells were measured using RT-qPCR, along with evaluations of the migration and invasion capabilities of breast cancer cells. The findings indicated that the expression of hsa_circ_0052112 was significantly higher in MDA-MB-231 cells compared to MCF-7 cells. Additionally, the overexpression of hsa_circ_0052112 promoted cell migration and invasion in breast cancer, whereas its downregulation suppressed these processes ¹³. As of now, no additional studies have been published on hsa_circ_0052112. Considering its significant implications and established relevance to breast cancer, researchers set out to build upon this research by investigating the expression of hsa_circ_0052112 in tumor and blood samples from breast cancer patients in comparison to normal controls. Additionally, the aim was to examine its expression across various clinicopathological groups for the first time.

The present research was designed in experimental and *in silico* phases. In the experimental phase, the expression of hsa_circ_0052112 in the tumor tissue and blood of breast cancer patients was investigated in comparison with the normal control group, and its association with the clinical characteristics of the disease was studied. In the *in silico* phase, the interaction of hsa_circ_0052112 with miRNA, RBPs, and target genes was investigated. Breast cancer remains a lead-

ing cause of cancer-related morbidity and mortality among women worldwide, necessitating the identification of novel biomarkers for improved diagnosis, prognosis, and treatment strategies. Circular RNAs have emerged as significant regulatory molecules in various biological processes, including tumorigenesis and cancer progression, highlighting their potential utility in oncology ¹⁴. This study specifically investigates the expression of hsa_circ_0052112 in breast cancer, aiming to elucidate its role as a biomarker. By analyzing hsa_circ_0052112 levels in tumor and blood samples from breast cancer patients and correlating these findings with clinical characteristics, researchers sought to establish a direct link between circRNA expression and breast cancer phenotypes. Furthermore, examination of the interactions between hsa_circ_0052112, RNA-binding proteins (RBPs), and microRNAs (miRNAs) aims to provide insights into the molecular mechanisms underlying its involvement in breast cancer. Ultimately, this research endeavors to contribute to the development of personalized treatment strategies by establishing hsa_circ_0052112 as a multifaceted biomarker with significant clinical relevance.

Materials and Methods

Experimental phase

Sampling: 40 blood and breast tissue (Tumor and Normal adjacent) samples were collected from breast cancer patients at Imam Khomeini Hospital in Tehran, Iran. A control blood group consisting of 40 samples was also collected from non-affected individuals. The inclusion criteria for the normal control group samples were female participants within the age range of the patients, devoid of any history of breast disease or malignancy in both themselves and their first-degree relatives. The inclusion criteria for the test group were patients who were diagnosed with ductal carcinoma of the breast and had not initiated any treatment at the time of sampling. The demographic and clinical data of the patients and controls have been included in table 1. Ethical considerations and informed consent paperwork had been acquired for this study (IR.MODARES.REC.1398.148).

RNA extraction: Peripheral blood samples (3-4 ml) were collected into EDTA tubes for plasma preparation. The tubes were centrifuged at 1000 *g* for 15 *min* within 30 *min* of collection. Subsequently, 1 ml of plasma was transferred to a 1.5-ml Eppendorf tube and centrifuged at 11,000 *g* for 10 *min* to remove any remaining cellular debris. Total RNA was extracted from plasma and breast tissue samples using the RNX-Plus solution (CinnaClone-Iran) following a multi-step protocol. The process began with the addition of RNX-Plus to lyse cells and release RNA, followed by the introduction of chloroform for phase separation, which involved vigorous shaking and centrifugation to isolate the RNA-containing aqueous phase. RNA was then

Table 1. Characteristics of breast cancer patients and controls

	Patient N (%)
Number	40
Age (years)	
Mean	45.9±11.6
Range	26-73
Stage at diagnosis	
Stage I	3 (7.5)
Stage II	17 (42.5%)
Stage III	14 (35%)
Stage IV	6 (15%)
Lymph node status	
N0	18 (45%)
N+	22(55%)
Distance metastasis	
yes	5 [1 bone, 4 lungs] (12.5%)
No	35 (87.5%)
Hormone receptor status (IHC)	
ER positive	24 (60%)
ER negative	16 (40%)
PR positive	21 (52.5%)
PR negative	19 (47.5%)
HER-2 status (IHC)	
+++	12 (30%)
Negative	24 (60%)
Triple-negative breast cancer	4 (10%)

precipitated by adding isopropanol or ethanol and incubating the mixture at -20°C. After precipitation, the RNA pellet was washed with 70% ethanol to eliminate impurities and air-dried, before being resuspended in RNase-free water or buffer. The concentration and purity of the extracted RNA were assessed using a NanoDrop ND-ONE spectrophotometer (Thermo-USA), measuring absorbance at 260 nm and calculating the 260/280 ratio to check for protein contamination.

cDNA synthesis: For cDNA synthesis, the YT4500 cDNA Synthesis Kit (Yekta Tajhiz Azma, Iran) was utilized. The process began with the preparation of a reaction mixture comprising extracted RNA, random hexamer primers from the kit, reverse transcriptase, dNTPs, and buffer components. This mixture was then incubated according to the manufacturer's protocol, which involved an initial primer annealing at 25°C for 10 min, followed by reverse transcription at 42°C for 60 min, and enzyme inactivation at 70°C for 5 min. The synthesized cDNA was subsequently stored at -20°C for future use in quantitative RT-PCR.

SYBR Green Quantitative real-time reverse transcription PCR: To quantify the expression levels of circRNA (Hsa_circ_0052112) and the internal control gene *β-actin*, SYBR Green real-time RT-PCR was utilized, allowing for the amplification and detection of specific DNA sequences in real-time. *β-actin* was selected due to its stable expression across various cell types and experimental conditions, demonstrated minimal varia-

bility in preliminary assessments compared to other housekeeping genes, and extensive literature validation, thereby ensuring our findings' robustness and reliability. Each PCR reaction contained a cDNA template, specific primers for hsa_circ_0052112, forward: AGAGGGCTTTATACAGGGCC; reverse: CCCTGA AAGTCAAGCATCCC, and *β-actin* (forward: CACCTTCTACAATGAGCTGCGTGTG; reverse: ATAGC ACAGCCTGGATAGCAACGTAC, SYBR Green dye, and PCR buffer. The thermal cycling conditions included an initial denaturation at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 30 s, annealing at 57°C for 30 s, and extension at 72°C for 30 s. Each reaction incorporated negative controls (lacking template DNA) to check for contamination and technical duplicates to ensure reproducibility. The relative expression of circRNA was normalized to *β-actin*, with primer efficiency evaluated using LinReg-PCR software to confirm approximately 100% efficiency. The comparative quantification of gene expression was conducted using the $2^{-\Delta\Delta CT}$ method, where ΔCT represents the difference in Cycle threshold (Ct) values between the target gene and *β-actin*. Expression changes were categorized as upregulated (two-fold or higher), normal (between 0.5 and 2-fold), or downregulated (0.5-fold or lower), ensuring accurate and reliable quantification of gene expression levels in the analyzed samples.

Statistical analyses

The acquired data were analyzed using Graph Pad Prism software version 8.2. Since the data showed a normal distribution, parametric tests were employed for statistical analysis. To compare circRNA expression between two groups and more than two groups, the T-test and Analysis of Variance (ANOVA) test were conducted. The significance level was set at $p < 0.05$. Results were reported as mean±standard deviation. Cohen's d-test was used for effect size calculation.

Insilco phase

Investigating RNA regulatory interactions using the interactome database: The Interactome site (<https://circinteractome.nia.nih.gov/>) was employed as a supply for investigating the interactions between circular RNA (circRNA), RNA-Binding Proteins (RBPs), and microRNAs (miRNAs) in this study. This database comprises experimentally validated interactions between these molecules, enabling researchers to find the convoluted relationships within the RNA regulatory network. By analyzing the interactions on the Interactome site, we gained insights into how circRNAs, RBPs, and miRNAs interact, which may indicate novel regulatory mechanisms and biological functions. This information was significant for realizing gene expression regulation, disease mechanisms, and identifying potential therapeutic targets in the study performed. Table 2 shows the characteristics of circular RNA taken from the interactome site.

Table 2. Characteristics of circular RNA taken from the interactome site

CircRNA ID	hsa_circ_0052112
Genomic length	5283 bp
Location	chr19:53158813-53164096
Spliced seq length	209 bp
Gene symbol	ZNF83

Investigating the interaction between RNA-binding proteins and circular RNA in post-transcriptional

Gene regulation: The correlation between RBPs and circRNAs is especially noteworthy in RNA biology. RBPs can bind to circRNAs, touching their stability, localization, and potential functions. This interaction may result in the formation of circRNA-protein complexes that can serve as regulators of gene expression or signaling pathways. Comprehension of the relationship between RBPs and circRNAs is crucial for clarifying the complex regulatory networks within cells and revealing the roles of circRNAs in various biological processes. RNA-binding protein sites matching flanking regions of circRNA, taken from the interactome site, are shown in table 3.

Exploring the regulatory interplay between microRNAs and circular RNAs using the interactome database

In this study, the interplay between miRNAs and circRNAs engaging the Interactome site to uncover the regulatory mechanisms within the RNA network was explored. miRNAs are small non-coding RNAs that regulate gene expression post-transcriptionally by binding to target mRNAs and regulating their translation or degradation. CircRNAs, alternatively, are manifesting as important gene expression regulators that can segregate miRNAs and modulate their activity. By evaluating the interactions between miRNAs and circRNAs on the Interactome site, researchers intended to reveal new regulatory pathways and potential therapeutic targets. Recognition of how circRNAs impact miRNA function can provide a significant understanding of the complicated dynamics of different RNA molecules and their impact on gene expression. This insight is important for understanding disease mecha-

nisms, identifying biomarkers, and developing targeted therapies for various disorders. Table 4 lists the miRNAs that are targeted by hsa_circ_0052112 taken from the interactome site.

Identification of breast cancer-related miRNAs targeted by circular RNA through enrichment analysis

Among miRNAs targeted by hsa_circ_0052112 (Table 4), enrichment was done using miRcancer database (<https://mircancer.ecu.edu/>), and miRNAs related to breast cancer were identified.

Visualization of circRNA_miRNA_mRNA relationships in a tripartite manner using cytoscape

In this research study, the target miRNA genes selected from the previous step were identified using the TargetScan database (<https://www.targetscan.org/>). Subsequently, genes related to breast cancer were filtered using DisGeNET. The breast cancer-related genes were represented as data points in a multidimensional space (string), with each dimension reflecting a specific gene expression value. K-means clustering was employed to group genes with similar expression patterns into clusters. The resulting clusters were visualized as a network with confidence of interaction score of 0.9, where genes were depicted as nodes and relationships between genes were shown as edges based on their expression similarities. This network analysis enabled the exploration of gene interactions and functional relationships within biological systems. Following the creation of the gene network in string format, the data was transferred to Cytoscape for further analysis. Gene hubs were identified, and the circRNA_miRNA_mRNA relationships were visualized using Cytoscape in a tripartite manner.

Exploring Hub genes enrichment by reactome



In this study, hub genes enrichment by the Reactome database (<https://reactome.org/>) involved the identification of key genes that hold significant importance in biological pathways or processes. Reactome, a curated database of biological pathways, offers detailed information on molecular events and their interrelationships across diverse cellular processes. Through hub gene enrichment analysis using Reactome, researchers can identify genes that serve as cen-

Table 3. RNA-binding protein sites matching flanking regions of circRNA hsa_circ_0052112

RBPs	Tag name	% identity	Alignment length	Mismatches	Gap openings	Tag start	Tag end	circRNA start	circRNA end	Upstream/downstream
AGO2	HHFCT_76827_cluster-12605_20_8_35	100.00	35	0	0	1	35	+65	+99	Downstream
AGO2	HHFCT_76828_cluster-12605_21_8_36	100.00	36	0	0	1	36	-902	-867	Upstream
EIF4A3	HHLE2_605111_eIF4AI II_rep2_605111_1_54	100.00	54	0	0	1	54	+210	+263	Downstream
EIF4A3	HHLE2_605114_eIF4AI II_rep2_605114_1_40	100.00	40	0	0	1	40	-485	-446	Upstream
PTB	HHFPT_135578_PTB_cluster-12605_68_19_47	100.00	47	0	0	1	47	-767	-721	Upstream

Table 4. List of miRNAs targeted by the TargetScan database for circular RNA hsa_circ_0052112

TargetScan miRNA predictions				
CircRNA Mirbase ID	CircRNA (Top) - miRNA (Bottom) pairing	Site type	CircRNA start	CircRNA end
hsa_circ_0052112 (5' ... 3') hsa-miR-142-5p (3' ... 5')	ACCCUGCACAGAGGGCUUUUAUAC UCAUCACGAAAGAU--GAAAUAC	7mer-1a	159	165
hsa_circ_0052112 (5' ... 3') hsa-miR-146b-3p (3' ... 5')	CACAGAGGGCUUUUAUACAGGGCC GGUCUUGACUCAGGUGUCCCGU AUUCCAAAGACUCAUGUUACAU	7mer-m8	165	171
hsa_circ_0052112 (5' ... 3') hsa-miR-194 (3' ... 5')	AGGUGUACCUCACGACAAUGU GAUGGCUCUCCUCAGGGAUGCU	7mer-1a	21	27
hsa_circ_0052112 (5' ... 3') hsa-miR-324-5p (3' ... 5')	UGUGGUUACGGGAUCCCCUACGC UACAGGGCCGUGAUGUUGGAGAA	7mer-m8	82	88
hsa_circ_0052112 (5' ... 3') hsa-miR-515-5p (3' ... 5')	GUCUUUCACGAAAGAAAACCUCUU CCUGCACAGAGGGCUUUUAUACAG	8mer-1a	179	186
hsa_circ_0052112 (5' ... 3') hsa-miR-568 (3' ... 5')	CACACAUAUGUAAAUAUGUA NNNNAUUAAUUUCCAAAGACUCA	7mer-m8	161	167
hsa_circ_0052112 (5' ... 3') hsa-miR-924 (3' ... 5')	CGUUCUGUAGUGUUCUGAGA 	8mer-1a	13	20

Figure 1. The flowchart of the research and step-by-step phases of the *in silico* phase of the study.Figure 2. The mean expression of hsa_circ_0052112 in tissue and blood samples of breast cancer patients compared to normal control. Data were normalized by β -actin gene expression. ANOVA test, $p < 0.0001$.

tral components within specific pathways or networks. This analysis aids in elucidating the functional relevance of these hub genes and their involvement in various biological processes. By leveraging Reactome's extensive pathway data, researchers can gain insights into the functional enrichment of hub genes and their potential roles in complex biological systems.

The flowchart of the research and step-by-step phases of the *in silico* phase of the study are shown in figure 1.

Results

hsa_circ_0052112 Expression in Breast Cancer Patients

Tumor tissue and blood plasma analysis: As seen in figure 2, the expression level of hsa_circ_0052112 in the samples of breast cancer patients, both in the tumor tissue and the blood plasma, shows a significant increase compared to the control group ($p < 0.0001$). The mean expression of circular RNA in breast tumors is 7.92 ± 2.85 and in plasma is 3.40 ± 1.68 .

Association between circular RNA expression and receptor status in breast cancer patients

Implications for disease progression: In table 5, the average expression of hsa_circ_0052112 in groups with estrogen and progesterone receptors, as well as HER2, demonstrates a significant increase in both tissue and blood plasma levels compared to the corresponding

Table 5. Mean expression of hsa_circ_0052112 in different pathological conditions and hormone receptor status in breast cancer patients

hsa_circ_0052112 expression	ER-	ER+	PR-	PR+	HER2-	HER2+	TN	Non-TN	LN-	LN+	Stage I&II	Stage III& IV
Tumor	6.49±1.93	10.05±2.68	5.88±1.31	10.17±2.37	6.45±1.72	11.34±1.87	7.78± 1.98	7.93±2.93	5.77±1.32	9.68±2.54	5.86±1.30	9.97±2.48
p-value	0.0002		<0.0001		<0.0001		0.6823		<0.0001		<0.0001	
d	1.522		2.232		2.706		0.062		1.924		1.486	
Plasma	2.67±0.95	4.48±1.93	2.43±0.61	4.47±1.83	2.62±0.82	5.21±1.77	3.19±0.81	3.42±1.75	2.37±0.59	4.23±1.81	2.39±0.57	4.40±1.82
p-value	0.0010		<0.0001		<0.0001		0.7316		<0.0001		<0.0001	
d	1.186		1.494		1.865		0.165		1.371		2.073	

p-value, and Effect size were calculated using T-test and Cohen's d test, respectively.

groups lacking these receptors. This finding suggests that hsa_circ_0052112 may play a role in the biology of hormone receptor-positive breast cancer, potentially influencing tumor behavior and response to hormonal therapies.

Moreover, the increase in hsa_circ_0052112 expression is particularly pronounced in patients with metastatic disease and those at advanced stages of cancer. This correlation raises important questions about the potential role of hsa_circ_0052112 as a biomarker for disease progression. Its elevated levels could indicate a more aggressive tumor phenotype, which may be associated with poorer prognoses and the need for more intensive therapeutic strategies.

Additionally, the relationship between hsa_circ_0052112 expression and lymph node involvement further emphasize its clinical relevance. In patients with lymph node metastasis, hsa_circ_0052112 expression levels were significantly higher than those without lymph node involvement, suggesting its potential utility as a prognostic marker. This association may provide insights into the metastatic potential of breast tumors and aid in the stratification of patients for tailored therapeutic interventions.

Identification of RNA-binding proteins associated with the flanking regions of hsa_circ_0052112

Using the Interactome database, the RBPs that bind to the flanking region of circular RNA hsa_circ_0052112 were identified, which are AGO2, AGO2, EIF4A3, and PTB. Their features and characteristics are listed in table 3.

Identification and characterization of miRNAs targeted by hsa_circ_0052112 with implications for breast cancer

Using the TargetScan database, miRNAs targeted by hsa_circ_0052112 were identified, which consists of 7 microRNAs named hsa-miR-142-5p, hsa-miR-146b-3p, hsa-miR-194, hsa-miR-324-5p, hsa-miR-515-5p, hsa-miR-568, and hsa-miR-924. Their characteristics are listed in table 4. Among miRNAs targeted by circular RNA, enrichment was done using miRcancer database and hsa-miR-194 related to breast cancer, were identified.




Figure 3. Gene Interaction Network and Hub Genes. The gene interaction network was constructed to identify hub genes. Nodes without any connections have been removed for clarity. Hub genes are indicated by squares, and specifically, genes with four or more interactions were selected as hub genes.

Tripartite visualization of circRNA-miRNA-mRNA interactions in breast cancer network

Using TargetScan, the genes targeted by hsa-miR-194 were identified. Among them, genes related to breast cancer were filtered using DisGeNET. In figure 3, the interaction network of these genes is depicted, as can be seen, genes *MAPK1*, *STAT1*, *TRAF6*, and *ERBB4* were identified as hub genes. A hub gene is a gene that plays a central role in a biological network, often exhibiting a high degree of connectivity with other genes or proteins. As shown in figure 4, the circRNA_miRNA_mRNA relationships were visualized using Cytoscape in a tripartite manner.

Enrichment analysis of hub genes and their role in PI3K/AKT signaling pathways

In this study, hub genes (*MAPK1*, *STAT1*, *TRAF6*, and *ERBB4*) enrichment by the Reactome database was done and their significant importance in biological pathways or processes was elucidated and summarized in figure 5. As seen in figure 5, the two important biological processes that statistically showed the highest correlation with these genes are PI3P, PP2A, and IER3 regulate PI3K/AKT signaling R-HSA-6811558 and




Figure 4. The interaction between hsa_circ_0052112_miR-194 and hub genes is visualized using Cytoscape.




Figure 5. Enrichment analysis of hub genes using the reactome database.

negative regulation of PI3K/AKT network R-HSA-199418.

Discussion

circRNAs are stable noncoding RNAs that regulate gene expression in various tissues, including body fluids. This makes them a critical and exciting point of novel research into the pathogenesis of numerous diseases¹⁵. A convergence of data supports that circRNAs are involved in cancer, making them novel biomarkers or potential therapeutic targets¹³.

In a study published in 2018, circRNA microarrays analysis indicated elevated hsa_circ_0052112 expression in invasive breast cancer cell lines, further, the expression of hsa_circ_0052112 in MCF-7 cells and MDA-MB-231 cells was verified by RT-qPCR assay¹³. The expression of this circRNA in breast cancer patients has not been reported.

The present study investigates the relationship between hsa_circ_0052112 expression and clinical parameters in breast cancer, highlighting its potential as a biomarker. Elevated levels of hsa_circ_0052112 were found in tumor tissues and plasma of breast cancer patients, particularly in advanced stages and those with lymph node involvement, suggesting its role in disease progression and tumor aggressiveness. The correlation between hsa_circ_0052112 expression and hormone receptor status indicate its utility not only as a marker for breast cancer presence, but also in understanding tumor behavior, which may inform treatment responses.

The non-invasive measurement of hsa_circ_0052112 in plasma offers a promising approach for monitoring disease progression and treatment efficacy, potentially allowing for its incorporation into routine clinical assessments. Should future studies confirm its prognostic value, hsa_circ_0052112 could aid in patient stratification, guiding personalized treatment strategies based on expression levels.

The traditional markers ER, PR, and HER2 have been extensively validated in clinical settings and play crucial roles in determining the therapeutic approach for breast cancer patients. However, these biomarkers often exhibit variability in expression and can be influenced by tumor heterogeneity, potentially limiting their reliability as sole indicators of disease status¹⁶. In contrast, hsa_circ_0052112 demonstrates a unique profile characterized by its circular RNA structure, which confers increased stability in circulation compared to linear RNA species. This stability is particularly advantageous for its potential application in liquid biopsies, as it minimizes degradation and allows for more accurate detection in blood samples. In the realm of emerging biomarkers, such as Circulating tumor DNA (ctDNA) and microRNAs, hsa_circ_0052112 could complement these modalities by offering insights into tumor dynamics. While ctDNA found in the bloodstream and refers to DNA from cancerous cells and tumors provides information about the genetic alterations within the tumor¹⁷, hsa_circ_0052112 may reflect the tumor's functional status and cellular processes more accurately. Furthermore, microRNAs are often less stable and can be influenced by various physiological states¹⁸, whereas the stability of hsa_circ_0052112 could lead to more consistent and reliable measurements.

The *in silico* phase of our study indicated that the RNA-binding protein sites matching flanking regions of hsa_circ_0052112 are AGO2, EIF4A3, and PTB. Understanding the RNA-binding protein sites that correspond to the flanking regions of circRNAs is essential for comprehending their biogenesis, function, and role in disease, as well as for advancing research methodologies in molecular biology. Identifying RBP target sites is an important step toward understanding the mechanisms by which they conduct post-transcriptional regulation¹⁹. AGO2 (Argonaute 2) is a vital pro-

tein taking part within the RNA-Induced Silencing Complex (RISC) and it plays a pivotal role in the regulation of gene expression through its function as an RNA-Binding Protein (RBP). It is well-known for its involvement in the mechanisms of miRNA and small interfering RNA (siRNA) pathways, facilitating the silencing of target mRNAs²⁰. AGO2 serves as a critical RBP for hsa_circ_0052112, influencing its function and stability and potentially playing a role in the regulatory networks involving miRNAs and gene expression.

EIF4A3 (Eukaryotic Translation Initiation Factor 4A3) is a member of the eIF4A family of RBPs and plays a critical role in the regulation of mRNA translation and stability. It was recently recognized as an oncogene. It is ordinarily recognized for its involvement within the Exon Junction Complex (EJC), that is vital for mRNA processing, transport, and translation²¹.

Polypyrimidine Tract-Binding Protein (PTB), also known as PTBP1, is an RNA-binding protein that is involved in various processes, including splicing, mRNA stability, and translation. Analysis indicated that overexpression of PTBP1 generally predicted poor overall survival in patients with tumors such as liver hepatocellular carcinoma, adrenocortical carcinoma, lung adenocarcinoma, and skin cutaneous melanoma²².

circRNAs are a group of endogenous RNAs that control gene expression at the transcriptional and post-transcriptional levels. Recent reports have indicated that circRNAs serve as innovative diagnostic biomarkers and favorable therapeutic targets for several cancer types by interacting with other non-coding RNAs such as miRNAs²³. The miRNAs are revealed as pivotal risk factors and regulatory elements in cancer by regulating the expression of their target genes. Genetic interactions between miRNAs and circular RNAs can generate complex regulatory networks with various carcinogenic processes that play significant roles in tumorigenesis, cancer development, and cancer progression²³. CircRNAs have recently been identified for their miRNA sponge function, which enables them to modulate the expression of miRNA target genes by acting as competitive endogenous RNAs (ce-circRNAs)²⁴.

In the *in silico* phase of our study, among the miRNAs that were introduced as targets of hsa_circ_0052112 by target scan, miR-194 was found to be associated with all subtypes of breast cancer even triple negatives^{25,26}. Studies have shown that miR-194 can act as a tumor suppressor in breast cancer by regulating the expression of target genes involved in cell cycle control and apoptosis²⁷.

On the other hand, miR-194 altered expression has been associated with poor prognosis in breast cancer patients. In some cases, downregulation of miR-194 has been linked to increased metastatic potential and resistance to therapy, implying that it may act as a potential biomarker for breast cancer progression and

treatment response²⁸. miR-194 dual role as a potential tumor suppressor and its association with clinical outcomes highlights its importance in breast cancer research. In this research article, miR-194 is highlighted as a key microRNA involved in the post-transcriptional regulation of gene expression, playing critical roles in various biological processes such as cell proliferation, differentiation, apoptosis, and metabolism. Notably, as discussed earlier, miR-194 has been identified as a potential tumor suppressor in several cancers, including breast cancer. It can inhibit oncogene expression and promote apoptosis, thus influencing the balance between cell survival and death. Additionally, miR-194 regulates genes involved in cell adhesion, migration, and invasion, thereby affecting cancer metastasis²⁹.

In silico part of our study revealed that *MAPK1*, *STAT1*, *TRAF6*, and *ERRB4* were identified as hub genes targeted by miR-194. Hub genes are central genes within a biological network that play a crucial role in regulating various cellular processes. They are often characterized by a high degree of connectivity, meaning that they interact with many other genes or proteins. In the context of gene networks, hub genes can influence the behavior of the network as a whole and are often implicated in critical biological functions, disease mechanisms, and responses to environmental changes. Identifying hub genes can be important for understanding complex biological systems and developing targeted medicine therapies.

The enrichment analysis identified significant associations between the hub genes and two critical biological processes: "PI3P, PP2A, and IER3 regulate PI3K/AKT signaling" and "negative regulation of the PI3K/AKT network". These findings underscore the potential roles of the hub genes in modulating the PI3K/AKT signaling pathway. For instance, *MAPK1* is linked to cell proliferation and survival, suggesting its involvement in tumor growth and therapy resistance³⁰. *STAT1*, as a transcription factor, may influence the expression of genes that regulate the PI3K/AKT pathway, impacting cellular responses to disease signals³¹. *TRAF6* is known to activate AKT, promoting cell survival in cancer contexts³², while *ERRB4* is associated with signaling cascades that intersect with this pathway, particularly in breast cancer progression³³. The miR-194 interacts with the PI3K/AKT signaling pathway, a crucial cascade that regulates cell growth, survival, and metabolism. Activation of this pathway promotes cell survival and proliferation, often exploited by cancer cells, and is associated with chemotherapy resistance³⁴. The interplay between miR-194 and the PI3K/AKT pathway is significant in breast cancer, as low levels of miR-194 and aberrant activation of the PI3K/AKT pathway can serve as prognostic biomarkers linked to poorer outcomes and aggressive tumor characteristics.

Overall, the findings underscore the importance of miR-194 and the PI3K/AKT pathway in breast cancer

biology, offering insights that could inform prognostic assessments and therapeutic strategies aimed at targeted interventions in breast cancer management. Additionally, hsa_circ_0052112 may modulate the PI3K/AKT pathway by acting as a sponge for microRNAs, thereby upregulating target hub genes, and influencing RNA-binding proteins and the transcription of nearby genes. The interactions between hsa_circ_0052112 and the hub genes could significantly affect breast cancer progression, as enhanced PI3K/AKT activation is commonly linked to increased cell proliferation, survival, and metastasis. These insights may inform future therapeutic strategies targeting the dysregulated PI3K/AKT pathway in breast cancer.

The phosphatidylinositol 3-kinase (PI3K)/Akt pathway plays a pivotal role in various cellular processes and is aberrantly engaged in cancers, contributing to the manifestation and progression of tumors. PI3K/AKT signaling pathway is a critical intracellular signaling pathway that regulates multiple cellular functions, including growth, proliferation, survival, and metabolism. This pathway is often activated in response to growth factors and hormones, playing a significant role in normal cellular processes³⁵.

The mechanism of action of PI3K enzymes PI3K/AKT signaling pathway is as follows: phosphorylate phosphatidylinositol to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3), which activates AKT, leading to its translocation to the plasma membrane and subsequent phosphorylation. This activation promotes cell survival, growth, and metabolism, thereby facilitating cell proliferation and inhibiting apoptosis, which can contribute to tumor growth, particularly in breast cancer³⁶. In hormone receptor-positive breast cancers, estrogen signaling can aberrantly activate this pathway, enhancing tumor growth and metastasis³⁷. As mentioned earlier The PI3K/AKT pathway is one of the most frequently over-activated intracellular pathways in several human cancers. This pathway, influencing various downstream target proteins, contributes to the carcinogenesis, proliferation, invasion, and metastasis of tumor cells. A multi-level impairment, including mutation, aberrant regulation of miRNAs, and aberrant phosphorylation of cascade factors, has been revealed in multiple cancer types. The deregulation of this pathway counteracts common therapeutic strategies and contributes to multidrug resistance. Interpretation of this pathway is essential for developing effective treatment strategies and personalized medicine approaches in breast cancer management³⁸.

The PI3K/AKT signaling pathway is crucial in breast cancer progression, affecting growth, survival, metabolism, and motility. It is activated by growth factors and hormones, leading to AKT activation, which promotes cell cycle progression and tumor growth while inhibiting apoptosis. This pathway supports the energy demands of rapidly proliferating tumor cells and enhances the tumor microenvironment

through angiogenesis and metastasis. In hormone receptor-positive breast cancers, estrogen further stimulates this pathway, contributing to resistance against therapies like tamoxifen. Targeting the PI3K/AKT pathway is a focus of therapeutic interventions, although resistance is common, highlighting the need for combination therapies. Mutations in related genes, such as PIK3CA, are prevalent in breast cancer, underscoring the pathway's significance³⁹.

Conclusion

The study highlights the significant upregulation of hsa_circ_0052112 in breast cancer patients, particularly in advanced stages and in association with specific receptor statuses, suggesting its potential role as a biomarker and therapeutic target. Additionally, the involvement of miR-194 and the PI3K/AKT signaling pathway underscores the complexity of gene regulation in breast cancer, emphasizing the need for further research to explore their implications in disease progression and treatment strategies.

While our study provides valuable insights into the role of hsa_circ_0052112, it is important to acknowledge certain limitations, including the relatively small sample size, which may affect the generalizability of our findings. Additionally, the absence of external validation in larger, more diverse cohorts restricts our ability to draw definitive conclusions regarding its clinical relevance. Future research should focus on expanding the sample size and conducting multicenter studies to validate our results in broader populations. Furthermore, exploring the therapeutic implications of hsa_circ_0052112 could lead to novel interventions, necessitating investigations into its functional mechanisms and potential applications in targeted therapies. This approach may enhance our understanding of its role in disease pathology and pave the way for innovative treatment strategies.

Acknowledgement

We sincerely thank all the individuals who participated in this study. We would like to acknowledge Prof. Mozdarani and Prof. Kaviani for their support. This work was supported by a grant from NIGEB (Grant No. 805). The ethical code number is IR.MO-DARES.REC.1398.148.

Conflict of Interest

No potential conflicts of interest were disclosed by the authors.

References

1. Feinberg AP, Vogelstein B. Hypomethylation distinguishes genes of some human cancers from their normal counterparts. *Nature* 1983;301(5895):89-92.
2. Baylin SB, Jones PA. A decade of exploring the cancer epigenome-biological and translational implications. *Nat Rev Cancer* 2011;11(10):726-34.

3. Schröder R, Illert AL, Erbes T, Flotho C, Lübbert M, Duque-Afonso J. The epigenetics of breast cancer - Opportunities for diagnostics, risk stratification and therapy. *Epigenetics* 2022;17(6):612-24.
4. Thakur C, Qiu Y, Fu Y, Bi Z, Zhang W, Ji H, et al. Epigenetics and environment in breast cancer: New paradigms for anti-cancer therapies. *Front Oncol* 2022;12: 971288.
5. Nasirpour MH, Salimi M, Majidi F, Minucheir Z, Mozdarani H. Study of DACH1 Expression and its Epigenetic Regulators as Possible Breast Cancer-Related Biomarkers. *Avicenna J Med Biotechnol* 2023;15(2):108-17.
6. Feng Y, Spezia M, Huang S, Yuan C, Zeng Z, Zhang L, et al. Breast cancer development and progression: Risk factors, cancer stem cells, signaling pathways, genomics, and molecular pathogenesis. *Genes Dis* 2018;5(2):77-106.
7. Wang Y, Zhang J, Yang Y, Liu Z, Sun S, Li R, et al. Circular RNAs in human diseases. *MedComm* (2020) 2024;5(9):e699.
8. Wan B, Liu B, Lv C. Progress of research into circular RNAs in urinary neoplasms. *PeerJ* 2020; 8:e8666.
9. Dragomir MP, Calin GA. CpG island hypermethylation go circular (RNA). *Oncotarget* 2018;9(69):33052-3.
10. Larkin L. Breast cancer genetics and risk assessment: an overview for the clinician. *Climacteric* 2023 Jun;26(3): 229-34.
11. Shayan S, Arashkia A, Bahramali G, Azadmanesh K. Investigating the Effects of HMGB1 Overexpression on Colorectal Cancer Cell Migration via Oncolytic Herpes simplex Virus Type 1 (oHSV-1). *Avicenna J Med Biotechnol* 2024;16(2):120-9.
12. Erdogan C, Suer I, Kaya M, Ozturk S, Aydin N, Kurt Z. Bioinformatics analysis of the potentially functional circRNA-miRNA-mRNA network in breast cancer. *PLoS One* 2024;19(4):e0301995.
13. Zhang HD, Jiang LH, Hou JC, Zhong SL, Zhou SY, Zhu LP, et al. Circular RNA hsa_circ_0052112 promotes cell migration and invasion by acting as sponge for miR-125a-5p in breast cancer. *Biomed Pharmacother* 2018; 107:1342-53.
14. Conn VM, Chinnaiyan AM, Conn SJ. Circular RNA in cancer. *Nat Rev Cancer* 2024;24(9):597-613.
15. Cieśla M, Darmochwal-Kolarz DA, Kwaśniak K, Pałka A, Kolarz B. Plasma Circular-RNA 0005567 as a Potential Marker of Disease Activity in Rheumatoid Arthritis. *Int J Mol Sci* 2023;25(1):417.
16. Vellan CJ, Islam T, De Silva S, Mohd Taib NA, Prasanna G, Jayapalan JJ. Exploring novel protein-based biomarkers for advancing breast cancer diagnosis: A review. *Clin Biochem* 2024;129:110776.
17. Xu J, Gao H, Guan X, Meng J, Ding S, Long Q, et al. Circulating tumor DNA: from discovery to clinical application in breast cancer. *Front Immunol* 2024;15: 1355887.
18. Zhu M, Gao Y, Zhu K, Yuan Y, Bai H, Meng L. Exosomal miRNA as biomarker in cancer diagnosis and prognosis: A review. *Medicine (Baltimore)* 2024;103(42): e40082.
19. Li X, Kazan H, Lipshitz HD, Morris QD. Finding the target sites of RNA-binding proteins. *Wiley Interdiscip Rev RNA* 2014;5(1):111-30.
20. Nakanishi K. Anatomy of four human Argonaute proteins. *Nucleic Acids Res* 2022; 50(12):6618-38.
21. Hu B, Chen R, Jiang M, Xiong S, Liu X, Fu B. EIF4A3 serves as a prognostic and immunosuppressive microenvironment factor and inhibits cell apoptosis in bladder cancer. *PeerJ*. 2023;11:e15309.
22. Huang Q, Gu S, Fang J, Li X, Lin L. A pan-cancer analysis of the oncogenic role of polypyrimidine tract binding protein 1 (PTBP1) in human tumors. *Medicine (Baltimore)* 2022;101(52):e32428.
23. Kim WR, Park EG, Lee DH, Lee YJ, Bae WH, Kim HS. The Tumorigenic Role of Circular RNA-MicroRNA Axis in Cancer. *Int J Mol Sci* 2023; 24(3):3050.
24. Fraboulet RM, Si Ahmed Y, Aubry M, Corre S, Galibert MD, Blum Y. Cirscan: a shiny application to identify differentially active sponge mechanisms and visualize circRNA-miRNA-mRNA networks. *BMC Bioinformatics* 2024;25(1):53.
25. Yen YT, Yang JC, Chang JB, Tsai SC. Down-Regulation of miR-194-5p for Predicting Metastasis in Breast Cancer Cells. *Int J Mol Sci* 2021;23(1):325.
26. Caetano S, Garcia AR, Figueira I, Brito MA. MEF2C and miR-194-5p: New Players in Triple Negative Breast Cancer Tumorigenesis. *Int J Mol Sci* 2023;24(18):14297.
27. Yang F, Xiao Z, Zhang S. Knockdown of miR-194-5p inhibits cell proliferation, migration and invasion in breast cancer by regulating the Wnt/ β -catenin signaling pathway. *Int J Mol Med* 2018; 42(6):3355-63.
28. Sereno M, Haskó J, Molnár K, Medina SJ, Reisz Z, Malhó R, et al. Downregulation of circulating miR 802-5p and miR 194-5p and upregulation of brain MEF2C along breast cancer brain metastasization. *Mol Oncol* 2020;14 (3):520-38.
29. Yu M, Du H, Zhang C, Shi Y. miR-192 family in breast cancer: Regulatory mechanisms and diagnostic value. *Biomed Pharmacother* 2024;175:116620.
30. Braicu C, Buse M, Busuioc C, Drula R, Gulei D, Raduly L, et al. A Comprehensive Review on MAPK: A Promising Therapeutic Target in Cancer. *Cancers (Basel)* 2019; 11(10):1618.
31. Yang B, Singh S, Bressani R, Kanmogne GD. Cross-talk between STAT1 and PI3K/AKT signaling in HIV-1-induced blood-brain barrier dysfunction: role of CCR5 and implications for viral neuropathogenesis. *J Neurosci Res* 2010 Nov 1;88(14):3090-101.
32. Feng L, Feng S, Nie Z, Deng Y, Xuan Y, Chen X, et al. TRAF6 Promoted Tumor Glycolysis in Non-Small-Cell Lung Cancer by Activating the Akt-HIF α Pathway. *Bio-med Res Int* 2021;2021:3431245.
33. Toomey S, Eustace AJ, Fay J, Sheehan KM, Carr A, Milewska M, et al. Impact of somatic PI3K pathway and ERBB family mutations on pathological complete re-

- sponse (pCR) in HER2-positive breast cancer patients who received neoadjuvant HER2-targeted therapies. *Breast Cancer Res* 2017;19(1):87.
34. Wang C, Fu R, Wang Y, Wei J, Yu Y, Hu L, Zhang C. miR-124-3p and miR-194-5p regulation of the PI3K/AKT pathway via ROR2 in medulloblastoma progression. *Cancer Gene Ther* 2024;31(6):941-54.
 35. He Y, Sun MM, Zhang GG, Yang J, Chen KS, Xu WW, et al. Targeting PI3K/Akt signal transduction for cancer therapy. *Signal Transduct Target Ther* 2021;6(1):425.
 36. Glaviano A, Foo ASC, Lam HY, Yap KCH, Jacot W, Jones RH, et al. PI3K/AKT/mTOR signaling transduction pathway and targeted therapies in cancer. *Mol Cancer* 2023;22(1):138.
 37. Khatpe AS, Adebayo AK, Herodotou CA, Kumar B, Nakshatri H. Nexus between PI3K/AKT and Estrogen Receptor Signaling in Breast Cancer. *Cancers (Basel)* 2021;13(3):369.
 38. Rascio F, Spadaccino F, Rocchetti MT, Castellano G, Stallone G, Netti GS, et al. The Pathogenic Role of PI3K/AKT Pathway in Cancer Onset and Drug Resistance: An Updated Review. *Cancers (Basel)* 2021;13(16):3949.
 39. Browne IM, André F, Chandarlapaty S, Carey LA, Turner NC. Optimal targeting of PI3K-AKT and mTOR in advanced oestrogen receptor-positive breast cancer. *Lancet Oncol* 2024 Apr;25(4):e139-e151. Erratum in: *Lancet Oncol* 2024;25(6):e234.