



## The Effect of Sodium Selenite on Expression of Mitochondrial Transcription Factor A during *In Vitro* Maturation of Mouse Oocyte

Tina Moshaashae<sup>1</sup>, Saeed Zavareh<sup>2</sup>, Shahram Pourbeiranvand<sup>1</sup>, and Mojdeh Salehnia<sup>1\*</sup>

1. Department of Anatomy, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Faculty of Biology, Damghan University, Damghan, Iran

### Abstract

**Background:** The aim of the present study was to investigate the effect of Sodium Selenite (SS) supplemented media on oocyte maturation, expression of mitochondrial transcription factor A (TFAM) and embryo quality.

**Methods:** Mouse Germinal Vesicle (GV) oocytes were collected after administration of Pregnant Mare Serum Gonadotropin (PMSG); in experimental group 1, oocytes were cultured and then subjected for *in vitro* maturation in the absence of SS, and in experimental group 2, they were matured *in vitro* in the presence of 10 ng/ml of SS up to 16 hr. The control group included MII oocytes obtained from the fallopian tubes after ovarian stimulation with PMSG, followed by human chorionic gonadotropin. Then, the expression of TFAM in MII oocytes in all three groups was investigated using real-time RT-PCR. The fertilization and embryo developmental rates were assessed, and finally the quality of the blastocysts was evaluated using propidium iodide staining.

**Results:** The oocyte maturation rate to MII stage in SS treated group was significantly higher than non-treated oocytes (75.65 vs. 68.17%,  $p < 0.05$ ). Also, the rates of fertilization, embryo development to blastocyst stage as well as the cell number of blastocyst in SS supplemented group were higher than other experimental group ( $p < 0.05$ ). There was a significant decrease in TFAM gene expression in both *in vitro* groups compared to the group with *in vivo* obtained oocytes ( $p < 0.05$ ). Moreover, there was a significant increase in TFAM gene expression in oocytes that matured in the presence of SS compared to that of the group without SS ( $p < 0.05$ ).

**Conclusion:** Supplementation of oocyte maturation culture media with SS improved the development rate of oocytes and embryo and also enhanced TFAM expression in MII oocytes which can affect the mitochondrial biogenesis of oocytes.

*Avicenna J Med Biotech 2021; 13(2): 81-86*

**Keywords:** Gene expression, *In vitro* oocyte maturation, Mice, Mitochondrial transcription factor A, Oocytes, Sodium selenite

### Introduction

*In Vitro* Maturation (IVM) of oocyte refers to development of immature germinal oocytes in the medium supplemented with some growth factors and hormones<sup>1</sup>. The numerous successful pregnancies following IVM have caused this therapeutic method to be considered low risk and effective in assisted reproduction<sup>1-5</sup>. There are several groups of candidates for this method such as patients with polycystic ovarian syndrome, low follicles reserves, repeated IVF failure, ovarian hypertension, resistant ovary syndrome and young women who suffer from different types of cancer<sup>1-5</sup>. Although the rate of pregnancy after IVM of oocytes increases in these patients, it is still considered as an experimental method<sup>5</sup>. It was reported that vari-

ous factors such as the culture media, supplementation of hormones, co-culture with somatic cells and adding some antioxidants could affect the outcome of oocyte IVM<sup>6-9</sup>.

For oocyte maturation, the mitochondria are required as the source of energy and for intracellular calcium storage<sup>10,11</sup>. It also regulates calcium fluctuations during fertilization which is very important for the fertilization and embryonic development<sup>11</sup>. The transcription of mitochondrial DNA is essential for its biogenesis during the oocyte development and mitochondrial DNA (mtDNA) content is correlated with oocyte fertilization potential<sup>12-15</sup> and immature oocytes have fewer copies of mtDNA<sup>13</sup>. Mitochondrial transcription

factor A (*TFAM*) is an important mitochondrial DNA transcription factor and is coded by nuclear DNA<sup>16</sup>. It creates structural stability and converts the genome to nucleotide-like structures by shortening the length of the mitochondrial genome through covering it and gives a regular curve to it<sup>16,17</sup>. Decline in the *TFAM* expression is associated with the decrease in mtDNA content; in other words, the mtDNA content is directly related to the expression of *TFAM*<sup>18</sup>. Moreover, the high level of Reactive Oxygen Species (ROS) causes damage to mtDNA<sup>19-22</sup>.

It has also been shown that Sodium Selenite (SS) as a non-enzymatic antioxidant improves development of follicles, oocyte and embryo by elevating the total antioxidant capacity and reducing the level of ROS<sup>13,23-26</sup>. In our previous studies, it was shown that adding SS to the oocyte maturation media increased mtDNA content of the oocytes<sup>13,26</sup>. This study was designed to bring some additional information regarding the effect of SS on the biogenesis of oocyte mitochondria through *TFAM* expression. Thus, the aim of the present study was to investigate the effect of SS supplementation in the culture media of immature oocyte on *TFAM* expression as well as oocyte and embryo development.

### Materials and Methods

All laboratory materials, except those mentioned, were purchased from Sigma Aldrich, England.

#### Animal model

In this study, 60 female NMRI mice, 6-8 weeks old and 10 male NMRI mice, 8-12 weeks old were used. The mice were kept under standard conditions of 12 hr light: 12 hr dark, 20-24°C, and 40-50% humidity in the animal house of Tarbiat Modares University. The mice had adequate access to food and water.

#### Study design

For investigating the effect of sodium selenite on *TFAM* expression of the oocyte, the cumulus cells were removed from oocytes in all study groups. The three groups in this study were (a) *in vivo* control group in which the metaphase II (MII) oocytes were collected from fallopian tubes, (b) experimental group 1 in which *in vitro* maturation of Germinal Vesicle (GV) oocytes was done in the absence of SS, and (c) experimental group 2 in which *in vitro* maturation of GV oocytes was done in the presence of SS.

In both *in vitro* matured groups, the oocyte maturation and embryonic developmental rates and blastocyst cell count were studied in comparison with *in vivo* obtained blastocysts. Also, *TFAM* expression in *in vitro* obtained MII oocytes was analyzed and compared with *in vivo* obtained MII oocytes. All experiments were approved by the ethics committee of Tarbiat Modares University, Tehran, Iran (IR.MODARES.REC.1398.006).

#### GV oocytes collection

In order to collect the GV oocytes, 10 IU of Preg-

nant Mare Serum Gonadotropin (PMSG) were injected to each mouse intraperitoneally and 48 hr later, the mentioned mice were sacrificed through cervical dislocation. Initially, by cutting the skin and then through a longitudinal incision in the abdominal region, two ovaries were extracted and placed within the culture media. Then, the oocytes at GV stage were isolated using the needle attached to the insulin syringe under a stereo microscope. The GV oocytes with a centrally located nucleus, clear cytoplasm, uniform zona pellucida and homogeneous perivitelline space were chosen<sup>13</sup>. The cumulus cells were detached using repeated pipetting and enzymatic methods. All GV oocytes were denuded and were considered for *in vitro* maturation in two experimental groups.

#### MII oocytes collection

To collect the MII oocytes, 10 IU of PMSG were injected intraperitoneally to adult female mice and 48 hr later, 10 IU of human Chorionic Gonadotropin (hCG) were also injected. Thereafter, 16 hr after the last injection, the mice were sacrificed through cervical dislocation and MII oocytes were collected from their fallopian tube, denuded, and then kept at -80°C for molecular study.

#### *In vitro* maturation

The collected GV oocytes were cultured in  $\alpha$ -MEM medium supplemented with 50 of  $\mu\text{g/ml}$  penicillin, 75 of  $\mu\text{g/ml}$  streptomycin, 0.23 mM of sodium pyruvate, 10% of fetal bovine serum, 75 mIU/ml of recombinant follicle stimulating hormone, 10 IU/ml of hCG and in the presence and absence of 10 ng/ml of SS according to the experimental design under mineral oil at 37°C, 100% humidity in 5% CO<sub>2</sub> for 16 hr<sup>13</sup>. The maturation of oocytes was assessed under inverted microscope. The oocytes with clear nucleus were categorized as GV oocytes; those with uniform cytoplasm and without nucleus were classified as Germinal Vesicle Breakdown (GVBD) or metaphase I, and the oocytes with first polar body were considered as the MII oocytes. This experiment was replicated at least seven times. MII oocytes were collected and considered for the subsequent studies.

#### *In vitro* fertilization

The sperm was prepared from adult male mice at the age of 8-12 weeks. The mice were sacrificed through cervical dislocation and their epididymis tail was separated and transferred to 300  $\mu\text{l}$  drops of culture medium under mineral oil. Using a needle attached to insulin syringe, several small cuts were made in the epididymis tail so that the sperm was expelled out of the epididymis and kept for 1.5 hr in incubator at 37°C, 5% CO<sub>2</sub> and 100% humidity until the completion of capacitation. The oocytes were transferred to drops containing a concentration of 10<sup>5</sup> sperm per ml. They were then kept under the following conditions for 4-6 hr and then, the oocytes were washed several times with fresh media and culture for 120 hr. During the culture peri-

od, the embryonic developmental stages including the two-cell, four-cell, morula and blastocyst were assessed using inverted microscope. The embryos at blastocyst stage were collected for staining with Propidium Iodide (PI).

#### Harvesting of *in vivo* blastocyst

In order to collect *in vivo* obtained blastocysts, female mice were injected with 10 IU of PMSG hormone followed by 10 IU of hCG after 48 hr. Then, they were placed alongside a male mouse. The morning of the next day, the pregnant mice were separated and 96 hr after hCG injection, the pregnant female mice were sacrificed through cervical dislocation and the embryos were collected by flushing uterine horns.

#### Blastocysts staining

For embryo staining, the collected blastocysts from both *in vitro* groups (n=3 in each group in 3 repeats) were put in 100 µg/ml of PI and 1% of Triton ×100 for 10 s. Then, the stained embryos were placed in glycerol and were investigated under fluorescence microscope<sup>27</sup>.

#### Molecular analysis

In order to investigate the expression of *TFAM* gene in MII oocytes, real-time RT-PCR was used. Total RNA was extracted from the MII oocytes (n=10 in each group) using the TRIzol<sup>®</sup> reagent (Invitrogen, USA) according to the manufacturer's guidance. The quality of RNA was evaluated using the spectrophotometry (Eppendorf, Hamburg, Germany). The ratio of optical density, A260/A280 ≥ 1.8, was accepted for RNA and used for reverse transcription. The cDNA synthesis was performed based on the cDNA synthesis kit (Takara, Japan) manual. The sequence of forward and reverse primers (Table 1) was designed using Allele ID software, version 7.5 (PREMIER Biosoft, USA). To ensure specificity for the mRNA of target genes and nonreplication of genomic DNA, the sequence of primers was blasted using Primer 3 software at NCBI website. In addition, no genomic DNA replication by each of the designed primers was confirmed

through a real-time RT-PCR reaction with 25 ng of total RNA solution as template and electrophoresis of PCR product on 1.5% agarose gel. In this reaction, 25 ng of cDNA synthesized for each gene was used as positive control in a separate microtube in qPCR reaction. The real-time RT-PCR reaction was performed by StepOnePlus™ Real-Time PCR System (Applied Biosystems, USA) and using SYBR Green qPCR Master Mix (Ampliqon, Denmark). The mixture of the reaction contained SYBER Green (5 µl), forward primer (0.5 µl), reverse primer (0.5 µl), and cDNA (4 µl) in 10 µl of final volume. PCR reaction was adjusted for 15 min at 95°C. It was followed by 40 cycles of denaturation at 95°C for 10 s and annealing/extension at 60°C for 45 s. The level of expression of each gene was normalized through the β-actin as internal control gene and then relative expression of genes was calculated using the  $-2^{-\Delta\Delta C_t}$  formula.

#### Statistical analysis

The statistical analysis was performed using SPSS<sup>®</sup> version 24 (SPSS Inc., Chicago, USA). After investigating the normal distribution of data using Shapiro-Wilk test, in order to compare the percentage of GV, GVBD, and MII oocytes as well as viability between two experimental groups, unpaired t-test was used. The real-time RT-PCR data were normalized using Livak method and one-way analysis of variance (ANOVA) and post hoc Tukey tests were used. The results have been presented based on Mean±SD and p≤0.05 was considered statistically significant.

## Results

#### Developmental and fertilization rates

As shown in table 2, after 16 hr of *in vitro* culture of GV oocytes, the mean total percentage of survived oocytes in the group treated with SS and in the group matured without SS supplementation was 90.97±4.69 and 87.82±5.75, respectively and there was no significant difference between the two studied groups in this regard. Further, the rate of MII oocytes in the SS treat-

Table 1. List of primer

Gene	Primer pair sequence(5'-3')	Accession number	PCR product size (bp)
<i>B-actin</i>	F: TGTGACGTTGACATCCGTAA R: GCTAGGAGCCAGAGCAGTAA	NM-007393	64
<i>Tfam</i>	F: AAGGGAATGGGAAAGGTAGA R: AACAGGACATGGAAAGCAGAT	NM-011045	76

Table 2. Developmental rates of *in vitro* cultured GV oocytes in the Sodium Selenite (SS) treated and non treated groups

Groups	Total number of GV oocytes	No. of survived (Mean%±SD)	No. of degenerated (Mean%±SD)	No. of GV (Mean%±SD)	No. of GVBD (Mean%±SD)	No. of MII (Mean%±SD)
Without SS	202	176 (87.82±5.75)	26 (12.18±5.75)	30 (15.10±5.40)	8 (7.96±2.90)	138 (67.17±2.47)
With SS	222	201 (90.97±4.69)	21 (9.03±4.69)	24 (15.33±6.93)	9 (8.42±6.62)	168 (75.65±6.56) *

\* Significant differences compared with other group in the same column (p<0.05). Seven experimental replicates were performed for each group. GV= Germinal Vesicle, GVBD=Germinal Vesicle Breakdown.

Table 3. Fertilization and embryonic developmental rates of oocytes treated with Sodium Selenite (SS) treated and non treated groups

Groups	Total no of Inseminated oocyte	No. of fertilized (Mean%±SD)	No. of 2 cell (Mean%±SD)	No. of 4 cell (Mean%±SD)	No. of morulla (Mean%±SD)	No. of blastocyst (Mean%±SD)
With SS	43	35 (81.39%±3.51)	25 (71.40%±3.51)	17 (48.50%±2.08)	8 (22.80%±1.52)	6 (17.40%±10)
Without SS	40	28 (70.00%±1.52) *	19 (67.80%±2.3) *	13 (46.40%±1.52) *	5 (17.80%±1.52) *	3 (10.71%±10) *

\* Significant differences compared with other group in the same column ( $p < 0.05$ ).

ed group ( $75.65 \pm 6.56$ ) was significantly higher than the group matured without SS ( $68.17 \pm 2.47$ ;  $p < 0.05$ ). Statistical analysis indicated that the rates of GV and GVBD oocytes did not have any significant difference between the two tested groups.

The mean percentage of fertilization, two-cell, four-cell, and morula embryos in all studied groups are summarized in table 3. The mean percentage of oocytes that fertilized and reached to blastocyst stages in the presence of SS was  $81.39 \pm 3.51$  and  $17.14 \pm 10$ , respectively, while those of group without SS treatment were  $70 \pm 1.52$  and  $10.71 \pm 10$ , respectively. According to statistical analysis, the rates of fertilization and embryo development in the groups undergoing SS treatment increased significantly compared to the group without it ( $p < 0.05$ ).

#### Blastocyst cell count

The representative figures of blastocysts obtained from *in vitro* matured oocytes in the absence and presence of SS and *in vivo* obtained blastocysts as controls that were stained with PI are demonstrated in figure 1 A-C. As these figures show, the cell number of blastocyst was significantly higher in SS treated group in comparison with the group matured without SS ( $61 \pm 0.81$  vs.  $38 \pm 0.81$ ;  $p < 0.05$ ). The *in vivo* obtained blastocysts significantly showed higher cell number




Figure 1. Staining of blastocysts derived from *in vivo* condition (A) and *in vitro* matured germinal vesicle oocytes in the absence (SS-; B) and presence of Sodium Selenite (SS+; C). The comparison of total cell number of blastocysts in previous groups was shown in part D. Letter a indicated the differences of each group with the control group and letter b showed significant differences between Sodium Selenite treated group with non-treated group ( $p < 0.05$ ).




Figure 2. The relative expression of *TFAM* and  $\beta$ -actin in MII oocytes obtained from *in vivo* and *in vitro* conditions in the presence and absence of Sodium Selenite (A). a: significant differences with *in vivo* group; b: significant differences with non-treated *in vitro* group ( $p < 0.05$ ). The representative figure of gel electrophoresis of PCR product in studied groups (B). No template control (NTC).

( $80.25 \pm 0.83$ ) in comparison with both *in vitro* matured blastocysts ( $p < 0.05$ ; Figure 1D).

#### Real-time RT-PCR

The relative expression of *TFAM* to housekeeping gene ( $\beta$ -actin) in the oocytes obtained *in vivo* and *in vitro* in the presence and absence of SS was  $1.02 \pm 0.03$ ,  $0.7 \pm 0.04$ , and  $0.25 \pm 0.07$ , respectively. The expression levels of *TFAM* decreased significantly in both IVM culture groups in comparison with *in vivo* obtained oocytes ( $p < 0.05$ ; Figure 2). Comparing the two *in vitro* groups, relative expression of *TFAM* was significantly higher in the oocytes undergoing SS treatment compared to the group without it ( $p < 0.05$ ; Figure 2). The representative figure of PCR product in the studied groups is demonstrated in figure 2B.

#### Discussion

The present study investigated the effect of SS on the maturation of GV oocytes as well as the expression of *TFAM* in *in vitro* cultured oocytes in comparison with MII oocytes obtained from fallopian tubes.

The results of the present study are in line with previous studies that showed SS can be considered as an effective factor in the oocyte maturation and the rates of developmental parameters of *in vitro* cultured oocytes treated with SS increased significantly compared to the oocytes cultured in the medium without SS<sup>13,25</sup>. The similar effects of SS supplement in the culture medium on the development of follicle and oocyte in some mammal species have also been previously reported<sup>23,26,28-31</sup>. The obtained results may be related to the antioxidant properties of SS by reducing the ROS<sup>13,26,28-31</sup>. A research has also shown that the reduction in the level of ROS occurred through the antioxidant activity of SS in the oocyte maturation medium<sup>26</sup>. Also Uhm *et al* demonstrated that adding SS to swine embryo culture medium caused reduction of ROS and glutathione peroxide level was elevated<sup>30</sup>. Recently, Xiong *et al* observed that the activity of glutathione peroxidase of oocytes and DNA integrity of cumulus cells increased significantly in the presence of SS during IVM of oocyte<sup>23</sup>. In contrast to our results, Lizaraga *et al*, demonstrated that adequate SS concentrations added during IVM of bovine oocytes increased viability and non-apoptotic cumulus cells<sup>24</sup>. Moreover, they revealed that neither nuclear maturation nor the developmental capacity of oocytes was modified by SS supplementation during IVM<sup>24</sup>. These differences with the results of our study may be related to the use of denuded oocytes in our study instead of cumulus oocyte complexes or different types of mammalian species.

Our results also showed the same positive effect of SS on the developmental rates of embryo to blastocyst stage. It is suggested that the SS improved the embryo quality by alteration in the ROS level during embryo culture. It was shown that adding 10 ng/ml of SS to the culture medium of bovine oocyte improved the embryo quality and caused an increase in hatching rate of embryos<sup>24</sup>. Also SS may affect embryo development through intracellular signaling by activation of phosphoinositide 3-kinase/Protein kinase B (PKB), Akt (PI3K/AKT), and extracellular signal-regulated kinases<sup>31</sup>.

Another factor contributing to the effect of SS on oocytes development is reduction of oocyte apoptosis. Some studies have shown positive effects of SS on the inhibition of apoptosis pathway in other cell types<sup>32-34</sup>. Our molecular analysis in this study indicated that in both IVM groups, *TFAM* gene expression of MII oocytes had decreased in comparison with *in vivo* obtained oocytes. This suggests the effect of culture medium conditions on the incidence of *TFAM* gene expression, which could influence the mitochondrial biogenesis of oocytes during culture period. On the other hand, previous studies have shown a direct relationship between diminished mtDNA levels and reduced number of mitochondria in poor quality oocytes<sup>13</sup>. Thus, the reduction of expression of *TFAM* in both *in vitro*

groups can be synonymous to decreased development of these oocytes. In this regard, Ghorbanmehr *et al* also showed that the obtained mitochondrial DNA (mtDNA) copy number of oocytes matured in the culture medium in comparison to the oocytes obtained from *in vivo* conditions can be due to the poor quality of these oocytes; moreover, in the oocyte treated with SS, this parameter (mtDNA) was improved and reached to the same value similar to *in vivo* group<sup>13</sup>. The SS may exert its effects through affecting the nuclear respiratory factor 1 as the main regulator of *TFAM* gene expression. For example, in hepatic cells of mice, it has been shown that the overexpression of *NRF-1* leads to an increase in the expression of *TFAM* and mtDNA content<sup>35</sup>. Another possibility is that the SS can indirectly influence the expression of *TFAM* by affecting the incidence of other factors such as miRNAs<sup>36</sup>. To prove these suggestions, more studies are required.

### Conclusion

In conclusion, supplementation of oocyte maturation culture media with SS improved the development rate of oocytes and embryo and also enhanced *TFAM* gene expression in MII oocytes that could affect the mitochondrial biogenesis of oocytes.

### Acknowledgement

This work was supported by Tarbiat Modares University of Medical Sciences as MSc thesis. The authors clarify there is no any conflict of interest.

### References

1. Hatirnaz S, Ata B, Hatirnaz ES, Dahan MH, Tannus S, Tan J, et al. Oocyte in vitro maturation: A systematic review. *J Turkish Soc Obstet Gynecol* 2018;15:112-25.
2. Costello MF, Garad RM, Hart R, Homer H, Johnson L, Jordan C, et al. A review of second- and third-line infertility treatments and supporting evidence in women with polycystic ovary syndrome. *Med Sci (Basel)* 2019;7(7):75.
3. Abbara A, Clarke SA, Dhillon WS. Novel concepts for inducing final oocyte maturation in in vitro fertilization treatment. *Endocr Rev* 2018;39(5):593-628.
4. Creux H, Monnier P, Son WY, Buckett W. Thirteen years' experience in fertility preservation for cancer patients after in vitro fertilization and in vitro maturation treatments. *J Assist Reprod Genet* 2018;35(4):583-92.
5. Son WY, Henderson S, Cohen Y, Dahan M, Buckett W. Immature oocyte for fertility preservation. *Front Endocrinol* 2019;10:464.
6. Sato A, Sarentonglaga B, Ogata K, Yamaguchi M, Hara A, Atchalalt K, et al. Effects of insulin-like growth factor-1 on the in vitro maturation of canine oocytes. *J Reprod Dev* 2018;64(1):83-8.
7. Alvarez GM, Barrios Exposito MJ, Elia E, Paz D, Morado S, Cetica PD. Effects of gonadotrophins and insulin on glucose uptake in the porcine cumulus-oocyte complex during IVM. *Reprod Fertil Dev* 2019.

8. Asaadi A, Kafi M, Atashi H, Azari M, Hostens M. Frozen-thawed ampullary cell monolayer improves bovine embryo in vitro development and quality. *Zygote* 2019; 27(5):337-46.
9. Heydarnejad A, Ostadhosseini S, Varnosfaderani SR, Jafarpour F, Moghimi A, Nasr-Esfahani MH. Supplementation of maturation medium with CoQ10 enhances developmental competence of ovine oocytes through improvement of mitochondrial function. *Mol Reprod Devel* 2019;86(7):812-24.
10. John JC. Mitochondria and female germline stem cells, a mitochondrial DNA perspective. *Cells* 2019;8(8):852.
11. Schatten H, Sun QY, Prather R. The impact of mitochondrial function/dysfunction on IVF and new treatment possibilities for infertility. *Reprod Biol Endocrinol* 2014;12:111.
12. Kasapoglu I, Seli E. Mitochondrial dysfunction and ovarian aging. *Endocrinology* 2020;161(2):bqaa001.
13. Ghorbanmehr N, Salehnia M, Amooshahi M. The effects of sodium selenite on mitochondrial DNA copy number and reactive oxygen species levels of in vitro matured mouse oocytes. *Cell J* 2018;20(3):396-402.
14. Sharma N, Pasala MS, Prakash A. Mitochondrial DNA: epigenetics and environment. *Environ Mol Mutagen* 2019;60(8):668-2.
15. Farge G, Falkenberg M. Organization of DNA in mammalian mitochondria. *Int J Mol Sci* 2019;20(11):2770.
16. Ngo HB, Lovely GA, Phillips R, Chan DC. Distinct structural features of TFAM drive mitochondrial DNA packaging versus transcriptional activation. *Nat Commun* 2014;5:3077.
17. Ghaffari Novin M, Noruzinia M, Allahveisi A, Saremi A, Fadaei Fathabadi F, Mastery Farahani R, et al. Comparison of mitochondrial-related transcriptional levels of TFAM, NRF1 and MT-CO1 genes in single human oocytes at various stages of the oocyte maturation. *Iran Biomed J* 2015;19(1):23-8.
18. Nie J, Yan K, Sui L, Zhang H, Zhang H, Yang X, et al. Mogroside V improves porcine oocyte in vitro maturation and subsequent embryonic development. *Theriogenology* 2020;141:35-40.
19. Dai X, Qiu L, Zhao B, Gao Y, Mu Y, Chu Z, et al. Melatonin ameliorates the fertilization capacity of oocytes exposed to 17alpha-ethynylestradiol. *Reprod Toxicol* 2020;93:61-7.
20. Venditti P, Meo SD. The role of reactive oxygen species in the life cycle of the mitochondrion. *Int J Mol Sci* 2020;21(6):2173.
21. Sena LA, Chandel NS. Physiological roles of mitochondrial reactive oxygen species. *Mol Cell* 2012;48(2):158-67.
22. Yang S, Lian G. ROS and diseases: role in metabolism and energy supply. *Mol Cell Biochem* 2020;467(1-2):1-12.
23. Xiong X, Lan D, Li J, Lin Y, Li M. Selenium supplementation during in vitro maturation enhances meiosis and developmental capacity of yak oocytes. *Anim Sci J* 2018;89(2):298-306.
24. Lizarraga RM, Anchordoquy JM, Galarza EM, Farnetano NA, Carranza-Martin A, Furnus CC, et al. Sodium selenite improves in vitro maturation of bos primigenius taurus oocytes. *Biol Trace Elem Res* 2020;197(1):149-58.
25. Malekzadeh M, Salehnia M, Zavareh S. Effect of sodium selenite on the apoptotic genes expression in in vitro matured mouse metaphase II oocyte. *Veter Res Forum* 2020:(In press).
26. Amoushahi M, Salehnia M, Ghorbanmehr N. The mitochondrial DNA copy number, cytochrome c oxidase activity and reactive oxygen species level in metaphase II oocytes obtained from in vitro culture of cryopreserved ovarian tissue in comparison with in vivo-obtained oocyte. *J Obstet Gynaecol Res* 2018;44(10):1937-46.
27. Ghaemi SR, Salehnia M, Valojerdi MR. The effect of progesterone and exogenous gonadotropin on preimplantation mouse embryo development and implantation. *Exp Anim* 2008;57(1):27-34.
28. Abedelahi A, Salehnia M, Allameh AA, Davoodi D. Sodium selenite improves the in vitro follicular development by reducing the reactive oxygen species level and increasing the total antioxidant capacity and glutathione peroxidase activity. *Hum Reprod* 2010;25(4):977-85.
29. Abedelahi A, Salehnia M, Allameh AA. The effects of different concentrations of sodium selenite on the in vitro maturation of preantral follicles in serum-free and serum supplemented media. *J Assist Reprod Genet* 2008;25(9-10):483-8.
30. Uhm SJ, Gupta MK, Yang JH, Lee SH, Lee HT. Selenium improves the developmental ability and reduces the apoptosis in porcine parthenotes. *Mol Reprod Devel* 2007;74(11):1386-94.
31. Liu H, Li X, Qin F, Huang K. Selenium suppresses oxidative-stress-enhanced vascular smooth muscle cell calcification by inhibiting the activation of the PI3K/AKT and ERK signaling pathways and endoplasmic reticulum stress. *J Biol Inorg Chem* 2014;19(3):375-88.
32. Ren Z, Fan Y, Zhang Z, Chen C, Chen C, Wang X, et al. Sodium selenite inhibits deoxynivalenol-induced injury in GPX1-knockdown porcine splenic lymphocytes in culture. *Sci Rep* 2018;8(1):17676.
33. Colle D, Santos DB, de Souza V, Lopes MW, Leal RB, de Souza Brocardo P, et al. Sodium selenite protects from 3-nitropropionic acid-induced oxidative stress in cultured primary cortical neurons. *Mol Biol Rep* 2019; 46(1):751-62.
34. Yang L, Cai YS, Xu K, Zhu JL, Li YB, Wu XQ, et al. Sodium selenite induces apoptosis and inhibits autophagy in human synovial sarcoma cell line SW982 in vitro. *Mol Med Rep* 2018;17(5):6560-8.
35. Piantadosi CA, Suliman HB. Mitochondrial transcription factor A induction by redox activation of nuclear respiratory factor 1. *J Biol Chem* 2006;281(1):324-33.
36. Battaglia R, Vento ME, Ragusa M, Barbagallo D, La Ferlita A, Di Emidio G, et al. MicroRNAs are stored in human MII oocyte and their expression profile changes in reproductive aging. *Biol Reprod* 2016;95(6):131.