



Molecular Mechanisms of Anti-Inflammatory Activities of the Extracts of *Ocimum gratissimum* and *Thymus vulgaris*

Ige Francis Olaoye ¹, Babatunde Joseph Oso ^{1*}, and Adepeju Aberuagba ²

1. Department of Biochemistry, McPherson University, Seriki Sotayo, Ogun State, Nigeria

2. Department of Biochemistry, University of Ilorin, Ilorin, Kwara State, Nigeria

Abstract

Background: A large body of literature suggests that the extracts of *Ocimum gratissimum* (*O. gratissimum*) and *Thymus vulgaris* (*T. vulgaris*) play protective roles against various inflammatory disorders. However, the possible mechanism of action with reference to the interactions of their respective phytochemical compositions with pro-inflammatory mediators as the indication of their therapeutic effects is less clear. Therefore, the immunomodulatory properties of *O. gratissimum* and *T. vulgaris* were investigated in this study.

Methods: The *in vitro* lipoxygenase inhibitory potentials of methanolic extracts of the selected plants were assessed through colorimetric analysis. The pharmacokinetics of some identified compounds in the botanicals were investigated *via* the Swiss ADME server while the molecular interactions of the compounds with lipoxygenase, IL-1, IL-6, TNF- α , IL-8, and CCL-2 were performed through molecular docking.

Results: The assessment of the lipoxygenase inhibition revealed the extracts could possess anti-inflammatory agents. The pharmacokinetic results of some selected compounds identified in the botanicals showed moderate toxic effects compared to indomethacin. The molecular docking study substantiated the report of the *in vitro* analysis as indicated in the binding score of all the selected compounds compared to indomethacin.

Conclusion: The phytochemical components of the extracts of *O. gratissimum* and *T. vulgaris* could be effective as anti-inflammatory agents that could be explored in preventing disorders associated with excessive activities of pro-inflammatory mediators.

Avicenna J Med Biotech 2021; 13(4): 207-216

Keywords: Anti-inflammatory agents, Lipoxygenase, Ocimum, Phytochemicals, Plant extracts, Thymus plant

Introduction

One of the most promising recent alternatives to classical treatment is the use of immunomodulators for the prevention of diseases that could be associated with dysregulated inflammatory responses. Numerous clinical documents evidenced that many pathological conditions closely related to impaired inflammatory responses mediated by cytokines and chemokines play a role in the pathogenesis of many infections such as nephritis, rheumatoid arthritis, uveitis, early stages of insulin-dependent diabetes mellitus, and thyroiditis ^{1,2}. Similarly, high expression levels of IL-1 β , IFN- γ , IP-10, and monocyte chemoattractant protein 1 (or CCL-2) have been observed in COVID-19 patients with the severity of the disease corresponding to the serum levels of IL-2R and IL-6 in some patients ³. Other reports revealed that COVID-19 patients in the Intensive Care Unit

(ICU) had high serum levels of granulocyte colony-stimulating factor, IP-10, MCP-1, macrophage inflammatory protein-1A, and TNF- α , suggesting that cytokine storm positively correlates with the disease severity. One of the major concerns that could be associated with the down-regulation of inflammatory activities is the suppression of the immune response to the pathogenic organisms; however, regulation of pro-inflammatory activities through inhibition of the biological effects of pro-inflammatory cytokines could reduce pathological deterioration associated with inflammatory disorders. Additionally, targeting the enzymes of the eicosanoid pathway such as Cyclooxygenase 2 (COX-2) and Lipoxygenase (LOX) could therapeutically provide benefit in the context of pathogenesis of certain inflammatory diseases as these enzymes play important

roles in pathophysiology following pathogenic infection and release of potent pro-inflammatory cytokines and chemokines⁴. Dietary modulation of the inflammatory responses and immune capacity are important tools to cope with twin maladies of excessive inflammatory reactions and suppressed immune system.

Research has focused on the exploration of efficient phytochemicals that can be used for the prevention of diseases and/or treatment purposes. Certain plant parts have been reported to inhibit the activities of pro-inflammatory mediators^{5,6}. Examples of such botanicals include the leaves of *Ocimum gratissimum* (*O. gratissimum*) and *Thymus vulgaris* (*T. vulgaris*), well-known herbs in the tropical and subtropical regions widely used as medicinal plants and food additives due to their health benefit potentials⁷. Despite frequent and regular use of these herbs, the possible mechanism of actions with reference to the interactions of their respective phytochemical compositions with pro-inflammatory mediators, indicating their therapeutic effects, is less clear. Thus, this study investigated the putative molecular mechanism of anti-inflammatory effects of *O. gratissimum* and *T. vulgaris*.

Materials and Methods

Chemicals

All the chemicals used were of analytical grade. Methanol, sodium dihydrogen phosphate, and disodium hydrogen phosphate were products of Guangzhou JHD Chemical Reagent Co., Ltd. Guangzhou, China.

Extracts preparation

The leaves of *O. gratissimum* and *T. vulgaris* were obtained from Ogunmakin market, Obafemi/Owode, Ogun State, Nigeria and authenticated at the Department of Biochemistry, McPherson University, Seriki Sotayo, Ogun State, Nigeria. The samples were air-dried at room temperature of 30±1 °C. Exactly 50 g of each dried samples were pulverized and soaked in 100 ml of absolute methanol for 48 hr at room temperature of 30±1 °C. The samples were filtered in Whatman TM No 1 and the filtrate was allowed to stand at room temperature of 30±1 °C for the removal of the solvent. The obtained extracts were used for the analysis of lipoxigenase inhibitory potential.

Lipoxigenase inhibitory potential

The lipoxigenase inhibitory potentials of varying concentrations (between 20 µg/ml and 100 µg/ml) of methanolic extracts of *O. gratissimum* and *T. vulgaris* were determined as described by Shinde *et al* and Sorkun *et al*^{8,9}, using linoleic acid as the substrate and crude lipoxigenase prepared from soybean as previously reported by Oso and Karigidi¹⁰ as the source of the enzyme. Accurately, 0.1 ml of each extract was added to a test tube containing 0.5 ml of 0.1 M phosphate buffer (pH=9.0) and 150 µl of the enzyme, lipoxigenase. The mixture was allowed to incubate at 29 °C for 5 min. Afterwards, 0.5 ml of 0.6 mM linoleic

acid solution was added to the mixture. The absorbance was measured at 234 nm. The results were presented as the values of percentage inhibition of lipoxigenase activity.

Phytochemical identification and ADMET properties of phytochemical contents

The major phytochemical contents of *O. gratissimum* and *T. vulgaris* were retrieved from Dr Duke's Phytochemical and Ethnobotanical Databases. Theoretical physicochemical properties of the identified compounds, as well as their corresponding ADME parameters, pharmacokinetic properties, drug-like nature, and medicinal chemistry friendliness of the selected compounds were predicted using the Swiss ADME server (<http://www.swissadme.ch/index.php>)¹¹.

Molecular docking

The 3D X-ray crystal structure of lipoxigenase was retrieved from the Protein Data Bank (<https://www.rcsb.org/>) with PDB ID: 3D3L. The crystal structure was prepared by removing the existing ligand using the DS BIOVIA tool. Similarly, the 3D crystal structures of the selected cytokines of IL-1, IL-6, TNF-α, IL-8, and CCL-2 with PDB IDs: 3NJ5, 4NI7, 1TNR, 6N2U, and 4ZK9, respectively were obtained and prepared for molecular docking. The 3D structure of identified major compounds in *O. gratissimum* and *T. vulgaris* were obtained from Zinc15 (<http://zinc15.docking.org>) in SDF format¹² and converted to PDB format using BIOVIA DS Visualizer. Each protein with all ligands on PyRx virtual screening was docked and the respective binding scores were obtained for each ligand¹³. The docked files for each protein-ligand interaction were viewed using the DS BIOVIA software tool to obtain the 2D and 3D views of the complexes. All the docked cytokines were combined with all ligands separately in PDB format using PyMOL ver. 1.1eVal (DeLano Scientific LLC, USA) for molecular dynamics study.

Molecular dynamics simulation

The conformational stability of the lipoxigenase-ligand complexes and the pro-inflammatory cytokines-ligand complexes obtained from the molecular docking was assessed. This assessment was done using molecular dynamics simulations analysis performed through iMODS server (<http://imods.chaconlab.org>) by normal Mode Analysis (NMA) in internal coordinates (Torsional space) predicting properties such as deformability, mobility profiles, eigenvalues, variance, co-variance map and elastic network of the protein-ligand interactions¹⁴.

Statistical analysis

The results were analyzed using a One-way Analysis of Variance (ANOVA) for mean differences among the various extracts followed by Duncan's multiple range tests for post hoc comparisons at p<0.05 and presented as means±standard deviation of three determinations.

Results and Discussion

In vitro lipoxygenase inhibition study

Inflammation is known as a normal biological process as a result of tissue injury, viral/microbial pathogen infection, and chemical irritation. Over-expression of LOX and the pro-inflammatory metabolites of arachidonic acid such as leukotrienes have been linked to many human pathological conditions such as inflammation, cardiovascular diseases, and cancer^{15,16}. The result of lipoxygenase percentage inhibition potential of the plant extracts compared to indomethacin reveals that *O. gratissimum* showed the least potential while indomethacin exhibited the highest potential at all doses (Figure 1). Indomethacin, a non-steroidal anti-inflammatory drug, is widely used in the management of inflammatory diseases¹⁷. However, the *in vitro* study revealed that indomethacin has higher percentage inhibition towards LOX compared to the two plant extracts. The extracts of *O. gratissimum* and *T. vulgaris* showed inhibition properties at all doses suggesting their anti-inflammatory potential which could inhibit the rate-limiting step in arachidonic acid metabolism thus reducing the synthesis of leukotrienes. This finding corresponds to the previous reports of Wei and Shibamoto¹⁸ on LOX inhibitory effects by *T. vulgaris* and *O. gratissimum*.




Figure 1. Lipoxygenase inhibitory potentials of *O. gratissimum* and *T. vulgaris* comparable to indomethacin.

Assessment of molecular mechanism of the LOX inhibitory potentials of *O. gratissimum*, *T. vulgaris*, and indomethacin through structure-based virtual screening revealed that indomethacin had the highest binding affinity due to its least binding score which is significantly different from all the phytochemicals identified in *O. gratissimum* and *T. vulgaris* through Dr Duke's Phytochemical and Ethnobotanical Databases (Table 1). However, all the investigated compounds had good interactions with LOX based on their negative values¹⁹.

These binding interactions are mainly due to pi-interactions between compounds and LOX as well as hydrogen bonds in all the evaluated compounds (Figure 2). Although the number of hydrogen bond interactions is low compared to the binding score, the high binding score could be as a result of pi-interactions from Val190, Leu178, Leu227, Leu361, and Phe352, His360, and His365 via promotion of better position and organization of the proteins^{20,21}. The selected compounds in *T. vulgaris* (Cymene, terpinene, and thymol) showed no significant difference in the binding score and interacted with His360 and His365. Conversely, indomethacin did not interact with His360 and His365; however, its high binding score with LOX could be due to interactions with several residues through covalent interactions such as a carbon-hydrogen bond, pi-pi stack, pi-alkyl, van der Waals with polar amino acids such as Gln435, His462, Asp512, and Arg641 and non-polar amino acids of Leu178, Ala433, Ala434, Leu447, Leu453, Pro456, Ala461, Leu465, Leu507, and Cys508. The interactions such as pi-pi stack, pi-alkyl, van der Waals between the hydrophobic residues and LOX could be responsible for the stability and activity of the proteins²².

Pharmacokinetic study

The *in silico* pharmacological assessments of the drug-likeness of the identified phytochemicals were carried out through SwissADME server. The solubility features revealed that all the compounds had consensus Log p-values less than 5 like their lipophilicity feature.

Table 1. Molecular interactions of some selected phytochemical components of *T. vulgaris* and *O. gratissimum* with lipoxygenase

Phytochemicals	Zinc15 ID	Binding score (kcal/mol)	Interacting residues
<i>O. gratissimum</i>			
Caryophyllene oxide	2039864	-6.50±0.06 ^a	Phe352
Cis-ocimene	1531619	-4.93±0.03 ^c	Val190, Leu194, Phe352, His360, Leu361, Ile593, Leu597
Eugenol	1411	-6.10±0.39 ^a	Leu227, Tyr230, Gly309, Lys310, Gln509, Trp647
<i>T. vulgaris</i>			
Cymene	968246	-5.43±0.15 ^b	Val190, His360, Leu361, His365
Terpinene	967594	-5.37±0.15 ^b	Val190, His360
Thymol	967597	-5.47±0.20 ^b	Val190, Glu356, His360, His365, Leu597
Indomethacin			
-	601283	-7.70±0.06 ^d	Leu178, Ala433, Ala434, Gln435, Leu447, Leu453, Pro456, Ala461, His462, Leu465, Leu507, Cys508, Asp512, Arg641

a-d: Values with different superscripts are significantly different at p<0.05.




Figure 2. 3D illustration of the molecular interactions of (i) caryophyllene oxide, (ii) cymene, (iii) eugenol, (iv) cis-ocimene, (v) terpinene, (vi) thymol, and (vii) indomethacin with lipoygenase.

Interestingly, all the compounds were soluble in water by Log S (ESOL), Log S (Ali), and Log S (SILICOS-IT) analysis except terpinene which was moderately soluble by Log S (Ali) analysis (Table 2). The results revealed equal lipophilicity suggesting a better absorption and permeability of the selected compounds in living organisms due to less than 5 consensus Log P values²³. In line with this, the prediction of water solubility using Sorkun and Khetan's⁹ report on three different models showed that none of the evaluated compounds' Log S values exceeded -4.00 except terpinene with Log S (Ali) value of -4.220 (0.008 mg/ml) suggesting better absorption and metabolism as a result of excellent water solubility. Also, caryophyllene oxide, ocimene, and eugenol showed high gastrointestinal absorption as revealed by their low affinity for permeability-glycoprotein except for ocimene. Meanwhile, cymene, terpinene, and thymol showed low gastrointestinal absorption except for thymol with a low affinity for permeability glycoprotein.

Additionally, the results of the pharmacokinetics and the interactions of the selected phytochemicals with P-glycoprotein (P-gp) and cytochromes P450 (CYPs) and the predicted drug-likeness are presented in table 3. Caryophyllene oxide and eugenol showed high GI absorption except for ocimene while cymene and terpinene showed low GI absorption except for thymol. All selected compounds irrespective of the plant's source can cross the blood-brain barrier and showed skin permeation values greater than -2.5 cm/s while none served as the P-gp substrate. In a similar pattern, all the compounds showed no inhibitory effect on CYP2C9 and CYP2C19 suggesting that all compounds except caryophyllene oxide could not hinder the synthesis of epoxyeicosatrienoic acids (an anti-inflammatory agent) and terminate the metabolism of therapeutic drugs like anti-ulcer, anti-malaria, anti-convulsant, anesthetic, and sedative drugs²⁴. In an almost similar pattern, only cymene could inhibit CYP2D6 and stop the metabolism of anti-hypersensitive and anti-arrhythmic drugs as well as β -blockers and anti-depressants. The drug-likeness appraised based on five different rule-based filters showed that none of the selected compounds violated the Lipinski, Egan, and Veber rules. However, all the compounds violated the Ghose and Muegge rules with lower molecular weight except for caryophyllene oxide and eugenol with molecular weights greater than 160 g/mol¹¹.

The radar plots of all the compounds were within the physicochemical ranges which indicate the suitability of the compounds through oral administration (Figure 3). This computation largely correlates with their observed lipophilicity and water-solubility properties.

Molecular docking study of cytokines

A relationship has been established between the pathogenic expressions of LOX and cytokine produc-

Table 2. Solubility characteristics of the selected phytochemicals

Photochemicals	Caryophyllene oxide	Cymene	Eugenol	Ocimene	Terpinene	Thymol
Lipophilicity						
Log P _{ow} (iLOGP)	3.10	2.51	2.37	2.80	2.73	2.32
Log P _{ow} (XLOGP3)	3.56	4.10	2.27	4.26	4.50	3.30
Log P _{ow} (WLOGP)	3.94	3.12	2.13	3.48	3.31	2.82
Log P _{ow} (MLOGP)	3.67	4.47	2.01	3.56	3.27	2.76
Log P _{ow} (SILICOS-IT)	4.07	3.29	2.48	2.88	2.95	2.79
Consensus Log P _{ow}	3.67	3.50	2.25	3.40	3.35	2.80
Water solubility						
Log S (ESOL)	-3.450	-3.630	-2.460	-3.170	-3.450	-3.190
Solubility (mg/ml)	0.078	0.031	0.569	0.092	0.048	0.097
Class	Soluble	Soluble	Soluble	Soluble	Soluble	Soluble
Log S (Ali)	-3.510	-3.810	-2.530	-3.970	-4.220	-3.400
Solubility (mg/ml)	0.068	0.021	0.490	0.015	0.008	0.060
Class	Soluble	Soluble	Soluble	Soluble	Insoluble	Soluble
Log S (SILICOS-IT)	-3.510	-3.570	-2.790	-2.040	-2.230	-3.010
Solubility (mg/ml)	0.068	0.036	0.265	1.240	0.806	0.146
Class	Soluble	Soluble	Soluble	Soluble	Soluble	Soluble

Table 3. The pharmacokinetics properties of the selected compounds

	Caryophyllene oxide	Cymene	Eugenol	Ocimene	Terpinene	Thymol
Pharmacokinetics						
GI absorption	High	Low	High	Low	Low	High
BBB permeant	Yes	Yes	Yes	Yes	Yes	Yes
P-gp substrate	No	No	No	No	No	No
CYP1A2 inhibitor	No	No	Yes	No	No	Yes
CYP2C19 inhibitor	Yes	No	No	No	No	No
CYP2C9 inhibitor	Yes	No	No	No	No	No
CYP2D6 inhibitor	No	Yes	No	No	No	No
CYP3A4 inhibitor	No	No	No	No	No	No
Log K _p (cm/s) *	-5.12	-4.21	-5.69	-4.11	-3.94	-4.87
Drug-likeness						
Lipinski	Yes	Yes	Yes	Yes	Yes	Yes
Ghose	Yes	No	Yes	No	No	No
Veber	Yes	Yes	Yes	Yes	Yes	Yes
Egan	Yes	Yes	Yes	Yes	Yes	Yes
Muegge	No	No	No	No	No	No
Bioavailability score	0.55	0.55	0.55	0.55	0.55	0.55

* Skin permeation.




Figure 3. Suitable physicochemical space for oral bioavailability of (i) caryophyllene oxide, (ii) cymene, (iii) eugenol, (iv) cis-ocimene, (v) terpinene, and (vi) thymol.

tion as LOX inhibitory agents had been shown to ameliorate Tumor Necrosis Factor- α (TNF- α)-induced cytokine and chemokine release^{25,26}. This indicates that LOX inhibitors may be better candidates for the treatment of inflammatory diseases that are associated with cytokine storms. The molecular docking analyses between the selected phytochemicals and some pro-inflammatory mediators such as interleukin 1, interleukin 6, TNF- α , interleukin 8, and monocyte chemoattractant protein-1 (or C-C chemokine ligand 2, CCL-2)

revealed that the phytochemicals had a reasonable binding affinity for the pro-inflammatory mediators (Table 4) suggesting the modulatory potential of the compounds. These inflammatory mediators play important roles in the natural responses to pathogenic threats and the development of pathological disorders which could lead to chronic inflammation characterized by an exaggerated cytokine "storm" and the release of damage-associated molecular patterns²⁷. It could be concluded from figure 4 and table 4 that the

Anti-Inflammatory Activities of the Extracts of *Ocimum gratissimum* and *Thymus vulgaris*

Table 4. Molecular interactions of some selected phytochemical components of *T. vulgaris* and *O. gratissimum* with pro-inflammatory cytokines

Cytokines	Phytocompounds	Zinc15 ID	Binding score (kcal/mol)	Interacting residues
IL-1				
	Caryophyllene oxide	2039864	-5.45±0.05 ^a	Leu113, Ser125, Phe128, Pro129, Gly130, Phe132
	Cymene	968246	-4.85±0.05 ^b	Leu113, Glu124, Pro129, Gly130, Phe132
	Eugenol	1411	-4.80±0.00 ^b	Leu113, Tyr111, Glu124, Pro129, Gly130
	Cis-ocimene	1531619	-3.70±0.00 ^c	Glu124, Pro129, Gly130, Phe132
	Terpinene	967594	-4.45±0.05 ^d	Glu124, Pro129, Gly130, Phe132
	Thymol	967597	-4.80±0.01 ^b	Leu113, Glu124, Pro129, Phe132
IL-6				
	Caryophyllene oxide	2039864	-5.50±0.00 ^a	Glu23, Ile25, Asp26, Lys27, Arg182
	Cymene	968246	-4.43±0.03 ^b	Leu19, Ser22, Glu23, Asp26, Lys27
	Eugenol	1411	-4.77±0.03 ^c	Val96, Pro141, Ala145, Leu148
	Cis-ocimene	1531619	-3.80±0.00 ^d	Val96, Ala145, Leu148,
	Terpinene	967594	-4.20±0.06 ^e	Lys27, Tyr31, Val96
	Thymol	967597	-4.70±0.10 ^c	Leu92, Glu95
TNF-α				
	Caryophyllene oxide	2039864	-5.15±0.15 ^a	Phe74, Ile103, Ser105, Gln107, Val112, Leu114, Ala138, Phe139, Gln140, Leu141, Phe169
	Cymene	968246	-4.80±0.00 ^b	Ala30, Phe169
	Eugenol	1411	-4.35±0.05 ^c	Ala30, His32, Phe53, Tyr76, Phe169
	Cis-ocimene	1531619	-3.80±0.00 ^d	Ala30, Tyr76, Phe169
	Terpinene	967594	-4.15±0.05 ^e	Ala30, His32, Phe53, Phe169
	Thymol	967597	-4.55±0.05 ^f	Ala30, Tyr76, Phe169
IL-8				
	Caryophyllene oxide	2039864	-5.10±0.10 ^a	His16
	Cymene	968246	-5.55±0.05 ^b	His16
	Eugenol	1411	-5.55±0.05 ^b	His16
	Cis-ocimene	1531619	-5.10±0.10 ^a	His16
	Terpinene	967594	-5.05±0.05 ^a	His16
	Thymol	967597	-5.30±0.10 ^c	His16, Lys62
CCL-2				
	Caryophyllene oxide	2039864	-4.60±0.10 ^a	Ile20, Ser21, Asp62, Trp59, Ser63
	Cymene	968246	-5.60±0.20 ^b	Ile20, Val22, Leu25, Ala53, Trp59, Val60
	Eugenol	1411	-5.45±0.25 ^b	Ile20, Leu25, Ser27, Phe43, Ile51, Ala53, Val60, Ser63, Leu67
	Cis-ocimene	1531619	-4.70±0.00 ^a	Ile20, Arg24, Leu25, Ile51, Trp59
	Terpinene	967594	-4.80±0.10 ^a	Leu25, Ala53, Trp59, Val60
	Thymol	967597	-5.05±0.05 ^c	Thr16, Ile20, Arg24, Thr45, Val47, Ile51, Trp59

a-f: Values with different superscripts are significantly different at p<0.05.

significant low binding scores observed in cis-ocimene and terpinene interaction with IL-1 might be due to absence of Leu113 in the interacting site.

The residue Pro129 was shown to be a probable and important residue required for IL-1 activity as it was found to interact with all the selected phytocompounds (Figure 4). Caryophyllene oxide had the highest binding affinity with IL-1, IL-6, and TNF-α. This could be due to the contribution of Van der Waals forces to the interaction between the cytokines. IL-1, IL-6, and TNF-α are potent cytokines involved in various human diseases²⁸. Moreover, TNF-α could also stimulate the expression of IL-1 and IL-6, causing the associated pathological disorders²⁹. Various studies had provided a rationale to explore their respective targets in the development of therapy for certain diseases associated with autoimmune disorders³⁰. This could prevent the

respective cascade of signaling processes through the JAK/STAT3 activation pathway and transcription of genes of inflammatory proteins including cytokines, receptors, and protein kinases³¹. Eugenol had the best binding affinity followed by thymol with IL-8. His16 was identified as an important residue in the interaction of the chemokine with the phytocompounds. IL-8, a chemoattractant usually released by macrophages, induces inflammation through the recruitment of neutrophils and other immune cells to the site of injury or infection³². IL-1 and TNF-α could also induce IL-8 production from a variety of cells³³. Cymene and eugenol depicted the highest binding affinity towards monocyte C-C chemokine Ligand 2 (CCL-2) followed by thymol. The observed binding scores might be due to the pi-pi stacking interactions of the compounds with CCL-2 through Ile20, Ala53, and Val60 (Figure




Figure 4. 3D illustration of the molecular interactions of (i) caryophyllene oxide, (ii) cymene, (iii) eugenol, (iv) cis-ocimene, (v) terpinene, and (vi) thymol with, A) IL-1, B) IL-6, C) TNF- α , D) IL-8, and E) CCL-2.

4). Similarly, the affinity of thymol with CCL-2 could be associated with the conventional hydrogen bond formed between thymol and CCL-2³⁴. Excessive inflammation connected with various diseases such as autoimmune disease, atherosclerosis, and neurological disorders had been linked with the production of CCL-2 and its interaction with its main chemokine receptor CCR2³⁵. CCL2 has been assumed to be a therapeutic target as its inhibition was found to decrease lung inflammation, metastases, and atherosclerosis in animals³⁶.

Molecular dynamics study

The effects of the identified phytochemicals on the flexibility of the cytokines and the chemokines upon binding were assessed with molecular dynamics simulations derived factors such as deformability, B-factor, eigenvalues, variance, covariance, and elastic factors

considered in the study (Figure 5). The interactions of the phytochemicals with the cytokines and chemokines did not induce changes in the flexibility and dynamics of the pro-inflammatory mediators. However, there were variances in the flexibility and dynamics of the pro-inflammatory cytokines and chemokines. The main-chain deformability was higher in the CCL-2 while IL-8 had the highest rigidity among the selected mediators. The energy required to deform the TNF- α is relatively high as indicated by the lowest eigenvalue (2.741269×10^{-5}) (Figure 5C). However, IL-1 had the lowest variance associated with each normal mode which is inversely related to the eigenvalue (Figure 5D). The interaction between pairs of residues of the pro-inflammatory mediators computed using the Ca Cartesian coordinates showed that the cytokines (IL-1, IL-6, and TNF- α) could experience moderately corre-

Anti-Inflammatory Activities of the Extracts of *Ocimum gratissimum* and *Thymus vulgaris*




Figure 5. Molecular dynamics simulation of (i) IL-1, (ii) IL-6, (iii) TNF- α , (iv) IL-8, and (v) CCL-2 showing the, A) Deformability, B) B-factor, C) Eigenvalues, D) Variance, E) Covariance map, and F) Elastic network.

lated motion between the residues comparable to the chemokines (IL-8 and CCL-2) (Figure 5E). The stiffness was found to be comparatively lower in IL-8 and CCL-2 as indicated by the lesser distribution of dark grey dots in the map (Figure 5F)^{14,37}.

Conclusion

This study showed that all the selected phytochemical components of *O. gratissimum* and *T. vulgaris* except cis-ocimene interacted adequately with the selected inflammatory mediators. Thus, repositioning these compounds as pro-inflammatory antagonists could be a promising strategy to alleviate inflammatory disorders that could arise from infectious diseases or dysfunc-

tional immune disorders. Further analyses are recommended to confirm the putative therapeutic effects of the selected compound.

Acknowledgement

We recognize and appreciate the assistance of the staff of the Department of Biological Sciences, MCPerson University, Seriki Sotayo.

References

1. Cheung CY, Poon LLM, Ng IHY, Luk W, Sia SF, et al. Cytokine responses in severe acute respiratory syndrome coronavirus-infected macrophages in vitro: possible relevance to pathogenesis. *J Virol* 2005;79(12):7819-26.

2. Lau SKP, Lau CCY, Chan KH, Li CPY, Chen H, Jin DY, et al. Delayed induction of proinflammatory cytokines and suppression of innate antiviral response by the novel Middle East respiratory syndrome coronavirus: implications for pathogenesis and treatment. *J Gen Virol* 2013; 94(12):2679-90.
3. Chen L, Liu HG, Liu W, Liu J, Liu K, Shang J.[Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi* 2020;6;43(0):E005. Chinese.
4. Radi ZA, Meyerholz DK, Ackermann MR. Pulmonary cyclooxygenase-1 (COX-1) and COX-2 cellular expression and distribution after respiratory syncytial virus and parainfluenza virus infection. *Viral Immunol* 2010;23(1): 43-8.
5. Amri O, Zekhnini A, Bouhaimi A, Tahrouch S, Hatimi A. Anti-inflammatory activity of methanolic extract from *Pistacia atlantica* Desf. leaves. *Pharmacog J* 2018;10(1): 71-6.
6. Oso BJ, Oyewo EB, Oladiji AT. Ethanolic, N-hexane and aqueous partitioned extracts of *Xylopiya aethiopia* fruit modulated inflammatory responses in turpentine oil induced acute inflammation in male wistar rats. *Int J Res Health Sci* 2017; 5(2):1-10.
7. Basch E, Ulbricht C, Hammerness P, Blevins A, Sollars D. Thyme (*Thymus vulgaris* L.) thymol. *J Herb Pharmacother* 2004;4(1):49-67.
8. Shinde UA, Phadke AS, Nair AM, Mungantiwar AA, Dikshit VJ, Saraf MN. Membrane stabilizing activity-a possible mechanism of action for the anti-inflammatory activity of *Cedrus deodara* wood oil Author links open overlay panel. *Fitoterapia* 1999;70(3):251-7.
9. Sorkun MC, Khetan A, Süleyman Er. AqSolDB, a curated reference set of aqueous solubility and 2D descriptors for a diverse set of compounds. *Sci Data* 2019;6(1):143.
10. Oso BJ, Karigidi KO. Inhibitory action of dried leaf of *Cassia alata* (Linn.) Roxb against lipoxygenase activity and nitric oxide generation. *Scien Agropec* 2019;10(2): 185-90.
11. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep* 2017;7:42717.
12. Sterling T, Irwin JJ. ZINC 15–Ligand discovery for everyone. *J Chem Inf Model* 2015;55(11):2324-37.
13. Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* 2010.;31(2):455-61.
14. López-Blanco JR, Aliaga JI, Quintana-Ortí ES, Chacón P. iMODS: Internal coordinates normal mode analysis server. *Nucleic Acids Res* 2014;42:W271-W276.
15. Adamek A, Jung S, Dienesch C, Laser M, Ertl G, Bauersachs J, Frantz S. Role of 5-lipoxygenase in myocardial ischemia-reperfusion injury in mice. *Eur J Pharmacol* 2007;571(1):51-4.
16. Farooqui AA, Horrocks LA, Farooqui T. Modulation of inflammation in brain: a matter of fat. *J Neurochem* 2007;101(3):577-99.
17. Regula J, Butruk E, Dekkers Cp, De Boer Sy, Raps D, Simon L, et al. Prevention of NSAID-associated Gastrointestinal Lesions: A comparison study pantoprazole versus Omeprazole. *Am J Gastroenterol* 2006;101(8):1747-55.
18. Wei A, Shibamoto T. Antioxidant/Lipoxygenase inhibitory activities and chemical compositions of selected essential oils. *J Agric Food Chem* 2010;58(12):7218-25.
19. Oso BJ, Olaoye IF. Comparative in vitro studies of anti-glycemic potentials and molecular docking of *Ageratum conyzoides* L. and *Phyllanthus amarus* L. methanolic extracts. *SN App Sci* 2020;2:629.
20. Brylinski M. Aromatic interactions at the ligand-protein interface: Implications for the development of docking scoring functions. *Chem Biol Drug Des* 2018;91(2):380-90.
21. Arthur DE, Uzairu A. Molecular docking studies on the interaction of NCI anticancer analogues with human Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit. *J King Saud University Science* 2019;31(4): 1151-66.
22. Yadav DK, Kumar S, Saloni, Misra S, Yadav L, Teli M, et al. Molecular insights into the interaction of RONS and Thieno[3,2-c]pyran analogs with SIRT6/COX-2: A molecular dynamics study. *Sci Rep* 2018;8(1):4777.
23. Fonteh P, Elkhadir A, Omondi B, Guzei I, Darkwa J, Meyer D. Impedance technology reveals correlations between cytotoxicity and lipophilicity of mono and bimetallic phosphine complexes. *Biometals* 2015;28(4):653-67.
24. Daly AK, Rettie AE, Fowler DM, Miners JO. Pharmacogenomics of CYP2C9: Functional and clinical considerations. *J Pers Med* 2018;8(1):1.
25. Zhao L, Cuff CA, Moss E, Wille U, Cyrus T, Klein EA, et al. Selective interleukin-12 synthesis defect in 12/15-lipoxygenase-deficient macrophages associated with reduced atherosclerosis in a mouse model of familial hypercholesterolemia. *J Biol Chem* 2002;277(38):35350-6.
26. Lin HC, Lin TH, Wu MY, Chiu YC, Tang CH, Hour MJ, et al. 5-Lipoxygenase inhibitors attenuate TNF- α -Induced inflammation in human synovial fibroblasts. *PLoS One* 2014;9(9):e107890.
27. Kim B, Lee Y, Kim E, Kwak A, Ryoo S, Bae SH, et al. The Interleukin-1 α precursor is biologically active and is likely a key alarmin in the IL-1 family of cytokines. *Front Immunol* 2013;4:391.
28. Lopetuso LR, Chowdhry S, Pizarro TT. Opposing functions of classic and novel IL-1 family members in gut health and disease. *Front Immunol* 2013;4:181.
29. Frangogiannis NG, Lindsey ML, Michael JH, Youker KA, Bressler RB, Mendoza LH, et al. Resident cardiac mast cells degranulate and release performed TNF- α , initiating the cytokine cascade in experimental canine myocardial reperfusion. *Circulation* 1998;98(7):699-710.

30. Masola V, Carraro A, Granata S, Signorini L, Bellin G, Violi P, et al. In vitro effects of interleukin (IL)-1 beta inhibition on the epithelial-to-mesenchymal transition (EMT) of renal tubular and hepatic stellate cells. *J Transl Med* 2019;17(1):12.
31. Wang Y, Van Boxel-Dezaire AH, Cheon H, Yang J, Stark GR. STAT3 activation in response to IL-6 is prolonged by the binding of IL-6 receptor to EGF receptor. *Proc Natl Acad Sci USA* 2013;110(42):16975-80.
32. Gauglitz GG, Finnerty CC, Herndon DN, Mlcak RP, Jeschke MG. Are serum cytokines early predictors for the outcome of burn patients with inhalation injuries who do not survive? *Crit Care* 2008;12(3):R81.
33. Kasahara T, Mukaida N, Yamashita K, Yagisawa H, Akahoshi T, Matsushima K. IL-1 and TNF alpha induction of IL-8 and monocyte chemoattractant and activating factor (MCAF) mRNA expression in a human astrocytoma cell line. *Immunology* 1991;74(1):60-7.
34. Elokely KM, Doerkse RJ. Docking challenge: protein sampling and molecular docking performance. *J Chem Inf Model* 2013;53(8):1934-5.
35. Severini C, Passeri PP, Ciotti M, Florenzano F, Possenti R, Zona C, et al. Bindarit, inhibitor of CCL2 synthesis, protects neurons against amyloid- β -induced toxicity. *J Alzheimers Dis* 2014;38(2):281-93.
36. Bonapace L, Coissieux M, Wyckoff J, Mertz KD, Varga Z, Junt T, et al. Cessation of CCL2 inhibition accelerates breast cancer metastasis by promoting angiogenesis. *Nature* 2014;515(7525):130-3.
37. Adeoye AO, Oso BJ, Olaoye IF, Tijjani H, Adebayo AI. Repurposing of chloroquine and some clinically approved antiviral drugs as effective therapeutics to prevent cellular entry and replication of coronavirus. *J Biomol Struct Dyn* 2020;1-11.