

The Phenotypic and Molecular Identification of Phyllospheric Bacteria Possessing Antimicrobial Activity from *Funtumia elastica* (Preuss) Stapf.

Bolanle A. Adeniyi ^{1*}, Mercy Ogunlana ¹, Christopher O. Igbokwe ¹, Bamidele Tajudeen ²
and Gail B. Mahady ³

1. Department of Pharmaceutical Microbiology, University of Ibadan, Nigeria

2. Nigerian Institute of Medical Research, Lagos, Nigeria

3. Department. Pharmacy Practice, College of Pharmacy, University of Illinois, Chicago, United States of America

Abstract

Background: Unlike plant phytochemicals, little has been done to explore the metabolites from phyllosphere bacterial flora, some of which enabled them to survive interspecific competition through amensalism. This study evaluated the antimicrobial activity of metabolites from Phyllospheric Bacteria (PB) isolated from *Funtumia elastica* (FE), against selected bacterial and fungal pathogens. Phenotypic and molecular methods were used to identify the isolated phyllo-microbiota.

Methods: The PB were aseptically isolated by sonication. Their metabolites were obtained from the fresh overnight culture of the organisms. The cell-free supernatants containing the metabolites were used for antimicrobial assays against the pathogens. The DNA of the bacterial isolates were isolated using a NIMR-BIOTECH DNA extraction kit, while their 16S rRNA was amplified with the primer: 799F 5'-AACACGGATTA GATACC-3', 1193R 5'-ACGTCATCCCCACCTTCC-3', using SolisFast* Master Mix, (Solis Biodyne-Estonia). The BLAST of the sequence was done from the NCBI Genbank. The PB strains identified were submitted to NCBI and accession numbers were assigned to them.

Results: The phyllosphere of FE yielded 21 bacterial isolates: 7 Gram-positives and 14 Gram-negatives. The metabolites from these isolates showed varying degrees of bioactivity against *Staphylococcus aureus* (ATCC29213), *Escherichia coli* (ATCC 25922) *Klebsiella pneumoniae* (ATCC 35659); *Trychophyton rubrum*, *Candida albicans* and *Microsporum canis*. Fifteen bioactive isolates sequenced yielded four genera, Enterobacter (*E. hormaechei* 98.44%), Bacillus (*B. cereus* 100%), Pontoea (*P. dispersa* 99.72%), Staphylococcus (*S. arlettae* 99.72%).

Conclusion: Bacteria from FE phyllosphere, produced metabolites antagonistic (*cidal*) to some human pathogens. This has great potential for possible drug discovery.

Keywords: 16S rRNA, Biotechnology, DNA, Drug discovery, Enterobacter, Microbiota, Phytochemicals

To cite this article: Adeniyi BA, Ogunlana M, Igbokwe CO, Tajudeen B, Mahady GB. The Phenotypic and Molecular Identification of Phyllospheric Bacteria Possessing Antimicrobial Activity from *Funtumia elastica* (Preuss) Stapf. Avicenna J Med Biotech 2024;16(3):180-186.

Introduction

In recent years, the search for and use of medicines and food supplements derived from plants has accelerated. Ethnopharmacologists, botanists, microbiologists, and natural product chemists are searching the earth for phytochemicals and "clues" that can be used to treat infectious diseases ¹. Phytochemicals are biologically active chemical compounds that occur naturally, are found in plants and are beneficial for human health ². Medicinal plants are rich in phytochemicals and can be

optimized in their structure and processed into new drugs ³. Medicinal plants commonly used in our community can be an excellent source of these new drugs ⁴. The resistance of micro-organisms to classic antibiotics and their rapid development have aroused great attention in the treatment of infectious diseases ⁵.

Plant phyllosphere consist of aerial or above-ground parts of plants and are primarily one of the most prevalent microbial habitats on earth ⁶. The phyllosphere is

* Corresponding author:
Bolanle A. Adeniyi, Ph.D.,
Department of Pharmaceutical
Microbiology, University of
Ibadan, Nigeria
Tel: +234 802 3451480
E-mail:
bolaadeniyi2022@gmail.com
Received: 23 Aug 2023
Accepted: 29 Nov 2023

dominated by leaves. The leaf surface habitat is very large with an estimated global leaf area of 508,630,100 km^2 which means 1,017,260,200 km^2 and implies a sufficiently large habitat for micro-organisms⁷. The complex composition of phyllosphere microbes is influenced by some external factors such as light, UV radiation, atmospheric temperature, less water, and nutrients, which make them adapt to harsh environmental conditions⁸.

Micro-organisms associated with plants may be endophytic or epiphytic depending on their location on the host plant. Endophytic organisms interact with and influence the internal part of plants, while epiphytic organisms interact and influence the exterior surface of plants. These organisms are not harmful to the plant but produce some useful substances that help in promoting the growth of the plant, providing resistance to pathogenic microbes and the production of secondary metabolites⁹.

Bacteria are the most abundant micro-organisms found in the phyllosphere. It is estimated that 6.4×10^8 km^2 of the global leaf surface harbours about 10^{26} bacteria, the most abundant colonizers of the phyllosphere¹⁰. The culture method is widely used for the identification of the different microbes in phyllospheres. Thompson *et al*¹¹ identified 78 different bacterial species from sugar beet. However, this cultural method is possibly inaccurate in determining diversity. This is because of the presence of uncultured organisms. The use of culture-independent methods such as 16S rRNA sequences would give a more accurate and complete overview of the structure⁸. Phenotypic characterization was based on comparing morphologic and phenotypic characteristics of type strains with that of the isolate to be identified. The 16S rRNA gene, being the highly conserved portion of the bacterial genome, is commonly used for taxonomic purposes in bacteria. The 16S rRNA is about 1550 *bp* long comprising of variable and conserved regions¹².

The average number of bacteria that are usually found on leaf surfaces is around 10^6 - 10^8 cells/cm. *Proteobacteria*, *Firmicutes*, *Bacteroides*, and *Actinobacteria* are the four major phyla that are generally found in the phyllosphere. Molecular studies have indicated that the highest bacterial inhabitants of the phyllosphere are *firmicutes* and alpha-, beta- and gamma-proteobacteria. *Cyanobacteria*, *Acidobacteria*, and *Actinobacteria* are also found to be frequently occurring in the phyllosphere. *Methylobacterium* and *Sphingomonas* are the most common genera of the class alpha-proteobacteria. Some of the factors that determine this bacterial assembly include the age of the plant, climatic conditions, the immune system of the plant, the species of plant and plant genotype and also, the soil type⁸.

Environmental variables can change, while the geographical location of the plant and the plant genotype remain constant and stable. At times, the genotype has a major influence on the microbiome composition.

However, geographical location has been reported to have the most significant impact. Some season-dependent microbial communities inhabiting the surface of perennial plants are almost the same from year to year, while others show significant variation with changing seasons¹³.

Funtumia elastica (*F. elastica*) (Preuss) Stapf (Figure 1), is from the Apocynaceae family. Its common name is "Ireh" in Yoruba, and it is a forest tree growing in West and Central Africa¹⁴. *Funtumia elastica* has a traditional history of ethnopharmacological use in the treatment of whooping cough, asthma, dysmenorrhea, wounds, and fungal infections. It is confirmed to exhibit significant antimicrobial activity against wide range of organisms such as *Bacillus subtilis*, *Candida albicans* (*C. albicans*), *Pseudomonas aeruginosa*, and *Escherichia coli* (*E. coli*), and diseases such as syphilis, gonorrhea, and hemorrhoids, amongst others¹⁵. The stem latex of *F. elastica* is used for washing wounds, its leaves to treat hemorrhoids, and its bark powder in the treatment of respiratory ailments including asthma¹⁴. It is also of great value in traditional medicine for the treatment and management of some diseases and disorders. A decoction from its bark is administered as a laxative and vermifuge. It is also included in prescriptions for problems associated with painful menstruation. The stem bark can be pounded and taken in spirits to cure hemorrhoids. The decoction from the stem bark is also used for treating chest infections like whooping cough. The decoction of its leaves is also used in treating mouth and venereal diseases¹⁶. Despite the well-known antimicrobial activities of *F. elastica*, and its readily availability, there are no published studies on phyllosphere organisms on its leaf surface. This study therefore was designed to isolate, identify, and test their activity against clinical pathogenic bacterial and fungal isolates.

Materials and Methods

Plant sample collection

The leaves of *F. elastica* were collected from the



Figure 1. *Funtumia elastica* (Preuss) Stapf plant. (Photo by C. Delnatte).

University of Ibadan Botanical Garden and aseptically transported in a sterile bag.

Isolation and identification of *F. elastica* phyllosphere bacteria

The plant leaves were aseptically collected into a pre-weighed sterile bag and then transported to the laboratory within an hour. The weight was determined. Eight grams (8 g) of the leaves were then transferred aseptically into a bottle containing 80 ml of sterile distilled water and aseptically sonicated for 10 min after which the sonicate was serially diluted into 10^{-1} , 10^{-2} , and 10^{-3} dilutions. The dilutions were inoculated into Tryptone Soy Agar (TSA) containing 0.8 ml of Nystatin in order to inhibit fungal growth. The inoculum was evenly distributed on the surface of the set agar using a sterile glass spreader. The plates were left on the bench for some minutes. They were then incubated on a transparent bench incubator, at room temperature for 4 days, close to the window for light accessibility.

Colony counts and subcultures from primary plates

The plates of the primary isolates were inspected and the colonies were counted. The morphologies of the various colonies were recorded. Isolated colonies were subcultured into fresh TSA plates without Nystatin. These plates were then incubated as described earlier. The results were then documented.

Biochemical tests on isolated organisms

Gram staining and biochemical tests were carried out and they include oxidase test, catalase test, and indole test. The growth of the organisms on differential media was also observed. These media include MacConkey agar, EMB, and MSA. The colour of colonies, texture, the appearance of colonies, and colour of plates were recorded.

Antimicrobial activity of cell-free supernatant (metabolites) of phyllospheric organisms

The antibacterial activity of the phyllospheric organisms against *Staphylococcus aureus* (*S. aureus*) (ATCC 29213), *E. coli* (ATCC 25922), and *Klebsiella pneumoniae* (*K. pneumoniae*) (ATCC 35659) was determined. The phyllospheric isolates were each cultured into 5 ml of sterile TSB and incubated for 48 hr. 20 ml of Muller Hinton agar (for bacteria), and Sabouraud Dextrose Agar (for fungi), were dispensed into MacCartney bottles and then sterilized. The molten agar was poured aseptically into sterile petri dishes and allowed to set. Pure colonies of the test organisms were resuspended in sterile distilled water, and standardized to 0.5 McFarland standard. A microbial lawn of the test organisms was made on the surface of the dried agar plates. A 6 mm diameter cork borer was used to bore wells on the plates. After 48 hr, of incubation of the phyllosphere isolates, about 2 ml of the broth culture of each organism, was dispensed into sterile cryovial tubes and centrifuged (Hospibrand-USA) at 1000 rpm for 5 min to obtain the cell-free supernatant. Sterile pasteur pipettes were used to dispense about 100 μ l of

Table 1. Primers used for the amplification and sequencing of the 16S rRNA in this study

Primer	Sequence (5'-3')
799F	5'-AACACGGATTAGATACC-3'
1193R	5'-ACGTCATCCCCACCTTCC-3'

the cell-free supernatant into the wells. Gentamicin (10 μ g/ml) was used as the positive control for bacterial species, while Ketoconazole (10 μ g/ml) for fungi. The plates were left for some minutes to allow for diffusion of the supernatant; after which they were incubated at 37°C for 24 hr, (for bacteria), and 25°C, for 48 hr (for fungi) ^{17,18}.

DNA extraction and 16S ribosomal RNA amplification

About 5 to 6 colonies of each isolate were harvested into 100 μ l of sterile molecular-grade water. The DNA from the cells was extracted by boiling in a water bath at 95°C set at 10 min ¹⁹. This was then stored in a freezer at -20°C awaiting Polymerase Chain Reaction (PCR).

Polymerase Chain Reaction (PCR)

Two sets of primers were used to run the different PCRs; 799F and 1193R, the primer sequences are stated in table 1.

The reaction was performed for 30 cycles in the PCR Thermal Cycler with initial denaturation at 95°C for 2 min, final denaturation at 94°C for 30 s, annealing of primers at 50°C for 30 s, initial extension at 72°C for 90 s and 10 min for final extension. The amplified product was visualized using an ultraviolet transilluminator. The nucleotide sequences of the 16S rRNA gene of 15 isolates were determined.

Results

Total colony forming units for each dilution

The total colony forming units (CFU/ml) for the 10^{-1} and the 10^{-2} dilutions were 4.0×10^4 and 5.4×10^4 , respectively.

Phenotypic identification of phyllosphere organisms

The physical characteristics of the phyllosphere organisms were determined by considering the form, elevation, appearance, pigmentation, and texture as shown in table 2. The cellular morphology of these organisms was also included in this table. The results of the biochemical test done on the isolates were recorded in table 3. This includes catalase, oxidase, and indole test.

16S ribosomal RNA amplification

Figure 2 shows the agar-gel electrophoresis result of the PCR amplification of the 16S rRNA of the bioactive isolates, while the isolated strains and their accession numbers are presented in table 4.

Antimicrobial activity of phyllosphere organisms

The result of the antimicrobial activity of the me-

Table 2. The Gram reaction, cellular and colonial morphology of phyllospheric organisms of *Funtumia elastic*

S/NO	Form	Elevation	Margin	Appearance	Optical property	Pigmentation	Texture	Gram reaction	Shape
FE-1	Circular	Convex	Entire	Dull	Opaque	Cream	Rough	G+ve	Cocci
FE-3	Irregular	Flat	Undulate	Glistening	Opaque	Cream	Mucoid	G+ve	Rod
FE-4	Circular	Raised	Entire	Shiny	Opaque	White	Mucoid	G-ve	Rod
FE-5a, FE-5b	Rhizoid	Umbonate	Undulate	Wrinkled	Opaque	Dull yellow	Mucoid	G-ve	Cocci
FE-7	Irregular	Umbonate	Undulate	Rough	Opaque	White	Mucoid	G-ve	Rod
FE-9a, FE-9b, FE-10	Circular	Convex	Entire	Rough	Opaque	Cream	Moist	G-ve	Cocci
FE-11	Irregular	Convex	Undulate	Dull	Opaque	White	Moist	G-ve	Cocci
FE-14, FE-15	Circular	Convex	Entire	Rough	Opaque	Cream	Soft	G+ve	Rod
FE-16	Irregular	Umbonate	Undulate	Rough	Opaque	White	Hard	G-ve	Rod
FE-17	Irregular	Flat	Undulate	Rough	Opaque	Yellow	Moist	G+ve	Cocci
FE-18	Circular	Convex	Entire	Shiny	Opaque	Yellow	Mucoid	G-ve	Rod
FE-19, FE-20	Circular	Flat	Entire	Shiny	Opaque	White	Moist	G+ve	Cocci
FE-21, FE-22	Irregular	Flat	Undulate	Dull	Opaque	White	Moist	G-ve	Cocci
FE-23, FE-24	Circular	Crateriform	Entire	Dull	Opaque	White	Mucoid	G-ve	Rod

Table 3. Biochemical tests of isolated phyllosphere organisms

S/NO	Oxidase	Catalase	Indole	MacConkey (colony pigment)	EMB	MSA (colony pigment)
FE-1	-ve	+ve	-ve	Pink	Growth (no gms)	Pink
FE-3	-ve	+ve	-ve	Pink	Growth (no gms)	White
FE-4, FE-5a, FE-5b, FE-7, FE-10, FE-11	-ve	+ve	-ve	Pink	Growth (no gms)	Cream
FE-9a, FE-9b	-ve	-ve	+ve	Cream	Growth with gms	Cream
FE-14	-ve	+ve	-ve	Colourless	Growth (no gms)	Pink
FE-15	-ve	-ve	-ve	Pink	Growth (no gms)	Pink
FE-16	-ve	+ve	-ve	Pink	Growth (no gms)	Yellow
FE-17	-ve	+ve	-ve	Colourless	Growth (no gms)	Pink
FE-18, FE-24	-ve	+ve	-ve	Pink	Growth (no gms)	Orange
FE-19, FE-23	-ve	+ve	-ve	Pink	Growth (no gms)	Cream
FE-20, FE-21	-ve	+ve	-ve	Pink	Growth (no gms)	Pink
FE-22	-ve	+ve	+ve	Pink	Growth (no gms)	Yellow

gms= green metallic sheen.




Figure 2. Agar-gel electrophoresis result of the PCR amplification of the 16S rRNA of the bioactive isolates. Size of DNA marker used = 394 bp.

tabolites from phyllospheric organisms against the test pathogenic bacteria are recorded in table 5, while table 6, contains their antifungal activity.

Discussion

The principle of interspecific antagonism amongst phyllospheric bacteria was applied in this study to determine the antimicrobial activity of metabolites obtained from bacteria isolated from *F. elastica* phyllospheres, against some selected human pathogens.

Twenty-one different colonies were isolated. These colonies observed primarily on TSA plates, ranged from white to cream to yellow and bright yellow in colour, indicating a mixed culture of different bacterial species. Of this number, *Bacillus* species were more prevalent, indicating their dominance of plant phyllo-

Table 4. The BLAST of the sequenced data and the assigned accession numbers

SN	Codes	Identity	Strain	% Identity	Accession number
2	Fe3	<i>Enterobacter hormaechei</i>	BAT2	98.44	OR268892
4	Fe5a	<i>Bacillus aerius</i>	BAT4	100	OR268893
5	Fe5b	<i>Enterobacter hormaechei</i>	BAT5	98.44	OR268894
7	Fe7	<i>Klebsiella pneumonia</i>	BAT7	97.49	OR268895
8	Fe9b	<i>Pantoea dispersa</i>	BAT8	99.72	OR268896
9	Fe10	<i>Enterobacter hormaechei</i>	BAT9	98.75	OR268897
11	Fe14	<i>Bacillus paramycoides</i>	BAT11	100	OR268898
12	Fe15	<i>Bacillus cereus</i>	BAT12	100	OR268899
16	Fe19	<i>Staphylococcus cohnii</i>	BAT16	95.31	OR268900
17	Fe20	<i>Staphylococcus arlettae</i>	BAT17	99.72	OR268901
18	Fe21	<i>Staphylococcus arlettae</i>	BAT18	99.18	OR268902
19	Fe22	<i>Bacillus cereus</i>	BAT19	100	OR268903
20	Fe23	<i>Bacillus aerius</i>	BAT20	99.73	OR268904
21	Fe24	<i>Pantoea stewartii</i>	BAT21	99.18	OR268905

Table 5. Antibacterial activity of the metabolites from phyllosphere organisms against test organisms

S/No	Diameter Zones of inhibition (Mean±SEM)		
	<i>Staphylococcus aureus</i> (ATCC 29213)	<i>Escherichia coli</i> (ATCC 25922)	<i>Klebsiella</i> (ATCC 35659)
FE-1 and FE-3: (<i>Enterobacter hormaechei</i>)	-	-	-
FE-4	-	-	-
FE-5a (<i>Bacillus aerius</i>)	-	-	-
FE-5b (<i>Enterobacter hormaechei</i>)	-	-	-
FE-7 (<i>Klebsiella pneumonia</i>)	-	-	13.0±3.0
FE-9a	18.0±1.0	-	-
FE-9b (<i>Pantoea dispersa</i>)	24.0±1.0	-	26.0±1.0
FE-10 (<i>Enterobacter hormaechei</i>)	20.0±1.0	-	-
FE-11	17.0±1.0	-	14.0±1.0
FE-14 (<i>Bacillus paramycoides</i>)	15.0±1.0	-	-
FE-15, FE-16, FE-17, FE-19, FE-22, FE-23 (<i>Bacillus aerius</i>)	-	-	-
FE-18	20.0±1.0	-	-
FE-20 (<i>Staphylococcus arlettae</i>)	10.0±0.5	-	15.0±1.0
FE-21 (<i>Staphylococcus arlettae</i>)	11.0±0.0	-	15.0±1.0
FE-24 (<i>Pantoea stewartii</i>)	-	23.0±1.0	13.0±1.0
Control (10 µg/ml gentamicin)	12±1.0	10±0.5	11±0.5

SEM = Standard Error of the Mean.

sphere as reported by ^{20,21}. Other Genera include *Enterococcus* sp, *Staphylococcus* sp, *Pantoea* sp, and *Klebsiella* sp, thus aligning our study with that of Ali M, *et al* ²² who reported similar genera in their study. The total colony forming units (CFU/ml) for the 10⁻¹ and the 10⁻² dilutions were 4.0×10⁴ and 5.4×10⁴, respectively, reflecting the reported numerical strength of bacteria in the phyllosphere ⁸.

There were 7 Gram-positives and 14 Gram-negative bacteria present, twelve of which were cocci in shape, while nine were rods (both long and short) (Tables 2 and 3). Earlier studies have shown a predominance of

Gram-negative bacteria over Gram-positive ones on plant phyllosphere ²³. It has been reported that quite a number of phyllospheric organisms are pigmented. This is advantageous to the phyllospheric bacteria as it helps to withstand the adverse effects of ultraviolet rays from the Sun ^{24,25}. In a study conducted by Mazinani *et al* ²¹, 18 bacterial strains were isolated from *Astragalus obtusifolius*, 23 from *Hippocrepis unisiliquosa*, 32 from *Prosopis juliflora* and 31 from *Xanthium strumarium*. From *F. elastica* used in this study, 21 bacterial strains were isolated.

Table 6. Antifungal activity of metabolites from phyllosphere organisms against test organisms

Diameter	Zones of inhibition (Mean±SEM)			
	<i>Candida albicans</i>	<i>Trychophyton rubrum</i>	<i>Meyerozyma guilliermondii</i>	<i>Microsporium canis</i>
S/NO				
FE-1 (<i>Enterobacter hormaechei</i>)	-	31.0±1.0	-	-
FE-3, FE-4, FE-5b, FE-7, FE-9a	-	-	-	-
FE-5a	-	-	-	20.0±1.0
FE-9b (<i>Pantoea dispersa</i>)	-	25.0±0.0	25.0±0.0	29.0±1.0
FE-10 (<i>Enterobacter hormaechei</i>)	-	-	23.0±1.0	30.0±1.0
FE-11, FE-14, FE-15, FE-16, FE-17, FE-18	-	-	-	-
FE-19, FE-20, FE-21, FE-22, FE-23	-	-	-	-
FE-24 (<i>Pantoea stewartii</i>)	19.0±1.0	-	53.5±11.5	48.5±3.5
Control (20 ug/ml-Terbinafine)	15±0.30	22±0.5	30±0.30	27±0.35

SEM = Standard error of the mean.

In this study, not all the phyllosphere bacteria's metabolites showed bioactivity against the test pathogens. Isolate Fe-9b (BAT8), identified as *Pantoea dispersa* had broad-spectrum antimicrobial activity against Gram-positive bacteria, Gram-negative bacterium, and fungal. *Pantoea dispersa* W18, isolated from contaminated soil, was found to exert antimicrobial activity against Mycobacterium species, including *Mycobacterium tuberculosis*, an important human pathogen ²⁶. Fe-24 which was eventually identified as *Pantoea stewartii* BAT21, showed activity against *E. coli* ATCC 25922, *K. pneumoniae* ATCC 35659, and *C. albicans*. Isolates Fe-7 (*K. pneumoniae* BAT7), and Fe-10 (*Enterobacter hormaechei* BAT9), etc, were observed to show activity against at least one of the test bacteria. Based on this result (Table 5), 42% of isolated bacteria showed activity against at least one or more of the test pathogens. Lower activity against the dermatophytes was observed (Table 6). However, some of the isolates (33%) had activity against at least one of the dermatophytes tested. More of the phyllospheric bacteria showed activity against the bacterial pathogens than the fungi. This finding tends to agree with the trend in the study of Goryluk *et al* ²⁰, who also reported low susceptibility of fungal pathogens to inhibitory effects of phyllospheric bacteria. In particular, the most active phyllospheric bacteria were Fe-9b (*Pantoea dispersa* BAT8) and Fe-24 (*Pantoea stewartii* BAT21). However, in the study by Mazinani *et al* ²¹, the most active strains were rods but Gram-positive.

Conclusion

The result of this study shows that the isolated bacteria from *F. elastica* phyllosphere, produced metabolites that are antagonistic (cidal) to some human pathogens, including bacteria and fungi. Therefore, further research is necessary to purify, identify, and characterize these metabolites for possible drug discovery and development against these target pathogens.

Acknowledgement

The authors wish to acknowledge the Technical Staff of the Department of Pharmaceutical Microbiology, University of Ibadan, Nigeria, for their technical input in this study.

Conflict of Interest

Authors declare no conflict of interest.

References

- Mintah SO, Asafo-Agyei T, Archer MA, Junior PA, Boamah D, Kumadoh D, et al. Medicinal plants for treatment of prevalent diseases. *Pharmacognosy-Medicinal Plants* 2019 Jan 17:1-9.
- Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry* 2013;1(6):168-82.
- Ugboko HU, Nwinyi OC, Oranusi SU, Fatoki TH, Omonhinmin CA. Antimicrobial importance of medicinal plants in Nigeria. *ScientificWorldJournal* 2020 Sep 22:2020:7059323.
- Manandhar S, Luitel S, Dahal RK. In vitro antimicrobial activity of some medicinal plants against human pathogenic bacteria. *J Trop Med* 2019 Apr 2:2019:1895340.
- Khameneh B, Iranshahy M, Soheili V, Fazly Bazzaz BS. Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrob Resist Infect Control* 2019 Jul 16:8:118.
- Bringel F, Couée I. Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. *Front Microbiol* 2015 May 22:6:486.
- Vorholt JA. Microbial life in the phyllosphere. *Nat Rev Microbiol* 2012;10(12):828-40.
- Sivakumar N, Sathishkumar R, Selvakumar G, Shyamkumar R, Arjunekumar K. Phyllospheric microbiomes: diversity, ecological significance, and biotechnological applications. *Plant Microbiomes for Sustainable Agriculture* 2020;25:113-72.
- Kumar J, Singh D, Ghosh P, Kumar A. Endophytic and

- epiphytic modes of microbial interactions and benefits. Plant-Microbe Interactions in Agro-Ecological Perspectives 2017;p.227-253.
10. Lindow SE, Brandl MT. Microbiology of the phyllosphere. Apple Environ Microbiol 2003;69(4):1875-83.
 11. Thompson IP, Bailey MJ, Fenlon JS, Fermor TR, Lilley AK, Lynch JM, McCormack PJ, McQuilken MP, Purdy KJ, Rainey PB, Whipps JM. Quantitative and qualitative seasonal changes in the microbial community from the phyllosphere of sugar beet (*Beta vulgaris*). Plant and Soil 1993 Mar;150:177-91.
 12. Clarridge JE 3rd. Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. Clin Microbiol Rev 2004; 17(4):840-62.
 13. Jia T, Yao Y, Wang R, Wu T, Chai B. Dynamics relationship of phyllosphere and rhizosphere bacterial communities during the development of *Bothriochloa ischaemum* in copper tailings. Front Microbiol 2020;11:869.
 14. Zirihi GN, Grellier P, Guédé-Guina F, Bodo B, Mambu L. Isolation, characterization and antiplasmodial activity of steroidal alkaloids from *Funtumia elastica* (Preuss) Stapf. Bioorg Med Chem Lett 2005 May 16;15(10): 2637-40.
 15. Agyare C, Koffuor GA, Boakye YD, Mensah KB. Anti-microbial and anti-inflammatory properties of *Funtumia elastica*. Pharm Biol 2013;51(4):418-25.
 16. Adekunle AA, Ikumapayi AM. Antifungal property and phytochemical screening of the crude extracts of *Funtumia elastica* and *Mallotus oppositifolius*. West Indian Med J 2006;55(4):219-23.
 17. Adeniyi BA, Odelola HA, Oso BA. Antimicrobial potentials of *Diospyros mespiliformis* (Ebenaceae). Afr J Med Med Sci 1996;25(3):221-4.
 18. Adeniyi BA, Onwubuc he BC, Anyiam FM, Ekundayo O, Mahady GB. Anti-Helicobacter pylori activities of *Eucalyptus grandis*: Effects on susceptibility, urease activity and cell surface hydrophobicity. Pharm Biol 2009; 47(1):13-7.
 19. Yashiro E, Spear RN, McManus PS. Culture-dependent and culture-independent assessment of bacteria in the apple phyllosphere. J Appl Microbiol 2011 May 1;110(5): 1284-96.
 20. Goryluk A, Rekosz-Burlaga H, Blaszczyk M. Isolation and characterization of bacterial endophytes of *Chelidonium majus* L. Pol J Microbiol 2009;58(4):355-61.
 21. Mazinani Z, Zamani M, Sardari, S. Isolation and identification of phyllospheric bacteria possessing antimicrobial activity from *Astragalus obtusifolius*, *Prosopis juliflora*, *Xanthium strumarium* and *Hippocrepis unisiliquosa*. Avicenna J Med Biotechnol 2017;9(1):31-7.
 22. Ali M, Walait S, Farhan Ul Haque M, Mukhtar S. Antimicrobial activity of bacteria associated with the rhizosphere and phyllosphere of *Avena fatua* and *Brachiaria reptans*. Environ Sci Pollut Res Int 2021;28:68846-61.
 23. Stoltzfus J, So R, Malarvithi P, Ladha J, Bruijn F. Isolation of endophytic bacteria from rice and assessment of their potential for supplying rice with biologically fixed nitrogen. Plant Soil 1997;194:25-36.
 24. Thapa S, Prasanna R. Prospecting the characteristics and significance of the phyllosphere microbiome. Annals of Microbiology 2018;68:229-45.
 25. Yurimoto H, Shiraishi K, Sakai Y. Physiology of methylo-trophs living in the phyllosphere. Microorganisms 2021;9(4):809.
 26. Teng T, Li X, Zhang L, Li Y. Identification and Characterization of pantocin wh-1 a novel cyclic polypeptide produced by *Pantoea dispersa* W18. Molecules 2020;25 (3):485.